

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email editorial.bmjopen@bmj.com

BMJ Open

Antenatal Care Services and its Implications for Vital and Health Outcomes of Children: Evidence from 171 Surveys in 69 Low- and Middle-Income Countries

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-017122
Article Type:	Research
Date Submitted by the Author:	02-May-2017
Complete List of Authors:	Kuhnt, Jana Vollmer, Sebastian; University of Goettingen, ; Harvard School of Public Health,
Primary Subject Heading:	Global health
Secondary Subject Heading:	Epidemiology
Keywords:	EPIDEMIOLOGY, PERINATOLOGY, NUTRITION & DIETETICS

SCHOLARONE™
Manuscripts

1
2
3 **ANTENATAL CARE SERVICES AND ITS IMPLICATIONS FOR VITAL AND HEALTH OUTCOMES**
4
5 **OF CHILDREN**

6
7 **Evidence from 171 Surveys in 69 Low- and Middle-Income Countries**
8
9

10
11
12
13 **Authors:**

14
15 **Jana Kuhnt**

16 M.Sc., Department of Economics and Centre for Modern Indian Studies, University of Göttingen,
17 Göttingen, Germany
18

19 **Sebastian Vollmer***

20 Prof. Dr., Department of Economics and Centre for Modern Indian Studies, University of Göttingen,
21 Göttingen, Germany
22 Department of Global Health and Population, Harvard T.H. Chan School of Public Health, Boston
23 MA, USA
24

25
26 *** Corresponding Author:**

27 Center for Modern Indian Studies
28 Waldweg 26, Altbau
29 Room 1.117
30 37073 Göttingen
31 Germany
32 Tel. +49 (0)551 / 39-8170
33 Fax +49 (0)551 / 39-20417
34 svollmer@uni-goettingen.de
35
36
37

38 **Acknowledgement**

39
40 The data for this research were collected by The Demographic and Health Surveys Program
41
42 (www.dhsprogram.com), under a contract from the U.S. Agency for International Development.
43
44
45

46 **Manuscript word count: 2928**
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Objectives: Antenatal care (ANC) is an essential part of basic primary health care and its provision has expanded worldwide. There is limited evidence of large-scale cross-country studies on impact of ANC offered to pregnant women on child health outcomes. We investigate the association of ANC in low- and middle-income countries on short- and long-term mortality and nutritional child outcomes.

Setting: Nationally representative health and welfare data from 171 Demographic and Health Surveys (DHS) conducted between Jan 1, 1990, and Dec 31, 2013 from 69 low- and middle-income countries for women of reproductive age (15-49), their children, and respective household.

Participants: Total sample: 1,044,450 children. Observations were lost due to missing data on outcome/exposure variables or covariates. Analytical sample: 751,871 for neonatal mortality, 574,017 for infant mortality, 399,769 for low birthweight, 501,251 for stunting, 512,196 for underweight. Only information from last-born live births considered.

Main Outcomes and Measures: Neonatal and infant mortality, low birth-weight, stunting (low height-for-age) and underweight (low weight-for-age).

Results: Prevalence of neonatal mortality higher among women not receiving any ANC check-up (3.06%) compared to attending at least one check-up (1.67%). For infant mortality the respective numbers are 4.20% and 2.21%. Women not attending ANC on average lower educated and poorer than women attending at least one ANC check-up.

ANC associated with 0.97 pp lower probability of neonatal mortality ($P < .001$), 1.05 pp lower probability of infant mortality ($P < .001$), 3.54 pp lower probability of low birthweight ($P < .004$), 4.29 pp lower probability of stunting ($P < .003$), 3.31 pp lower probability of underweight ($P < .002$).

Conclusions: Currently existing and accessed ANC services in low- and middle-income countries are directly associated with improved birth outcomes and longer-term reductions of child mortality and malnourishment. This emphasizes the importance of ANC within the continuum of maternal/child health care. Quality and quantity of ANC services associated with further improvements of short- and long-term vital and nutritional outcomes.

STRENGTHS AND LIMITATIONS

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

* First study on the association of antenatal care with child health and vital outcomes that for all low- and middle-income countries for which high quality and comparable data are available

* The analysis does not allow a causal interpretation of the results.

For peer review only

INTRODUCTION

Despite strong international efforts to expand the worldwide coverage of basic primary health services for women, pregnancy and childbirth still represent a high-risk period for mother and child, especially in low- and middle-income countries. Nearly three million babies die every year during their first month of life. Especially in low- and middle-income countries many of those deaths and morbidities are due to easily preventable causes. Undetected infections during pregnancy, like malaria, syphilis, tuberculosis, tetanus, or HIV/AIDS, as well as high blood pressure, diabetes, and other pre-existing health conditions often complicate or aggravate a pregnancy and pose significant risk for mother and child. Antenatal care (ANC) – the services offered to mother and unborn child during pregnancy – is an essential part of the basic primary health care offering a mosaic of services that can prevent, detect, and treat such risk factors early on in the pregnancy. The detection of high-risk pregnancies through the analysis of socioeconomic, medical and obstetrical factors represents a key element of antenatal care. It is also often used as a platform for additional interventions that have been shown to positively influence the maternal and child health status, like immunization and nutrition programs, breastfeeding counselling, or to educate women about the possibilities of family planning and birth spacing.¹⁻⁵ Additionally, antenatal care programs are used to provide care and information that is not directly related to pregnancy but can reduce possible maternal risk factors, like promoting healthy lifestyles, to tackle malnutrition or inform about gender-based violence. Hence, antenatal care is a potentially important determinant in reducing maternal and child morbidity and mortality.⁶⁻¹³ Within the last decades, the provision of antenatal care services has increased worldwide and during 2006-2013 the antenatal care coverage defined as at least one antenatal care visit with a skilled provider was around 75% in low- and lower-middle income countries.^{14,15} To our knowledge, there exists no global study for all low- and middle-income countries, which analyses the association of existing antenatal care services that are offered to pregnant women in low- and middle-income countries on child health outcomes.

Numerous studies have helped to develop an internationally accepted set of so-called, essential antenatal care services' by evaluating the effects of single interventions, like tetanus and malaria prevention programs, on maternal and neonatal health¹⁶⁻¹⁸ or by studying the optimal number and content of antenatal care visits.^{19,20} But the de facto offered and utilized set of antenatal care services

1
2
3 can deviate greatly from those recommended interventions across as well as within countries. A
4
5 couple of studies evaluate the relationship between the utilization of antenatal care services and
6
7 perinatal outcomes in individual low- and middle-income countries. The majority have shown
8
9 positive effects of antenatal care on newborn mortality, the occurrence of stillbirth and pre-term
10
11 labour, and low birthweight.²¹⁻²⁶ However, they exclusively focus on single countries, are often
12
13 conducted at the clinic level, and have small sample sizes. This limits their external validity. We
14
15 identified only one study that focuses on a larger regional sample. Conde-Agudelo et al. (2000)
16
17 studied 837,232 births in Latin America between 1985 and 1997.²⁷ One major risk factor associated
18
19 with fetal death was the lack of antenatal care. We could not find a study that took into account
20
21 possible long-term effects of the utilization of antenatal care services on children's nutritional and
22
23 vital status.

24
25 With up to 171 Demographic and Health Surveys (DHS) from up to 69 low- and middle-income
26
27 countries we use the most comprehensive data for low- and middle-income countries in the world
28
29 that currently exist. Specifically, we investigate whether the attendance of mothers at antenatal care
30
31 services was associated with improved short- and long-term survival rates or reductions in the
32
33 prevalence of low birthweight, stunting and underweight in their children.

34 35 36 37 38 METHODS

39 40 **Data**

41
42 We used data from the Demographic and Health Surveys (DHS), which are cross-sectional household
43
44 surveys that use a harmonized questionnaire to facilitate between-country comparisons. The DHS
45
46 collect nationally representative health and welfare data for women of reproductive age, age 15-49,
47
48 their children and the respective household.

49
50 They have been conducted at different time intervals in 90 low- and middle-income countries since
51
52 1985. We included all surveys, which have information for the relevant outcome and explanatory
53
54 variables. The final sample consists of pooled data from up to 171 surveys in 69 low- and middle-
55
56 income countries worldwide conducted between Jan 1, 1990, and Dec 31, 2013. The DHS used a
57
58 multistage stratified sampling, countries were divided into regions, within which populations were
59
60

1
2
3 stratified by area of residence and from which a random sample was taken according to the most
4 recent population census. Those are the primary sampling units (clusters with on average 250
5 households), which are equally likely to be selected to the proportion the specific cluster's population
6 contributed to the total population. At a second stage, after a complete listing of all households
7 within the cluster, an average of 20-30 households was randomly selected by equal probability.
8 Sampling weights in order to calculate nationally representative statistics are provided by DHS.
9 For each live birth the women were among other things asked about date of birth, birthweight, vital
10 status at the time of the interview, and either current age or age of death of the child. Furthermore,
11 the DHS collected information on the height and weight of the children born during the last five
12 years. For each last-born live birth there is information on the attendance rates and quality of
13 antenatal care visits during the last pregnancy that led to a live birth. Considering the on-going
14 debates on the importance of the quality and number of antenatal care visits^{20,28-30} we specified three
15 different main explanatory variables, including the mere attendance of antenatal care (irrespective of
16 number and quality of the visits), the provision of service by a skilled attendant, and the attendance
17 of at least four ANC visits as recommended by the WHO. Antenatal care visits with a doctor,
18 midwife, nurse, auxiliary midwife, obstetrician, health professional, or trained (traditional) birth
19 attendant, we considered as skilled antenatal care services. Whereas antenatal care with a traditional
20 birth attendant, relatives, any other person or none of the mentioned was classified as unskilled
21 antenatal care.

22 Outcomes

23 We analysed the data for short- and long-term vital outcomes and low birthweight of all last-born
24 live births as well as stunting and underweight for the last-born children aged 0-59 months at the
25 time of the interview. Mortality outcomes were neonatal death, defined as death within the first
26 month of life and infant death, defined as death after the first month but within the first year of life.
27 Nutritional outcomes were low birthweight, stunting and underweight. We used the WHO
28 classification that defines low birthweight as a birthweight below 2,500g at birth. Following WHO
29 and UNICEF suggestions, we only included biologically plausible birthweights from 500 – 5999g. To
30 calculate stunting and underweight, we used anthropometric data defined by WHO standards and
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 classifications. Comparing the child's height and weight to those of a well-nourished reference
4
5 population of the same age and sex, allows us to calculate the z-scores of height-for-age and weight-
6
7 for-age. Stunting is defined by a height-for-age z-score of less than -2 and underweight is defined by
8
9 a weight-for-age z-score of less than -2. Biologically implausible values of the z-scores were excluded
10
11 following WHO guidelines.

12 13 14 15 **Statistical Analysis**

16 We used linear probability regression models to investigate the association between antenatal care
17
18 services and short- and longer-term vital and health outcomes of children. We adjusted the
19
20 regressions for confounding factors and controlled for PSU fixed effects. Standard errors were
21
22 clustered at the PSU level. They capture characteristics of local enumeration areas that are common
23
24 to all respondents from that area. We used sex, birth order, birth spacing, birth month, and whether
25
26 the child was a multiple birth; the mother's age at birth, education, work status, relation to the
27
28 household head, and her marital status; and household wealth quintile as covariates. The wealth
29
30 quintile variable is based on the ownership of household assets and indicates the household's wealth
31
32 relative to other households within the respective country in that survey. Additionally, by including
33
34 variables indicating place of delivery, mode of delivery (vaginal or caesarean), status of tetanus
35
36 injection of mother before birth, and if she breastfed at least one month after birth (only applicable
37
38 for long-term outcomes), we inspected possible mediator variables meaning that the uptake of
39
40 antenatal care services might starkly influence those variables, which themselves might affect the
41
42 outcome variables.

43
44 Using Stata (version 14.0) for all statistical analyses, we also took into account the stratified survey
45
46 design by using the Stata svy command. We used sampling weights provided by DHS in all our
47
48 regressions.

49 50 51 52 **RESULTS**

53
54 The total sample included observations for 1,044,450 children. Observations were lost due to missing
55
56 data on outcome variables, missing data on the ANC variables (including the dummies indicating the
57
58 mere attendance of ANC visits, the attendance at skilled providers, and whether at least four visits
59
60

1
2
3 were accomplished during the pregnancy) or missing data on covariates. The final analytical sample
4 is 751,871 for neonatal mortality, 399,769 for low birthweight, 574,017 for infant mortality, 501,251 for
5 stunting and 512,196 for underweight (see Figure 1).
6
7

8
9
10 The prevalence of newborn death was higher among women that did not receive any antenatal care
11 check-up (3.06 percent) compared to those attending at least one check-up (1.67 percent). For infant
12 mortality the respective numbers are 4.20 percent and 2.21 percent. Prevalence of all outcome
13 variables was higher among women not attending any antenatal care visit than those attending at
14 least one antenatal check-up (Table 1). Pregnant women that did not attend antenatal care visits
15 were on average lower educated and poorer than those women that attended at least one ANC
16 check-up (see Table A1 in the appendix).
17
18
19
20
21
22
23

24
25
26 In Table 2 we report the association between antenatal care take-up and short- and long-term
27 mortality outcomes. For each outcome, we show the results from five different specifications where
28 PSU fixed effects are included in all five. The first column shows the mere association between the
29 attendance of at least one ANC visit without controlling for any covariates. The second column
30 shows the association adjusted for control variables and the third column shows the association that
31 in addition to control variables is also adjusted for potential transmission channels of antenatal care
32 (place of delivery, mode of delivery, status of tetanus toxoid (TT) injection of mother before birth,
33 and for the long-term outcomes we also include if the mother breastfed at least for one month after
34 birth). The fourth column reports the coefficients while adding whether the mother has attended her
35 antenatal care visits at a skilled provider and if she received at least four visits before the delivery.
36 The last column includes potential transmission channels. We will focus on the second and fourth
37 specification adjusted for control variables and only refer to the other three specifications for
38 comparison purposes.
39
40
41
42
43
44
45
46
47
48
49
50
51

52 Women attending at least one antenatal care visit have a 0.98 percentage points reduced probability
53 of their newborn dying within the first month after birth and a 1.05 percentage points lower
54 probability of experiencing death of their child within the first year of life. While the quality of the
55 provider is not significantly related to neonatal mortality, the quantity is. Having at least four ANC
56
57
58
59
60

1
2
3 visits is associated with 0.54 percentage points fewer neonatal deaths. The association between
4
5 antenatal care check-ups and infant mortality, seems to be driven by the quality and quantity of
6
7 prenatal visits. Once controlling for the quality and number of visits, the coefficient for at least one
8
9 ANC visit is not statistically significant anymore. Going to a skilled provider is associated with 0.69
10
11 percentage points fewer infant deaths. Additionally, having at least four ANC check-ups, is
12
13 associated with 0.48 percentage points fewer infant deaths.

14
15 The DHS dataset also provides information on several variables that are well established in the
16
17 literature to impact mortality and morbidity outcomes of children and that simultaneously are
18
19 potentially influenced by ANC attendance. When controlling for these potential transmission
20
21 channels of antenatal care services, it can be seen that some of the ANC coefficients are somewhat
22
23 attenuated when controlling for these additional variables but not by much. Results for all covariates
24
25 are provided in the appendix (Table A2).

26
27
28 In Table 3 we report the association between antenatal care and short- and long-term nutritional
29
30 outcomes of the child. If the mother chooses to have at least one ANC visit, this is associated with a
31
32 3.54 percentage points reduced probability of giving birth to a low birthweight baby. Stunting and
33
34 underweight outcomes are reduced by 4.29 and 3.31 percentage points respectively. The association
35
36 between antenatal care check-ups and low birthweight as well as underweight, seem also to be
37
38 driven by the quality and quantity of prenatal visits. Once controlling for the quality and number of
39
40 visits, the coefficient for at least one ANC visit loses its significance. Attendance at a skilled provider
41
42 reduces the probability of having a low birthweight baby by 1.37 percentage points and for
43
44 underweight by 2.18 percentage points. Additionally having at least four ANC check-ups, reduces the
45
46 outcomes by 2.76 percentage points and 2.06 percentage points respectively. In case of stunting the
47
48 coefficient indicating at least one ANC visit remains significant when adding quality and quantity
49
50 indicators. Going to a skilled provider is associated with 2.17 percentage points reduced stunting
51
52 probability and attending at least four ANC check-ups reduces stunting probability by 1.57
53
54 percentage points.

1
2
3 Adding potential transmission channels of antenatal care services to the regression slightly
4 attenuates the three ANC coefficients (more so in the case of LBW). Results for all covariates are
5 provided in the appendix (Table A2).
6
7
8
9

10 DISCUSSION

11
12 Most existing evidence on the effect of antenatal care on child health is based on data from high-
13 income countries and their conclusions are not easily transferable to low- and middle-income
14 settings. The existing studies for low- and middle-income countries often focus on individual
15 countries. Furthermore, the studied effects of antenatal care have been limited to direct short-term
16 maternal and child delivery outcomes. This is the first large-scale cross-country study for all low-and
17 middle-income countries with availability of comparable data for ANC, which systematically
18 investigates the association of ANC with short- and long-term mortality and nutritional child
19 outcomes.
20
21
22
23
24
25
26
27

28 Using child vital data and child anthropometry from up to 171 surveys in 69 low- and middle-income
29 countries we have shown that antenatal care is associated with reductions in neonatal and infant
30 mortality, low birthweight, stunting and underweight. This association can be seen for all outcomes
31 in almost all world regions though it is especially strong in Latin America and Caribbean, Sub-
32 Saharan Africa, and South and Southeast Asia (which constitute about 90% of our sample). Receiving
33 prenatal care by skilled providers is significantly associated with prevalence reductions of all
34 outcome variables (except for neonatal mortality) and hence, plays an important role in the provision
35 of ANC services. The same is true for the quantity of antenatal care visits during pregnancy.
36
37
38
39
40
41
42
43

44 Receiving at least four ANC check-ups, improves the outcomes of all short- and long-term child
45 health indicators. The magnitude of the association is quantitatively important, as it varies around -
46 0.98 and -4.29 percentage points. Currently, 47 children out of 1000 livebirths die in the developing
47 world before their fifth birthday.³¹
48
49
50
51

52 Many pregnant women in low- and middle-income countries have no access or do not attend ANC
53 services regularly enough (4+ visits) or at skilled providers.^{14,32} According to our results, improving the
54 coverage and uptake of antenatal care services could be an important tool to improve short- and
55 even long-term mortality and nutritional outcomes of children.
56
57
58
59
60

1
2
3 There are a couple of self-selection issues and limitations, which we have attempted to address. For
4 instance, we controlled for mother's education and household wealth, since more educated or more
5 affluent mothers might be more likely to seek antenatal care and at the same time have a better
6 overall health status. Similarly, we adjusted for PSU fixed effects to control for community
7 characteristics, overall health status in the region and the local availability and quality of health care
8 services as well as other characteristics that are common to the local area. However, there are a few
9 maternal characteristics, which we did not observe and therefore were not able to control for. For
10 instance, if pregnant women feel that there could be something wrong, they might be more likely to
11 seek antenatal care. Similarly, if women had negative birth outcomes in the past, they might also be
12 more likely to seek antenatal care to avoid the repetition of the negative birth outcome. Both cases
13 would downward bias our estimates, and the true association would be even stronger than the
14 association, which we found in our analysis. It is important to point out again that we are only
15 including the outcomes of livebirths. The attendance at antenatal care services might lead to better
16 survival chances of those babies that would have otherwise died before birth. This might impose a
17 downward bias on our estimates. However, there are also potential selection issues, which could bias
18 the results in the other direction. For instance it is unclear how women would behave in case of an
19 unwanted pregnancy, they might be less likely to seek antenatal care and have worse overall health
20 behaviour compared to women in planned pregnancies. In a robustness check we controlled for an
21 indicator variable if the pregnancy was wanted and this did not change the results.
22
23 In summary, our study provides evidence for the potential importance of antenatal care for
24 improving child health and vital outcomes in low- and middle-income countries.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 A. CONTRIBUTOR SHIP STATEMENT

4 JK and SV conceptualized the study and developed the analytical strategy. JK wrote the first draft of
5 the paper, conducted the statistical analysis and interpreted the data. SV contributed to the
6 interpretation of the results and critically revised it.
7

8 B. COMPETING INTERESTS

9 The authors have read and understood BMJ policy on declaration of interests and declare that they
10 have no competing interests.
11

12 C. FUNDING

13 This research received no specific grant from any funding agency in the public, commercial or not-
14 for-profit sectors.
15

16 D. DATA SHARING STATEMENT

17 This study used data that was collected by the Demographic and Health Surveys Program
18 (www.dhsprogram.com), under a contract from the U.S. Agency for International Development.
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

REFERENCES

1. Imdad A, Bhutta Z. Effects of calcium supplementation during pregnancy on maternal, fetal and birth outcomes. *Paediatric and perinatal epidemiology*. 2012;26(5):138-152.
2. Gupta A, Sharma D. Breastfeeding and Child Health. *Economic and Political Weekly*. 2006;April 15.
3. Oddy W, Kendall G, Li J, et al. The long-term effects of breastfeeding on child and adolescent mental health: a pregnancy cohort study followed for 14 years. *Journal of Pediatrics*. 2010;156(4):568-574.
4. Black R, Victora C, Walker S, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *The Lancet*. 2013;382:427-51.
5. Titalay C, Dibley M. Antenatal iron/folic acid supplements, but not postnatal care, prevents neonatal deaths in Indonesia: analysis of Indonesia Demographic and Health Surveys. *BMJ Open*. 2012;2(6).
6. Abou-Zahr C, Wardlaw T. Antenatal care in developing countries: promises, achievements and missed opportunities. Geneva: World Health Organization; 2003.
7. Chen XK, Wen SW, Yang Q, Walker M. Adequacy of prenatal care and neonatal mortality in infants born to mothers with and without antenatal high-risk conditions. *Australian and New Zealand Journal of Obstetrics and Gynaecology*. 2007;47(2): 122-7.
8. Khan K, Wojdyla D, Say L, Gulmezoglu A, VanLook P. WHO analysis of causes of maternal death: a systematic review. *The Lancet*. 2006;367:1066-1074.
9. Lincetto O, Mothebesoane-Anoh S, Gomez P, Munjanja S. Antenatal Care. In Opportunities for Africa's Newborns. PMNCH; 2006.
10. World Health Organization. WHO Statement on Antenatal Care. World Health Organization: Geneva, Switzerland; 2011.
11. Zanconato G, Msolomba R, Guarenti L, Franchi M. Antenatal care in developing countries: the need for a tailored model. *Seminars in Fetal and neonatal Medicine*. 2006;11(1):15-20.

12. PMNCH. The PMNCH Report 2012. Analysing Progress on Commitments to the Global Strategy for Women's and Children's Health. World Health Organization: Geneva, Switzerland; 2012.
13. Moss W, Darmstadt G, Marsh D, Black R, Santosham M. Research Priorities for the Reduction of Perinatal and Neonatal Morbidity and Mortality in Developing Country Communities. *Journal of Perinatology*. 2002;22(6): 484-495.
14. World Health Organization. World Health Statistics. World Health Organization: Geneva, Switzerland; 2014.
15. Wang W, Alva S, Wang S, Fort A. Levels and Trends in the Use of Maternal Health Services in Developing Countries. DHS Comparative Reports No. 26. Calverton, Maryland, USA: ICF Macro; 2011.
16. Bhutta Z, Das J, Bahl R, et al. Can available interventions end preventable deaths in mothers, newborn babies, and stillbirths, and at what cost? *The Lancet*. 2011;384(9940):347-370.
17. Blencowe H, Lawn J, Vandelaer J, Roper M, Cousens S. Tetanus toxoid immunization to reduce mortality from neonatal tetanus. *International Journal of Epidemiology*. 2010;39(suppl 1): i102-i109.
18. Blencowe H, Cousens S, Kamb M, Berman S, Lawn J. Lives Saved Tool supplement detection and treatment of syphilis in pregnancy to reduce syphilis related stillbirths and neonatal mortality. *BMC Public Health*. 2011;11(Suppl 3).
19. Villar J, Ba'aqel H, Piaggio G, et al. WHO antenatal care randomised trial for the evaluation of a new model of routine antenatal care. *The Lancet*. 2001;357:1551-1564.
20. Carroli G, Villar J, Piaggio G, et al. WHO systematic review of randomized controlled trials of routine antenatal care. *The Lancet*. 2001;357:1565-70.
21. Hong R, Ruiz-Beltran M. Impact of Prenatal Care on Infant Survival in Bangladesh. *Maternal and Child Health Journal*. 2007;11,199-206.
22. Kapoor S, Anand K, Kumar G. Risk Factors for Stillbirths in A Secondary Level Hospital at Ballabgarh, Haryana: A Case Control Study. *Indian Journal of Pediatrics*. 1994;61:161-66.

- 1
2
3 23. Mbuagbaw L, Gofin R. A new measurement for optimal antenatal care: determinants and
4 outcomes in Cameroon. *Maternal and Child Health Journal*. 2011;15(8):1427-34.
5
6
7 24. Brown C, Sohani S, Khan K, Lilford R, Mukhwana W. Antenatal care and perinatal outcomes
8 in Kwale district, Kenya. *BMC Pregnancy and Childbirth*. 2008;8(2).
9
10
11 25. Celik Y, Younis M. Effects of antenatal care services on birthweight: importance of model
12 specification and empirical procedure used in estimating the marginal productivity of health
13 inputs. *Journal of Medical Systems*. 2007;31(3):197-204.
14
15
16 26. McCaw Binns A, Greenwood R, Ashley D, Golding J. Antenatal and Perinatal-Care in Jamaica
17 - do They Reduce Perinatal Death Rates? *Paediatric and Perinatal Epidemiology*.
18 1994;8(Suppl 1):86-97.
19
20
21 27. Conde-Agudelo A, Belizán J, Díaz-Rossello J. Epidemiology of fetal death in Latin America.
22 *Acta Obstetrica et Gynecologica Scandinavica*. 2000;79:371-378.
23
24
25 28. Vogel P, Habib N, Souza J, et al. Antenatal care packages with reduced visits and perinatal
26 mortality: a secondary analysis of the WHO Antenatal Care Trial. *Reproductive Health*.
27 2013;10(19).
28
29
30 29. Hofmeyr GJ, Hodnett ED. Antenatal care packages with reduced visits and perinatal
31 mortality: a secondary analysis of the WHO Antenatal Care Trial – Comentary: routine
32 antenatal visits for healthy pregnant women do make a difference. *Reproductive Health*.
33 2013;10(20).
34
35
36 30. Hodgins S, D'Agostino A. The quality-coverage gap in antenatal care: toward better
37 measurement of effective coverage. *Global Health: Science and Practice*. 2014.
38
39
40 31. You D, Hug L, Ejdmeyr S, Beise J. Levels and Trends in Child Mortality. Report 2015. United
41 Nations Children's Fund: New York, USA; 2015.
42
43
44 32. World Health Organization. World Health Statistics. World Health Organization: Geneva,
45 Switzerland; 2016.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Descriptive Statistics

Outcome variable	Mother attended at least 1 ANC visit; % (n=615621)	Mother did not attend ANC visit, % (n=136250)
Neonatal Mortality	1.67	3.06*
Infant Mortality	2.21	4.20*
Low Birthweight	9.69	14.24*
Stunting	31.37	47.11*
Underweight	15.72	30.73*

n values denote the number of observations in the neonatal mortality sample.
* Proportions in the two groups were significantly different from each other, $p \leq 0.05$ (t test).

Table 2: Associations between antenatal care visits and mortality outcomes

Neonatal Mortality					
	(1)	(2)	(3)	(4)	(5)
at least 1 ANC visit	-0.0101*** [0.001]	-0.00977*** [0.001]	-0.00468*** [0.001]	-0.00567*** [0.002]	-0.00376** [0.002]
skilled ANC provider				-0.00162 [0.002]	0.00151 [0.002]
at least 4 ANC visits				-0.00537*** [0.001]	-0.00532*** [0.001]
facility delivery			0.00193*** [0.001]		0.00244*** [0.001]
mother had TT injct			-0.00822*** [0.001]		-0.00792*** [0.001]
delivery by C-section			0.0102*** [0.001]		0.0103*** [0.001]
N	751871	751871	688612	751871	688612
Infant Mortality					
	(1)	(2)	(3)	(4)	(5)
at least 1 ANC visit	-0.0126*** [0.001]	-0.0105*** [0.001]	-0.00614*** [0.001]	-0.00160 [0.003]	-0.000395 [0.003]
skilled ANC provider				-0.00690*** [0.002]	-0.00423* [0.002]
at least 4 ANC visits				-0.00475*** [0.001]	-0.00400*** [0.001]
facility delivery			-0.00185** [0.001]		-0.00138* [0.001]
mother had TT injct			-0.00541*** [0.001]		-0.00503*** [0.001]
delivery by C-section			0.00127 [0.001]		0.00140* [0.001]
child was breastfed			-0.0498*** [0.002]		-0.0497*** [0.002]
N	574017	574017	522419	574017	522419
Controls	no	yes	yes	yes	yes
All regressions include PSU fixed effects. ***, ** and * denote significance at the 1%, 5%, and 10% levels, respectively. Std. Errors in square brackets and are clustered at PSU level. Control variables include: Mother's age, marital status and educational achievement, whether she heads the HH, child's sex and birth order and spacing, month of birth, whether it was a multiple birth and HH wealth quintile.					

Table 3: Associations between antenatal care visits and nutritional outcomes

Low Birthweight					
	(1)	(2)	(3)	(4)	(5)
at least 1 ANC visit	-0.0403*** [0.004]	-0.0354*** [0.004]	-0.0323*** [0.004]	-0.00249 [0.009]	0.00166 [0.009]
skilled ANC provider				-0.0137* [0.008]	-0.0158* [0.008]
at least 4 ANC visits				-0.0276*** [0.002]	-0.0277*** [0.002]
facility delivery			0.00181 [0.002]		0.00349 [0.002]
mother had TT injct			-0.00646*** [0.002]		-0.00457** [0.002]
delivery by C-section			0.0204*** [0.002]		0.0209*** [0.002]
N	399769	399769	367490	399769	367490
Stunting					
	(1)	(2)	(3)	(4)	(5)
at least 1 ANC visit	-0.0693*** [0.003]	-0.0429*** [0.003]	-0.0375*** [0.003]	-0.0145** [0.007]	-0.0155** [0.008]
skilled ANC provider				-0.0217*** [0.007]	-0.0172** [0.007]
at least 4 ANC visits				-0.0157*** [0.002]	-0.0128*** [0.002]
facility delivery			-0.0363*** [0.002]		-0.0347*** [0.002]
mother had TT injection			0.00185 [0.002]		0.00277 [0.002]
delivery by C-section			0.00441* [0.003]		0.00483* [0.003]
child was breastfed			-0.0282*** [0.005]		-0.0282*** [0.005]
child ever got vaccine			0.0544*** [0.004]		0.0550*** [0.004]
N	501251	501251	457622	501251	457622
Underweight					
	(1)	(2)	(3)	(4)	(5)
at least 1 ANC visit	-0.0505*** [0.002]	-0.0331*** [0.002]	-0.0226*** [0.002]	-0.00225 [0.006]	-0.00224 [0.007]
skilled ANC provider				-0.0218*** [0.006]	-0.0136** [0.006]
at least 4 ANC visits				-0.0206*** [0.002]	-0.0168*** [0.002]
facility delivery			-0.0252*** [0.002]		-0.0232*** [0.002]
mother had TT injct			-0.00443** [0.002]		-0.00338* [0.002]
delivery by C-section			-0.00569*** [0.002]		-0.00515*** [[0.002]
child was breastfed			-0.0164*** [0.004]		-0.0163*** [0.004]

child ever got vaccine			-0.00336 [0.003]		-0.00269 [0.003]
N	512196	512196	467001	512196	467001
Controls	no	yes	yes	yes	yes
All regressions include PSU fixed effects. ***, ** and * denote significance at the 1%, 5%, and 10% levels, respectively. Std. Errors in square brackets and are clustered at PSU level. Control variables include: Mother's age, marital status and educational achievement, whether she heads the HH, child's sex and birth order and spacing, month of birth, whether it was a multiple birth and HH wealth quintile.					

For peer review only

Table A1: Covariates across exposure category

Covariate	Mother attended at least 1 ANC visit; % (n=615621)	Mother did not attend ANC visit, % (n=136250)
Maternal education		
<i>None</i>	45.40	82.97*
<i>Primary or incompl. Secondary</i>	33.91	13.33*
<i>Secondary or higher</i>	20.69	3.70*
Maternal age at birth		
<i>≤17 years</i>	3.51	4.90*
<i>17 - 19 years</i>	12.87	12.51*
<i>20 - 24 years</i>	28.23	25.15*
<i>25 - 29 years</i>	24.53	23.08*
<i>30 years or older</i>	30.86	34.35*
Mother is working	57.21	50.97*
Mother is HH Head	9.13	7.24*
Mother is married	72.25	85.79*
Child is female	48.71	48.74
Multiple birth	0.88	0.98*
Birth order		
<i>First child</i>	31.24	19.37*
<i>Second child</i>	21.60	16.19*
<i>Third child</i>	15.57	14.58*
<i>Fourth child</i>	10.41	12.64*
<i>Fifth or later born child</i>	21.18	37.23*
Birth spacing ≤ 18 mths	3.55	5.57*
Wealth quintiles		
<i>First quintile</i>	15.16	32.46*
<i>Second quintile</i>	17.54	26.84*
<i>Third quintile</i>	19.89	20.42
<i>Fourth quintile</i>	22.32	13.85*
<i>Fifth quintile</i>	25.10	6.42*
N values denote the number of observations in the neonatal mortality sample. Values were population weighted. * Proportions in the two groups were significantly different from each other, $p \leq 0.05$ (t test).		

Table A2: Associations between the exposure variables and full set of control variables

	Neonatal Mortality	Infant Mortality	Low Birthweight	Stunting	Underweight
Maternal education [reference: no education]					
Primary or incompl. Secondary	-0.00247*** [0.001]	-0.00544*** [0.001]	-0.0183*** [0.002]	-0.0448*** [0.002]	-0.0363*** [0.002]
Secondary or higher	-0.00516*** [0.001]	-0.00916*** [0.001]	-0.0346*** [0.002]	-0.0778*** [0.003]	-0.0453*** [0.002]
Maternal age at birth [reference: ≤					
17 - 19 years	-0.00504*** [0.001]	-0.00684*** [0.002]	-0.00847** [0.004]	-0.0503*** [0.005]	-0.0352*** [0.004]
20 - 24 years	-0.00781*** [0.001]	-0.0104*** [0.002]	-0.0211*** [0.004]	-0.0754*** [0.004]	-0.0499*** [0.004]
25 - 29 years	-0.00697*** [0.001]	-0.00990*** [0.002]	-0.0214*** [0.004]	-0.101*** [0.005]	-0.0576*** [0.004]
30 years or older	-0.000944 [0.001]	-0.00903*** [0.002]	-0.00921** [0.004]	-0.118*** [0.005]	-0.0597*** [0.004]
Mother is working	0.00333*** [0.000]	0.00144** [0.001]	-0.000679 [0.001]	0.0169*** [0.002]	0.00674*** [0.002]
Mother is HH Head	-0.000707 [0.001]	-0.000768 [0.001]	-0.000502 [0.002]	0.00555** [0.003]	-0.00265 [0.002]
Mother is married	-0.00172*** [0.001]	-0.00607*** [0.001]	-0.00955*** [0.002]	-0.00749*** [0.002]	-0.00868*** [0.002]
Child is female	-0.00453*** [0.000]	-0.00191*** [0.001]	0.0200*** [0.001]	-0.0394*** [0.002]	-0.0223*** [0.001]
Multiple birth	0.100*** [0.004]	0.0649*** [0.005]	0.384*** [0.010]	0.144*** [0.009]	0.146*** [0.008]
Birth order [reference: First born child]					
Second child	-0.00888*** [0.001]	-0.00455*** [0.001]	-0.0275*** [0.002]	0.0290*** [0.002]	0.00553*** [0.002]
Third child	-0.0104*** [0.001]	-0.00487*** [0.001]	-0.0324*** [0.002]	0.0462*** [0.003]	0.0149*** [0.002]
Fourth child	-0.0113*** [0.001]	-0.00538*** [0.001]	-0.0365*** [0.003]	0.0582*** [0.003]	0.0202*** [0.003]
Fifth or later born child	-0.0108*** [0.001]	-0.00353*** [0.001]	-0.0396*** [0.003]	0.0837*** [0.003]	0.0294*** [0.003]
Birth spacing ≤ 18 months	0.0199*** [0.001]	0.0228*** [0.002]	0.0228*** [0.004]	0.0491*** [0.004]	0.0308*** [0.003]
Wealth quintiles [reference: poorest 20%]					
Second quintile	-0.000750 [0.001]	-0.00202** [0.001]	-0.00730*** [0.002]	-0.0173*** [0.003]	-0.0123*** [0.002]
Third quintile	-0.0000752 [0.001]	-0.00254** [0.001]	-0.0123*** [0.003]	-0.0367*** [0.003]	-0.0308*** [0.002]

<i>Fourth quintile</i>	-0.0000543 [0.001]	-0.00561*** [0.001]	-0.0173*** [0.003]	-0.0586*** [0.003]	-0.0501*** [0.003]
<i>Fifth quintile</i>	-0.00113 [0.001]	-0.00653*** [0.001]	-0.0216*** [0.003]	-0.0988*** [0.003]	-0.0766*** [0.003]
Month of birth	0.00000989 [0.000]	-0.0000732 [0.000]	-0.000215 [0.000]	-0.00375*** [0.000]	-0.00294*** [0.000]
N	751871	574017	399769	501251	512196
Results are from the main results regressions (4). ***, ** and * denote significance at the 1%, 5%, and 10% levels, respectively. Std. Errors in square brackets and are clustered at PSU level.					

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Neonatal Mortality	Low Birthweight	Infant Mortality	Stunting	Underweight
1044450	1044450	1044450	1044450	1044450
39394 missing data for control variables	89394 missing data for control variables			
955056	955056	955056	955056	955056
96096 missing data for ANC variables	196096 missing data for ANC variables	196096 missing data for ANC variables	196096 missing data for ANC variables	196096 missing data for ANC variables
758960	758960	758960	758960	758960
7089 missing data for outcome variable	359191 missing data for outcome variable	184943 missing data for outcome variable	257709 missing data for outcome variable	246764 missing data for outcome variable
751871	399769	574017	501251	512196
69 LMIC 171 surveys	68 LMIC 161 surveys	69 LMIC 171 surveys	64 LMIC 153 surveys	64 LMIC 152 surveys

Figure 1: Sample deduction

297x209mm (300 x 300 DPI)

ew only

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Page #
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2-3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4-5
Objectives	3	State specific objectives, including any prespecified hypotheses	5
Methods			
Study design	4	Present key elements of study design early in the paper	6
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	6
Participants	6	(a) <i>Cohort study</i> —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up <i>Case-control study</i> —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls <i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	6
		(b) <i>Cohort study</i> —For matched studies, give matching criteria and number of exposed and unexposed <i>Case-control study</i> —For matched studies, give matching criteria and the number of controls per case	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	6-7
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	6
Bias	9	Describe any efforts to address potential sources of bias	7-8
Study size	10	Explain how the study size was arrived at	8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	7-8
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	7-8
		(b) Describe any methods used to examine subgroups and interactions	8
		(c) Explain how missing data were addressed	6,8
		(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed <i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	8
		(e) Describe any sensitivity analyses	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Continued on next page

For peer review only

Results			Page #
Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram	8 Figure 1
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest (c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	9 (table A1)
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time <i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure <i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	9
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	9-10
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	9-10
Discussion			
Key results	18	Summarise key results with reference to study objectives	11
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	12-13
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	12-13
Generalisability	21	Discuss the generalisability (external validity) of the study results	
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	14

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

BMJ Open

Antenatal Care Services and its Implications for Vital and Health Outcomes of Children: Evidence from 193 Surveys in 69 Low- and Middle-Income Countries

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-017122.R1
Article Type:	Research
Date Submitted by the Author:	15-Aug-2017
Complete List of Authors:	Kuhnt, Jana Vollmer, Sebastian; University of Goettingen, ; Harvard School of Public Health,
Primary Subject Heading:	Global health
Secondary Subject Heading:	Epidemiology
Keywords:	EPIDEMIOLGY, PERINATOLOGY, NUTRITION & DIETETICS

SCHOLARONE™
Manuscripts

1
2
3 **ANTENATAL CARE SERVICES AND ITS IMPLICATIONS FOR VITAL AND HEALTH OUTCOMES**
4
5 **OF CHILDREN**

6
7 **Evidence from 193 Surveys in 69 Low- and Middle-Income Countries**
8
9

10
11
12
13 **Authors:**

14
15 **Jana Kuhnt**

16 M.Sc., Department of Economics and Centre for Modern Indian Studies, University of Göttingen,
17 Göttingen, Germany
18

19
20 **Sebastian Vollmer***

21 Prof. Dr., Department of Economics and Centre for Modern Indian Studies, University of Göttingen,
22 Göttingen, Germany
23 Department of Global Health and Population, Harvard T.H. Chan School of Public Health, Boston
24 MA, USA
25

26 *** Corresponding Author:**

27 Center for Modern Indian Studies
28 Waldweg 26, Altbau
29 Room 1.117
30 37073 Göttingen
31 Germany
32 Tel. +49 (0)551 / 39-8170
33 Fax +49 (0)551 / 39-20417
34 svollmer@uni-goettingen.de
35
36
37

38 **Acknowledgement**

39
40 The data for this research were collected by The Demographic and Health Surveys Program
41
42 (www.dhsprogram.com), under a contract from the U.S. Agency for International Development.
43
44
45

46 **Manuscript word count: 3402**
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Objectives: Antenatal care (ANC) is an essential part of primary health care and its provision has expanded worldwide. There is limited evidence of large-scale cross-country studies on the impact of ANC offered to pregnant women on child health outcomes. We investigate the association of ANC in low- and middle-income countries on short- and long-term mortality and nutritional child outcomes.

Setting: We used nationally representative health and welfare data from 193 Demographic and Health Surveys (DHS) conducted between 1990 and 2013 from 69 low- and middle-income countries for women of reproductive age (15-49), their children, and respective household.

Participants: The analytical sample consisted of 752,635 observations for neonatal mortality, 574,675 observations for infant mortality, 400,426 observations for low birthweight, 501,484 observations for stunting, 512,424 observations for underweight.

Main Outcomes and Measures: Outcome variables are neonatal and infant mortality, low birthweight, stunting and underweight.

Results: At least one antenatal care visit was associated with a 1.04 percentage points reduced probability of neonatal mortality and a 1.07 percentage points lower probability of infant mortality. Having at least four ANC visits and at having at least once seen a skilled provider reduced the probability by an additional 0.56 and 0.42 percentage points respectively. At least one ANC visit is associated with a 3.82 percentage points reduced probability of giving birth to a low birthweight and a 4.11 and 3.26 percentage points reduced stunting and underweight probability. Having at least four ANC visits and at having at least once seen a skilled provider reduced the probability by an additional 2.83, 1.41 and 1.90 percentage points respectively.

Conclusions: The currently existing and accessed ANC services in low- and middle-income countries are directly associated with improved birth outcomes and longer-term reductions of child mortality and malnourishment.

STRENGTHS AND LIMITATIONS

* First study on the association of antenatal care with child health and vital outcomes for all low- and middle-income countries for which high quality and comparable data are available.

1
2
3 * First study investigating possible long-term effects of the utilization of ANC on children's
4 nutritional and vital status.
5

6
7 * The study focuses on the association between the antenatal care services effectively available
8 and accessible to women in low- and middle-income countries and hence, generates knowledge
9 on the current status quo and effect of the ANC services on child health.
10

11
12
13
14 * The analysis does not allow a causal interpretation of the results.
15

16
17 * Data availability limits the investigation of the association of more disaggregated quality
18 indicators of ANC with the outcome variables.
19

INTRODUCTION

Despite strong international efforts to expand the worldwide coverage of basic primary health services for women, pregnancy and childbirth still represent a high-risk period for mother and child, especially in low- and middle-income countries. As part of the Sustainable Development Goal 3, reductions in maternal and early child mortality are remaining high on the global development policy agenda.¹ Still nearly three million babies die every year during their first month of life while especially in low- and middle-income countries many of those deaths and morbidities are due to easily preventable causes.^{2,3} Undetected infections during pregnancy, like malaria, syphilis, tuberculosis, tetanus, or HIV/AIDS, as well as high blood pressure, diabetes, and other pre-existing health conditions often complicate or aggravate a pregnancy and pose significant risk for mother and child. Antenatal care (ANC) – the services offered to mother and unborn child during pregnancy – is an essential part of the basic primary health care offering a mosaic of services that can prevent, detect, and treat such risk factors early on in the pregnancy. The detection of high-risk pregnancies through the analysis of socioeconomic, medical and obstetrical factors represents a key element of antenatal care. It is also often used as a platform for additional interventions that have been shown to positively influence the maternal and child health status, like immunization and nutrition programs, breastfeeding counselling, or to educate women about the possibilities of family planning and birth spacing.⁴⁻¹³ Additionally, antenatal care programs are used to provide care and information that is not directly related to pregnancy but can reduce possible maternal risk factors, like promoting healthy lifestyles, to tackle malnutrition or inform about gender-based violence. Hence, antenatal care is a potentially important determinant in reducing maternal and child morbidity and mortality.¹⁴⁻²² Within the last decades, the provision of antenatal care services has increased worldwide and during 2010-2015 the antenatal care coverage defined as the percentage of women aged 15-49 that attended at least one antenatal care visit with a skilled provider was around 85% globally and around 77% in the least developed countries.^{23,24} To our knowledge, there exists no global study for all low- and middle-income countries, which analyses the association of existing antenatal care services that are offered to pregnant women in low- and middle-income countries on child health outcomes.

1
2
3 Numerous studies have helped to develop an internationally accepted set of so-called, essential
4 antenatal care services by evaluating the effects of single interventions, like tetanus and malaria
5 prevention programs, on maternal and neonatal health²⁵⁻³⁰ or by studying the optimal number and
6 content of antenatal care visits.³¹⁻³⁴ But the de facto offered and utilized set of antenatal care services
7
8 can deviate greatly from those recommended interventions across as well as within countries. A
9 couple of studies evaluate the relationship between the utilization of antenatal care services and
10 perinatal outcomes in individual low- and middle-income countries. The majority have shown
11 positive effects of antenatal care on newborn mortality, the occurrence of stillbirth and pre-term
12 labour, and low birthweight.³⁵⁻⁴³ However, they exclusively focus on single countries, are often
13 conducted at the clinic level, and have small sample sizes. This limits their external validity. We
14 identified only one study that focuses on a larger regional sample. Conde-Agudelo et al. (2000)
15 studied 837,232 births in Latin America between 1985 and 1997.⁴⁴ One major risk factor associated
16 with fetal death was the lack of antenatal care. We could not find a study that took into account
17 possible long-term effects of the utilization of antenatal care services on children's nutritional and
18 vital status.

19
20
21
22
23
24
25
26
27
28
29
30
31
32
33 With up to 193 Demographic and Health Surveys (DHS) from up to 69 low- and middle-income
34 countries we use the most comprehensive data for low- and middle-income countries in the world
35 that currently exist. Specifically, we investigate whether the attendance of mothers at antenatal care
36 services was associated with improved short- and long-term survival rates or reductions in the
37 prevalence of low birthweight, stunting and underweight in their children.

38 39 40 41 42 43 44 METHODS

45 46 **Data**

47
48 We used data from the Demographic and Health Surveys (DHS), which are publicly available
49 (<http://dhsprogram.com>). The DHS are cross-sectional household surveys that use a harmonized
50 questionnaire to facilitate between-country comparisons. The DHS collect nationally representative
51 health and welfare data for women of reproductive age (15-49 years), their children and the
52 respective household.
53
54
55
56
57
58
59
60

1
2
3 They have been conducted at different time intervals in 90 low- and middle-income countries since
4
5 1985. We included all surveys, which have information for the relevant outcome and explanatory
6
7 variables. The final sample consists of pooled data from up to 193 surveys in 69 low- and middle-
8
9 income countries worldwide conducted between Jan 1, 1990, and Dec 31, 2013. The DHS used a
10
11 multistage stratified sampling, countries were divided into regions, within which populations were
12
13 stratified by area of residence and from which a random sample was taken according to the most
14
15 recent population census. Those are the primary sampling units (clusters with on average 250
16
17 households), which are equally likely to be selected to the proportion the specific cluster's population
18
19 contributed to the total population. At a second stage, after a complete listing of all households
20
21 within the cluster, an average of 20-30 households was randomly selected by equal probability.
22
23 Sampling weights in order to calculate nationally representative statistics are provided by DHS.
24
25 For each live birth within the five (or in some cases three) years previous to the survey the women
26
27 were among other things asked about date of birth, birthweight, vital status at the time of the
28
29 interview, and either current age or age at death of the child. Furthermore, the DHS collected
30
31 information on the height and weight of the children born during the last three or five years.
32
33 For each last-born live birth of the previous three or five years there is information on the attendance
34
35 rates and quality of antenatal care visits during the last pregnancy that led to a live birth. Considering
36
37 the on-going debates on the importance of the quality and number of antenatal care visits^{33,45-48} we
38
39 specified two different main explanatory variables. Firstly, the mere attendance of antenatal care (a
40
41 dummy variable indicating whether the woman attended at least one ANC visit during her last
42
43 pregnancy leading to a live birth) irrespective of the total number of visits and the type of provider. In
44
45 order to proxy the WHO recommendations regarding prenatal care (at least four visits at a skilled
46
47 provider³), we specified a variable indicating whether the woman saw at least once a skilled provider
48
49 during her at least four ANC visits. Unfortunately, we were unable to identify whether all ANC visits
50
51 were provided by a skilled professional. Antenatal care visits to a doctor, midwife, nurse, auxiliary
52
53 midwife, obstetrician, health professional, or trained (traditional) birth attendant, we considered as
54
55
56
57

58 ¹ In 2016 the WHO updated their recommendations to at least eight prenatal care visits at skilled providers⁴⁹.
59
60

1
2
3 skilled antenatal care services. Whereas antenatal care with a traditional birth attendant, relatives,
4 any other person or none of the mentioned was classified as unskilled antenatal care.
5
6
7

8 9 **Outcomes**

10 We analysed the data for short- and long-term vital outcomes and low birthweight of all last-born
11 live births as well as stunting and underweight for the last-born children aged 0-59 (in some surveys
12 0-36) months at the time of the interview. Hence, each woman is represented only once in the
13 dataset with the information of her last born child (in case of a live birth). Mortality outcomes were
14 neonatal death, defined as death of a live birth within the first month of life and infant death, defined
15 as death after the first month but within the first year of life. The latter excludes neonatal deaths and
16 is restricted to children that aged at least one year. Nutritional outcomes were low birthweight,
17 stunting and underweight. We used the WHO classification that defines low birthweight as a
18 birthweight below 2,500g at birth. Following WHO and UNICEF suggestions, we only included
19 biologically plausible birthweights from 500 – 5999g. To calculate stunting and underweight, we
20 used anthropometric data defined by WHO standards and classifications (using the Stata package
21 'igrowup_stata'). Comparing the child's height and weight to those of a well-nourished reference
22 population of the same age and sex, allows us to calculate the z-scores of height-for-age and weight-
23 for-age. Stunting is defined by a height-for-age z-score of less than -2 and underweight is defined by
24 a weight-for-age z-score of less than -2. Biologically implausible values of the z-scores were excluded
25 following WHO guidelines.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43

44 **Statistical Analysis**

45 We used linear probability regression models to investigate the association between antenatal care
46 services and short- and longer-term vital and health outcomes of children. We adjusted the
47 regressions for confounding factors and controlled for PSU fixed effects. The PSU fixed effects are
48 survey specific and herewith, we control for common factors faced by households in the same PSU at
49 one point in time, such as the local availability and quality of
50 health providers, and other local factors. Standard errors were clustered at the PSU level as
51 respondents in the same PSU might experience common shocks. They capture characteristics of
52
53
54
55
56
57
58
59
60

1
2
3 local enumeration areas that are common to all respondents from that area. We used sex, birth order
4
5 (5 categories: ranging from "First born" to "Fifth or later born" child), birth spacing (5 categories:
6
7 ranging from "no preceding birth" to "equal or more than 36 months"), birth month, and whether the
8
9 child was a multiple birth; the mother's age at birth (5 categories: ranging from "below 17" to "equal
10
11 or above 30" years), education (5 categories: ranging from "no education" to "higher education"),
12
13 work status, relation to the household head (dummy indicating whether mother is household head),
14
15 and her marital status; and household wealth quintile as covariates. The wealth quintile variable is
16
17 constructed by using a principal component analysis and is based on the ownership of household
18
19 assets (e.g. electricity, television, quality of dwelling) and indicates the household's wealth relative to
20
21 other households within the respective country in that survey. Additionally, by including variables
22
23 indicating place of delivery, mode of delivery (vaginal or caesarean), status of tetanus injection of
24
25 mother before birth, and if she breastfed at least one month after birth (several only applicable for
26
27 long-term outcomes), we inspected possible mediator variables meaning that the uptake of
28
29 antenatal care services might starkly influence those variables, which themselves might affect the
30
31 outcome variables.

32
33 Using Stata (version 14.0) for all statistical analyses, we also took into account the stratified survey
34
35 design by using the Stata svy command. We used sampling weights provided by DHS in all our
36
37 regressions.
38
39

40 RESULTS

41
42 Our initial samples consisted of surveys for which the respective outcome variable and the
43
44 information on antenatal care visits were collected and consisted of children between 0 and 59
45
46 months at the time of the interview and that were permanent members of the respective household.
47
48 The total sample for neonatal and infant mortality included observations for 1,019,463 children. In
49
50 some survey rounds data on birthweight and in others on anthropometric measures was not
51
52 systematically collected. This left us with data of 947,365 children, where information on their
53
54 birthweight could have potentially been recorded. For stunting this amounted up to 865,959
55
56 children and for underweight to 857,908 children.
57
58
59
60

1
2
3 Observations were lost due to missing data on outcome variables, missing data on the ANC variables
4 (including the dummies indicating the mere attendance of ANC visits, and the attendance of at least
5 4 ANC visits while the woman at least once saw a skilled provider) or missing data on covariates. The
6 final analytical sample was 752,635 for neonatal mortality, 400,426 for low birthweight, 574,675 for
7 infant mortality, 501,484 for stunting and 512,424 for underweight (see Figure 1 and Table A1).
8 The prevalence of newborn death was higher among women that did not receive any antenatal care
9 check-up (3.12 percent) compared to those attending at least one check-up (1.67 percent). For infant
10 mortality the respective numbers are 4.23 percent and 2.21 percent. Prevalence of all outcome
11 variables was higher among women not attending any ANC visit than those attending at least one
12 ANC check-up and compared to those that received at least four ANC visits while they at least once
13 have seen a skilled provider (Table 1). Pregnant women that did not attend antenatal care visits were
14 on average less educated and poorer than those women that attended at least one ANC check-up
15 (see Table A2 in the appendix).
16
17
18
19
20
21
22
23
24
25
26
27
28
29

30 In Table 2 we report the association between antenatal care take-up and short- and long-term
31 mortality outcomes. For each outcome, we show the results from three different specifications
32 where PSU fixed effects are included in all three. The first column shows the mere association
33 between the attendance of at least one ANC visit without controlling for any covariates. The second
34 column shows the association adjusted for control variables and the third column reports the
35 coefficients while adding whether the mother has received at least four antenatal care visits during
36 pregnancy while at least having once seen a skilled provider. The interpretation of this additional
37 term follows the logic of an interaction term as it overlaps in its definition with the variable indicating
38 the mere attendance of ANC. Hence, it shows the additional effect if the mother followed more
39 closely the WHO recommendations. In the appendix (Tables A3 & A4) we report the regression
40 results where in addition to control variables we also adjusted for potential transmission channels of
41 antenatal care (place of delivery, mode of delivery, status of tetanus toxoid (TT) injection of mother
42 before birth, and whether the mother breastfed at least for one month after birth). We will focus on
43 the second and third specification adjusted for control variables and only refer to the other
44 specifications for comparison purposes.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Women attending at least one antenatal care visit have a 1.04 percentage points reduced probability
4 of their newborn dying within the first month after birth and a 1.07 percentage points lower
5 probability of experiencing death of their child within the first year of life. Following the WHO
6 recommendations on ANC visits is significantly related to lower mortality outcomes. Compared to
7 the mere attendance of less than four ANC visits (irrespective of the quality of the provider), having
8 at least four ANC visits and at having at least once seen a skilled provider is reducing the probability
9 of neonatal deaths by an additional 0.56 percentage points and is associated with an additional 0.42
10 percentage points reduction in the probability of infant deaths.

11
12 The DHS dataset also provides information on several variables that are well established in the
13 literature to impact mortality and morbidity outcomes of children and that simultaneously are
14 potentially influenced by ANC attendance. When controlling for these potential transmission
15 channels of antenatal care services, it can be seen that the majority of the ANC coefficients are
16 somewhat attenuated when controlling for these additional variables but not by much (Table A3).
17 Results for all covariates are provided in the appendix (Table A5).

18
19
20
21
22 In Table 3 we report the association between antenatal care and short- and long-term nutritional
23 outcomes of the child. If the mother attends at least one ANC visit, this is associated with a 3.82
24 percentage points reduced probability of giving birth to a low birthweight baby. Stunting and
25 underweight outcomes are reduced by 4.11 and 3.26 percentage points respectively. Attendance at a
26 skilled provider during at least one of at least four ANC visits further reduces the probability of having
27 a low birthweight baby by 2.83 percentage points, for stunting by 1.41 percentage points and for
28 underweight by 1.90 percentage points.

29
30
31 Adding potential transmission channels of antenatal care services to the regression slightly
32 attenuates the ANC coefficients in case of low birthweight and underweight (Table A4). Results for
33 all covariates are provided in the appendix (Table A5).

34 35 36 DISCUSSION

37
38 Most existing evidence on the effect of antenatal care on child health is based on data from high-
39 income countries and their conclusions are not easily transferable to low- and middle-income
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 settings. The existing studies for low- and middle-income countries often focus on individual
4
5 countries. Furthermore, the studied effects of antenatal care have been limited to direct short-term
6
7 maternal and child delivery outcomes. This is the first large-scale cross-country study for all low-and
8
9 middle-income countries with availability of comparable data for ANC, which systematically
10
11 investigates the association of ANC with short- and long-term mortality and nutritional child
12
13 outcomes.

14
15 Using child vital data and child anthropometry from up to 193 surveys in 69 low- and middle-income
16
17 countries we have shown that antenatal care is associated with reductions in neonatal and infant
18
19 mortality, low birthweight, stunting and underweight. While we measure average effects across
20
21 countries and years, we find that this association remains relatively stable across survey rounds
22
23 (Table A6) and can be seen for all outcomes in almost all world regions (Figures A1 & A2) though it is
24
25 especially strong in Latin America and Caribbean, Sub-Saharan Africa, and South and Southeast Asia
26
27 (which constitute about 90% of our sample). Receiving prenatal care by skilled providers and
28
29 attending at least four ANC visits is significantly associated with additional prevalence reductions of
30
31 all outcome variables and hence, plays an important role in the provision of ANC services. The
32
33 magnitude of the association is quantitatively important, as it varies around -1.04 and -4.11
34
35 percentage points.

36
37 Many pregnant women in low- and middle-income countries have no access or do not attend ANC
38
39 services regularly enough (4+ visits) or at skilled providers.^{23,50,51} According to our results, improving
40
41 the coverage and uptake of antenatal care services could be an important tool to improve short- and
42
43 even long-term mortality and nutritional outcomes of children.

44
45 There are a couple of self-selection issues and limitations, which we have attempted to address.
46
47 Unfortunately, we do not have disaggregated information on the type of provider (skilled / unskilled)
48
49 for each ANC visit. We try to proxy this by including whether the woman has at least once seen a
50
51 skilled provider during her pregnancy. Further, we controlled for mother's education and household
52
53 wealth, since more educated or more affluent mothers might be more likely to seek antenatal care
54
55 and at the same time have a better overall health status. Similarly, we adjusted for PSU fixed effects
56
57 to control for community characteristics, overall health status in the region and the local availability
58
59 and quality of health care services as well as other characteristics that are common to the local area.
60

1
2
3 However, there are a few maternal characteristics, which we did not observe and therefore were not
4 able to control for. For instance, if pregnant women feel that there could be something wrong, they
5 might be more likely to seek antenatal care. Similarly, if women had negative birth outcomes in the
6 past, they might also be more likely to seek antenatal care to avoid the repetition of the negative
7 birth outcome. Both cases would downward bias our estimates, and the true association would be
8 even stronger than the association, which we found in our analysis. It is important to point out again
9 that we are only including the outcomes of livebirths. The attendance at antenatal care services
10 might lead to better survival chances of those babies that would have otherwise died before birth.
11 This might impose a downward bias on our estimates. However, there are also potential selection
12 issues, which could bias the results in the other direction. For instance it is unclear how women would
13 behave in case of an unwanted pregnancy, they might be less likely to seek antenatal care and have
14 worse overall health behaviour compared to women in planned pregnancies. In a robustness check
15 we controlled for an indicator variable if the pregnancy was wanted and this did not change the
16 results. Also we cannot further approximate the quality of care received by the women. As the
17 quality of care will influence the effect of ANC this limits our study. By including PSU level fixed
18 effects we absorb indicators that are similar across this geographical unit and survey. Assuming that
19 the quality of ANC available to women within the same PSU is comparable, we successfully address
20 this data limitation. We also assume that missing data in our sample was not systematically
21 correlated with the true unobserved child health and vital outcomes and the availability and
22 accessibility of ANC services.

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42 In summary, our study provides evidence for the potential importance of antenatal care for
43 improving child health and vital outcomes in low- and middle-income countries and might be an
44 important tool to reach the third Sustainable Development Goal by 2030.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 A. CONTRIBUTOR SHIP STATEMENT

4 JK and SV conceptualized the study, developed the analytical strategy and interpreted the data. JK
5 conducted the statistical analysis and wrote the first draft of the manuscript. SV critically revised the
6 manuscript.

7
8 B. COMPETING INTERESTS

9 The authors have read and understood BMJ policy on declaration of interests and declare that they
10 have no competing interests.

11
12 C. FUNDING

13 This research received no specific grant from any funding agency in the public, commercial or not-
14 for-profit sectors.

15
16 D. DATA SHARING STATEMENT

17 This study used data that was collected by the Demographic and Health Surveys Program
18 (www.dhsprogram.com), under a contract from the U.S. Agency for International Development.

19
20 E. ETHICAL APPROVAL

21 Procedures and questionnaires for standard DHS surveys have been approved by the ICF Institutional
22 Review Board (IRB) as well as by the relevant body in each country.

23 ICF IRB ensures that the survey complies with the U.S. Department of Health and Human Services
24 regulations for the protection of human subjects (45 CFR 46), while the host country IRB ensures that
25 the survey complies with laws and norms of the nation. Oral informed consent for the interview was
26 obtained from respondents by interviewers.

REFERENCES

1. UN. Sustainable Development Goals. Goal 3: Ensure healthy lives and promote well-being for all at all ages. United Nations Sustainable Development Homepage. 2017 [accessed on the 12.07.2017: <http://www.un.org/sustainabledevelopment/health/>].
2. WHO. Global Health Observatory (GHO) data: Neonatal Mortality. World Health Organization. 2017 [accessed on 12.07.2017: http://www.who.int/gho/child_health/mortality/neonatal_text/en/].
3. UNICEF. Committing to Child Survival: A Promise Renewed. Progress Report 2015. United Nations Children's Fund (UNICEF). 2015.
4. Imdad A, Bhutta Z. Effects of calcium supplementation during pregnancy on maternal, fetal and birth outcomes. *Paediatric and perinatal epidemiology*. 2012;26(5):138-152.
5. Gupta A, Sharma D. Breastfeeding and Child Health. *Economic and Political Weekly*. 2006;April 15.
6. Oddy W, Kendall G, Li J, et al. The long-term effects of breastfeeding on child and adolescent mental health: a pregnancy cohort study followed for 14 years. *Journal of Pediatrics*. 2010;156(4):568-574.
7. Black R, Victora C, Walker S, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *The Lancet*. 2013;382:427-51.
8. Titaley C, Dibley M. Antenatal iron/folic acid supplements, but not postnatal care, prevents neonatal deaths in Indonesia: analysis of Indonesia Demographic and Health Surveys. *BMJ Open*. 2012;2(6).
9. Ota E, Hori H, Mori R, Tobe-Gai R, Farrar D. Antenatal dietary education and supplementation to increase energy and protein intake. *Cochrane Database Systematic Review*. 2015;(6).
10. Pena-Rosas JP, De-Regil LM, Garcia-Casal MN, Dowswell T. Daily oral iron supplementation during pregnancy. *Cochrane Database Systematic Review*. 2015;(7).
11. Haider BA, Bhutta ZA. Multiple-micronutrient supplementation for women during pregnancy. *Cochrane Database Systematic Review*. 2015;(11).

12. de Jongh TE, Gurol-Urganci I, Allen E, Zhu NJ, Atun R. Integration of antenatal care services with health programmes in low- and middle-income countries: systematic review. *Journal of Global Health*. 2016;6(1).
13. McNellan CR, Dansereau E, Colombara D, Palmisano E, Wallace M, Johanns C et al. Uptake of Antenatal Care, and its Relationship with Participation in Health Services and Behaviors: An Analysis of the Poorest Regions of Four Mesoamerican Countries. *Annals of Global Health*. 2017;83(1):193-194.
14. Carroli G, Rooney C, Villar J. How effective is antenatal care in preventing maternal mortality and serious morbidity? An overview of the evidence. *Paediatric perinatal Epidemiology*. 2011;15(Suppl 1):1-42.
15. Abou-Zahr C, Wardlaw T. Antenatal care in developing countries: promises, achievements and missed opportunities. Geneva: World Health Organization; 2003.
16. Chen XK, Wen SW, Yang Q, Walker M. Adequacy of prenatal care and neonatal mortality in infants born to mothers with and without antenatal high-risk conditions. *Australian and New Zealand Journal of Obstetrics and Gynaecology*. 2007;47(2): 122-7.
17. Khan K, Wojdyla D, Say L, Gulmezoglu A, VanLook P. WHO analysis of causes of maternal death: a systematic review. *The Lancet*. 2006;367:1066-1074.
18. Lincetto O, Mothebesoane-Anoh S, Gomez P, Munjanja S. Antenatal Care. In Opportunities for Africa's Newborns. PMNCH; 2006.
19. World Health Organization. WHO Statement on Antenatal Care. World Health Organization: Geneva, Switzerland; 2011.
20. Zanconato G, Msolomba R, Guarenti L, Franchi M. Antenatal care in developing countries: the need for a tailored model. *Seminars in Fetal and neonatal Medicine*. 2006;11(1):15-20.
21. PMNCH. The PMNCH Report 2012. Analysing Progress on Commitments to the Global Strategy for Women's and Children's Health. World Health Organization: Geneva, Switzerland; 2012.
22. Moss W, Darmstadt G, Marsh D, Black R, Santosham M. Research Priorities for the Reduction of Perinatal and Neonatal Morbidity and Mortality in Developing Country Communities. *Journal of Perinatology*. 2002;22(6): 484-495.

23. UNICEF. UNICEF Data: Monitoring the Situation of Children and Women. UNICEF. 2017. [accessed on 12.07.2017: <https://data.unicef.org/topic/maternal-health/antenatal-care/#>].
24. Wang W, Alva S, Wang S, Fort A. Levels and Trends in the Use of Maternal Health Services in Developing Countries. DHS Comparative Reports No. 26. Calverton, Maryland, USA: ICF Macro; 2011.
25. Bhutta Z, Das J, Bahl R, et al. Can available interventions end preventable deaths in mothers, newborn babies, and stillbirths, and at what cost? *The Lancet*. 2011;384(9940):347-370.
26. Blencowe H, Lawn J, Vandelaer J, Roper M, Cousens S. Tetanus toxoid immunization to reduce mortality from neonatal tetanus. *International Journal of Epidemiology*. 2010;39(suppl 1): i102-i109.
27. Demicheli V, Barale A, Rivetti A. Vaccines for women for preventing neonatal tetanus. *Cochrane Database Systematic Reviews*. 2015;(7).
28. Blencowe H, Cousens S, Kamb M, Berman S, Lawn J. Lives Saved Tool supplement detection and treatment of syphilis in pregnancy to reduce syphilis related stillbirths and neonatal mortality. *BMC Public Health*. 2011;11(Suppl 3).
29. Kayentao K, Garner P, van Eijk AM, Naidoo I, Roper C, Mulokozi A et al. Intermittent preventive therapy for malaria during pregnancy using 2 vs 3 or more doses of sulfadoxine-pyrimethamine and risk of low birth weight in Africa: systematic review and meta-analysis. *JAMA*. 2013;309:594-604.
30. Althabe F, Belizán JM, McClure EM, Hemingway-Foday J, Berrueta M, Mazzoni A et al. A population-based, multifaceted strategy to implement antenatal corticosteroid treatment versus standard care for the reduction of neonatal mortality due to preterm birth in low-income and middle-income countries: the ACT cluster-randomised trial. *The Lancet*. 2015;9968:629-639.
31. Villar J, Ba'aqel H, Piaggio G, et al. WHO antenatal care randomised trial for the evaluation of a new model of routine antenatal care. *The Lancet*. 2001;357:1551-1564.
32. Carroli G, Villar J, Piaggio G, et al. WHO systematic review of randomized controlled trials of routine antenatal care. *The Lancet*. 2001;357:1565-70.

- 1
2
3 33. Lavin T, Pattinson RC. Does antenatal care timing influence stillbirth risk in the third
4 trimester? A secondary analysis of perinatal death audit data in South Africa. An
5 International Journal of Obstetrics and Gynaecology (BJOG). 2017.
6
7
8
9 34. Nimi T, Fraga S, Costa D, Campos P, Barros H. Prenatal care and pregnancy outcomes: A
10 cross-sectional study in Luanda, Angola. *International Journal of Gynecology and Obstetrics*.
11 2016; 135(S1):S72-S78.
12
13
14 35. Hong R, Ruiz-Beltran M. Impact of Prenatal Care on Infant Survival in Bangladesh. *Maternal*
15 *and Child Health Journal*. 2007;11:199-206.
16
17
18 36. Kapoor S, Anand K, Kumar G. Risk Factors for Stillbirths in A Secondary Level Hospital at
19 Ballabgarh, Haryana: A Case Control Study. *Indian Journal of Pediatrics*. 1994;61:161-66.
20
21
22 37. Mbuagbaw L, Gofin R. A new measurement for optimal antenatal care: determinants and
23 outcomes in Cameroon. *Maternal and Child Health Journal*. 2011;15(8):1427-34.
24
25
26 38. Brown C, Sohani S, Khan K, Lilford R, Mukhwana W. Antenatal care and perinatal outcomes
27 in Kwale district, Kenya. *BMC Pregnancy and Childbirth*. 2008;8(2).
28
29
30 39. Celik Y, Younis M. Effects of antenatal care services on birthweight: importance of model
31 specification and empirical procedure used in estimating the marginal productivity of health
32 inputs. *Journal of Medical Systems*. 2007;31(3):197-204.
33
34
35 40. McCaw Binns A, Greenwood R, Ashley D, Golding J. Antenatal and Perinatal-Care in Jamaica
36 - do They Reduce Perinatal Death Rates? *Paediatric and Perinatal Epidemiology*.
37 1994;8(Suppl 1):86-97.
38
39
40 41. Cruz Barbosa IR, Merencio Silva WB, Gutierrez Cerqueira GS, Ferreira Novo N, Almeida FA,
41 Garcia Novo JLV. Maternal and fetal outcome in women with hypertensive disorders of
42 pregnancy: the impact of prenatal care. *Therapeutic Advances in Cardiovascular Disease*.
43 2015;9(4):140-146.
44
45
46 42. Abbas AM, Rabeea M, Abdel Hafiz HA, Ahmed NH. Effects of irregular antenatal care
47 attendance in primiparas on the perinatal outcomes: a cross sectional study. *Proceedings in*
48 *Obstetrics and Gynecology (POG) in Press*. 2017;3.
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3 43. Ballard K, Belete Z, Kinfu H, Tadesse M, Amin M, Atnafu H, The effect of prenatal and
4 intrapartum care on the stillbirth rate among women in rural Ethiopia. *International Journal*
5 *of Gynaecology and Obstetrics*. 2016;133(2):164-167.
6
7
8
9 44. Conde-Agudelo A, Belizán J, Díaz-Rossello J. Epidemiology of fetal death in Latin America.
10 *Acta Obstetrica et Gynecologica Scandinavica*. 2000;79:371-378.
11
12 45. Vogel P, Habib N, Souza J, et al. Antenatal care packages with reduced visits and perinatal
13 mortality: a secondary analysis of the WHO Antenatal Care Trial. *Reproductive Health*.
14 2013;10(19).
15
16
17 46. Dowswell T, Carroli G, Duley L, Gates S, Gülmezoglu AM, Khan-Neelofur D, Piaggio G.
18 Alternative versus standard packages of antenatal care for low-risk pregnancy. *Cochrane*
19 *Database of Systematic Reviews*. 2015;7.
20
21 47. Hofmeyr GJ, Hodnett ED. Antenatal care packages with reduced visits and perinatal
22 mortality: a secondary analysis of the WHO Antenatal Care Trial – Commentary: routine
23 antenatal visits for healthy pregnant women do make a difference. *Reproductive Health*.
24 2013;10(20).
25
26
27 48. Hodgins S, D’Agostino A. The quality-coverage gap in antenatal care: toward better
28 measurement of effective coverage. *Global Health: Science and Practice*. 2014.
29
30
31 49. World Health Organization. WHO recommendations on antenatal care for a positive
32 pregnancy experience. World Health Organization: Geneva, Switzerland. 2016.
33
34
35 50. You D, Hug L, Ejdmeyr S, Beise J. Levels and Trends in Child Mortality. Report 2015. United
36 Nations Children’s Fund: New York, USA. 2015.
37
38
39 51. World Health Organization. World Health Statistics. World Health Organization: Geneva,
40 Switzerland. 2016.
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Descriptive Statistics

Outcome variable	Mother attended at least 1 ANC visit; % (n= 616347)	Mother attended at least 4 ANC visits & has seen skilled provider; (n=416350)	Mother did not attend ANC visit, % (n=136288)
Neonatal Mortality	1.67*	1.40*	3.12
Infant Mortality	2.21*	1.81*	4.23
Low Birthweight	9.69*	8.94*	14.35
Stunting	31.35*	27.18*	47.08
Underweight	15.70*	12.02*	30.72
n values denote the number of observations in the neonatal mortality sample.			
* Proportions to the group "Mother did not attend ANC visit" were significantly different from each other, $p \leq 0.05$ (t test).			

Table 2: Associations between antenatal care visits and mortality outcomes

	Neonatal Mortality			Infant Mortality		
	(1)	(2)	(3)	(1)	(2)	(3)
<i>at least 1 ANC visit</i>	-0.0107*** [0.001]	-0.0104*** [0.001]	-0.00783*** [0.001]	-0.0127*** [0.001]	-0.0107*** [0.001]	-0.00873*** [0.001]
<i>at least 4 ANC visits & skilled ANC provider</i>			-0.00557*** [0.001]			-0.00424*** [0.001]
<i>N</i>	752635	752635	752635	574675	574675	574675
<i>Adjusted for confounding</i>	no	yes	yes	no	yes	yes

All regressions include PSU fixed effects. ***, ** and * denote significance at the 1%, 5%, and 10% levels, respectively. Std. Errors in square brackets and are clustered at PSU level. Control variables include: Mother's age, marital status and educational achievement, whether she heads the HH, child's sex and birth order and spacing, month of birth, whether it was a multiple birth and HH wealth quintile.

Table 3: Associations between antenatal care visits and nutritional outcomes

Low Birthweight	(1)	(2)	(3)
<i>at least 1 ANC visit</i>	-0.0428*** [0.004]	-0.0382*** [0.004]	-0.0187*** [0.004]
<i>at least 4 ANC visits & skilled ANC provider</i>			-0.0283*** [0.002]
<i>N</i>	400426	400426	400426
Stunting	(1)	(2)	(3)
<i>at least 1 ANC visit</i>	-0.0677*** [0.003]	-0.0411*** [0.003]	-0.0345*** [0.003]
<i>at least 4 ANC visits & skilled ANC provider</i>			-0.0141*** [0.002]
<i>N</i>	501484	501484	501484
Underweight	(1)	(2)	(3)
<i>at least 1 ANC visit</i>	-0.0502*** [0.002]	-0.0326*** [0.002]	-0.0237*** [0.002]
<i>at least 4 ANC visits & skilled ANC provider</i>			-0.0190*** [0.002]
<i>N</i>	512424	512424	512424
<i>Adjusted for confounding</i>	no	yes	yes

All regressions include PSU fixed effects. ***, ** and * denote significance at the 1%, 5%, and 10% levels, respectively. Std. Errors in square brackets and are clustered at PSU level. Control variables include: Mother's age, marital status and educational achievement, whether she heads the HH, child's sex and birth order and spacing, month of birth, whether it was a multiple birth and HH wealth quintile.

1
2
3 Figure 1: Sample deduction
4
5
6

7 Own calculation.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Neonatal Mortality	Low Birthweight	Infant Mortality	Stunting	Underweight
1019463	947365	1019463	865959	857908
8071 missing data for outcome variable	471146 missing data for outcome variable	215343 missing data for outcome variable	257991 missing data for outcome variable	231246 missing data for outcome variable
1011392	476219	804120	607968	626662
11652 missing data for ANC variables	61722 missing data for ANC variables	194395 missing data for ANC variables	79298 missing data for ANC variables	80304 missing data for ANC variables
799740	414497	609725	528670	546358
17105 missing data for control variables	14071 missing data for control variables	35050 missing data for control variables	27186 missing data for control variables	33934 missing data for control variables
752635	400426	574675	501484	512424
69 LMIC 193 surveys	68 LMIC 180 surveys	69 LMIC 193 surveys	64 LMIC 171 surveys	64 LMIC 170 surveys

Figure 1

609x457mm (300 x 300 DPI)

ew only

Table A1: Sample size and surveys across countries

	Phase (Year of survey)	No. obs. (% of total)
Albania	Phase 5 (2008/09)	1,152 (0.15)
Armenia	Phase 4 (2000), Phase 5 (2005), Phase 6 (2010)	2,778 (0.37)
Azerbaijan	Phase 5 (2006)	1,174 (0.16)
Bangladesh	Phase 3 (1993/94, 1996, 1997)*, Phase 4 (1999, 2000)*, Phase 5 (2004, 2007)*, Phase 6 (2011)	23,459 (3.12)
Benin	Phase 3 (1996), Phase 4 (2001)*, Phase 5 (2006), Phase 6 (2011/12)	16,274 (2.16)
Bolivia	Phase 3 (1993/94), Phase 4 (2003/04), Phase 5 (2008)	17,034 (2.26)
Brazil	Phase 2 (1991/92), Phase 3 (1996)	5,339 (0.71)
Burkina Faso	Phase 2 (1992/93), Phase 4 (1998/99, 2003)*, Phase 6 (2010)	15,095 (2.01)
Burundi	Phase 6 (2010/11)	3,224 (0.43)
Cambodia	Phase 4 (2000), Phase 5 (2005/06), Phase 6 (2010/11)	14,645 (1.95)
Cameroon	Phase 2 (1991), Phase 3 (1998), Phase 4 (2004), Phase 6 (2011)	10,344 (1.37)
Central Afr. Republic	Phase 3 (1994/95)	1,451 (0.19)
Chad	Phase 3 (1996/97), Phase 4 (2004)	5,821 (0.77)
Colombia	Phase 3 (1995), Phase 5 (2004, 2005)*, Phase 6 (2009/10)	24,944 (3.31)
Comoros	Phase 3 (1996)	595 (0.08)
Congo (DRC)	Phase 5 (2007)	3,297 (0.44)
Congo (Brazzaville)	Phase 5 (2005), Phase 6 (2011/12)	6,281 (0.83)
Dom. Republic	Phase 2 (1991), Phase 3 (1996), Phase 4 (2002), Phase 5 (2007)	17,845 (2.37)
Egypt	Phase 2 (1992/93), Phase 3 (1995/96), Phase 4 (2000, 2003)*, Phase 5 (2005, 2008)*	37,847 (5.03)
Ethiopia	Phase 4 (2000), Phase 5 (2005), Phase 6 (2011)	14,694 (1.95)
Gabon	Phase 4 (2000/01), Phase 6 (2012)	4,039 (0.54)
Ghana	Phase 3 (1993/94), Phase 4 (1998/99, 2003), Phase 5 (2008)	7,067 (0.94)
Guatemala	Phase 3 (1995), Phase 4 (1998/99)	8,340 (1.11)
Guinea	Phase 4 (1999), Phase 5 (2005), Phase 6 (2012)	8,518 (1.13)
Guyana	Phase 5 (2005, 2009)*	1,004 (0.13)
Haiti	Phase 3 (1994/95), Phase 4 (2000), Phase 5 (2005/06), Phase 6 (2012)	11,927 (1.58)
Honduras	Phase 5 (2005/06), Phase 6 (2011/12)	12,725 (1.69)
India	Phase 4 (1998/99/00), Phase 5 (2005/06)	46,435 (6.17)
Indonesia	Phase 2 (1991), Phase 3 (1994, 1997), Phase 4 (2002/03), Phase 5 (2007), Phase 6 (2012)	64,890 (8.62)
Ivory Coast	Phase 3 (1994), Phase 4 (1998/99), Phase 5 (2005), Phase 6 (2011/12)	8,020 (1.07)
Jordan	Phase 3 (1997), Phase 5 (2007), Phase 6 (2012)	13,205 (1.75)
Kazakhstan	Phase 3 (1995, 1999)*	1,454 (0.19)
Kenya	Phase 2 (1993), Phase 3 (1998), Phase 4 (2003), Phase 5 (2008/09)	9,250 (1.23)
Kyrgyz Republic	Phase 3 (1997), Phase 6 (2012)	2,859 (0.38)
Lesotho	Phase 5 (2004/05), Phase 6 (2009/10)	4,522 (0.60)
Liberia	Phase 5 (2006/07)	2,229 (0.30)
Madagascar	Phase 2 (1992), Phase 3 (1997), Phase 4 (2003/04), Phase 5 (2008/09)	13,746 (1.83)
Malawi	Phase 4 (2000), Phase 5 (2004/05), Phase 6 (2010)	20,486 (2.72)
Maldives	Phase 6 (2009)	2,041 (0.27)

Mali	Phase 3 (1995/96), Phase 4 (2001), Phase 5 (2006)	12,855 (1.71)
Moldova	Phase 5 (2005)	1,153 (0.15)
Morocco	Phase 2 (1992), Phase 5 (2003/04)	6,520 (0.87)
Mozambique	Phase 3 (1997), Phase 4 (2003/04), Phase 6 (2011)	12,008 (1.60)
Namibia	Phase 2 (1992), Phase 4 (2000), Phase 5 (2006/07)	5,961 (0.79)
Nepal	Phase 3 (1996), Phase 4 (2001), Phase 5 (2006), Phase 6 (2011)	11,688 (1.55)
Nicaragua	Phase 3 (1997/98), Phase 4 (2001)	8,658 (1.15)
Nigeria	Phase 2 (1990), Phase 4 (1999, 2003), Phase 5 (2008)	17,896 (2.38)
Niger	Phase 2 (1992), Phase 3 (1998), Phase 5 (2006), Phase 6 (2012)	13,275 (1.76)
Pakistan	Phase 2 (1990/91), Phase 5 (2006/07), Phase 6 (2012/13)	11,236 (1.49)
Paraguay	Phase 2 (1990)	2,421 (0.32)
Peru	Phase 2 (1991/92), Phase 3 (1996), Phase 4 (2000), Phase 5 (2005/06, 2007/08, 2009), Phase 6 (2010/11/12)	57,103 (7.59)
Philippines	Phase 3 (1993, 1998)*, Phase 4 (2003), Phase 5 (2008)	17,568 (2.33)
Rwanda	Phase 2 (1992), Phase 4 (2000), Phase 5 (2005), Phase 6 (2010/11)	15,041 (2.00)
Sao Tome and Principe	Phase 5 (2008/09)	1,072 (0.14)
Senegal	Phase 2 (1992/93), Phase 3 (1997), Phase 4 (2005), Phase 6 (2010/11, 2012/13)*	13,834 (1.84)
Sierra Leone	Phase 5 (2008)	2,202 (0.29)
South Africa	Phase 4 (1998)	3,075 (0.41)
Swaziland	Phase 5 (2006/07)	1,396 (0.19)
Tanzania	Phase 2 (1991/2), Phase 3 (1996), Phase 4 (1999), Phase 5 (2004/5, 2007/8)*, Phase 6 (2009/10)	16,392 (2.18)
Timor-Leste	Phase 6 (2009/10)	3,612 (0.48)
Togo	Phase 4 (1998)	2,196 (0.29)
Turkey	Phase 2 (1993), Phase 4 (1998), Phase 5 (2003/04)	7,980 (1.06)
Uganda	Phase 3 (1995), Phase 4 (2000/01), Phase 5 (2006), Phase 6 (2011)	11,743 (1.56)
Ukraine	Phase 5 (2007)	800 (0.11)
Uzbekistan	Phase 3 (1996)	768 (0.10)
Vietnam	Phase 4 (1997 / 2002)*	2,437 (0.32)
Yemen	Phase 2 (1991/2)	3,369 (0.45)
Zambia	Phase 2 (1992) Phase 3 (1996/97), Phase 4 (2001/02), Phase 5 (2007)	12,572 (1.67)
Zimbabwe	Phase 3 (1994), Phase 4 (1999), Phase 5 (2005/06), Phase 6 (2010/11)	9,754 (1.30)
TOTAL		752,635 (100)
Total numbers and frequencies are based on the neonatal mortality sample. * indicates that more than one survey was conducted within the phase		

Table A2: Covariates across exposure category

Covariate	Mother attended at least 1 ANC visit; % (n=616347)	Mother did not attend ANC visit, % (n=136288)
Maternal education		
<i>None</i>	45.41	82.92*
<i>Primary or incompl. Secondary</i>	33.89	13.38*
<i>Secondary or higher</i>	20.70	3.70*
Maternal age at birth		
<i>≤ 17 years</i>	3.51	4.89*
<i>17 - 19 years</i>	12.89	12.52*
<i>20 - 24 years</i>	28.13	25.31*
<i>25 - 29 years</i>	24.58	22.99*
<i>30 years or older</i>	30.88	34.29*
Mother is working	57.25	51.02*
Mother is HH Head	9.13	7.20*
Mother is married	72.24	85.77*
Child is female	48.65	48.88
Multiple birth	0.91	0.10*
Birth order		
<i>First child</i>	31.23	19.41*
<i>Second child</i>	21.60	16.16*
<i>Third child</i>	15.55	14.67*
<i>Fourth child</i>	10.43	12.53*
<i>Fifth or later born child</i>	21.19	37.23*
Birth spacing ≤ 18 mths	3.56	5.58*
Wealth quintiles		
<i>First quintile</i>	15.17	32.49*
<i>Second quintile</i>	17.50	26.81*
<i>Third quintile</i>	19.88	20.39
<i>Fourth quintile</i>	22.30	13.91*
<i>Fifth quintile</i>	25.14	6.40*
N values denote the number of observations in the neonatal mortality sample. Values were population weighted. * Proportions in the two groups were significantly different from each other, $p \leq 0.05$ (t test).		

Table A3: Associations between mortality outcomes and mediator variables

	Neonatal Mortality		Infant Mortality	
	(1)	(2)	(1)	(2)
<i>at least 1 ANC visit</i>	-0.00520*** [0.001]	-0.00288*** [0.001]	-0.00562*** [0.001]	-0.00416*** [0.001]
<i>at least 4 ANC visits & skilled ANC provider</i>		-0.00556*** [0.001]		-0.00342*** [0.001]
<i>facility delivery</i>	0.00211*** [0.001]	0.00273*** [0.001]	-0.00137* [0.001]	-0.000999 [0.001]
<i>mother had TT injection</i>	-0.00856*** [0.001]	-0.00814*** [0.001]	-0.00619*** [0.001]	-0.00594*** [0.001]
<i>delivery by C-section</i>	0.0101*** [0.001]	0.0103*** [0.001]	0.00106 [0.001]	0.00117 [0.001]
<i>child was breastfed</i>			-0.0501*** [0.002]	-0.0501*** [0.002]
<i>N</i>	689319	689319	522970	522970
Adjusted for confounding	yes	yes	yes	yes

All regressions include PSU fixed effects. ***, ** and * denote significance at the 1%, 5%, and 10% levels, respectively. Std. Errors in square brackets and are clustered at PSU level. Control variables include: Mother's age, marital status and educational achievement, whether she heads the HH, child's sex and birth order and spacing, month of birth, whether it was a multiple birth and HH wealth quintile.

Table A4: Associations between nutritional outcomes and mediator variables

	Low Birthweight		Stunting		Underweight	
	(1)	(2)	(1)	(2)	(1)	(2)
<i>at least 1 ANC visit</i>	-0.0349*** [0.004]	-0.0168*** [0.004]	-0.0458*** [0.003]	-0.0388*** [0.003]	-0.0308*** [0.002]	-0.0225*** [0.003]
<i>at least 4 ANC visits & skilled ANC provider</i>		-0.0274*** [0.002]		-0.0159*** [0.002]		-0.0190*** [0.002]
<i>mother had TT injection</i>	-0.00721*** [0.002]	-0.00532*** [0.002]	0.0000754 [0.002]	0.000984 [0.002]	-0.00566*** [0.002]	-0.00458*** [0.002]
<i>child was breastfed</i>			-0.0264*** [0.005]	-0.0264*** [0.005]	-0.0154*** [0.004]	-0.0153*** [0.004]
<i>child ever got vaccine</i>			0.0539*** [0.004]	0.0546*** [0.004]	-0.00196 [0.003]	-0.00107 [0.003]
<i>N</i>	380570	380570	475936	475936	486040	486040
Adjusted for confounding	yes	yes	yes	yes	yes	yes

All regressions include PSU fixed effects. ***, ** and * denote significance at the 1%, 5%, and 10% levels, respectively. Std. Errors in square brackets and are clustered at PSU level. Control variables include: Mother's age, marital status and educational achievement, whether she heads the HH, child's sex and birth order and spacing, month of birth, whether it was a multiple birth and HH wealth quintile.

Table A5: Associations between the exposure variables and full set of control variables

	Neonatal Mortality	Infant Mortality	Low Birthweight	Stunting	Underweight
Maternal education [reference: no education]					
Primary or incompl. Secondary	-0.00247*** [0.001]	-0.00539*** [0.001]	-0.0191*** [0.002]	-0.0447*** [0.002]	-0.0356*** [0.002]
Secondary or higher	-0.00566*** [0.001]	-0.00890*** [0.001]	-0.0350*** [0.002]	-0.0785*** [0.003]	-0.0460*** [0.002]
Maternal age at birth [reference: ≤ 17yrs]					
17 - 19 years	-0.00474*** [0.001]	-0.00682*** [0.002]	-0.00926** [0.004]	-0.0477*** [0.005]	-0.0340*** [0.004]
20 - 24 years	-0.00735*** [0.001]	-0.0105*** [0.002]	-0.0205*** [0.004]	-0.0714*** [0.004]	-0.0477*** [0.004]
25 - 29 years	-0.00645*** [0.001]	-0.00963*** [0.002]	-0.0218*** [0.004]	-0.0970*** [0.005]	-0.0545*** [0.004]
30 years or older	-0.000352 [0.001]	-0.00881*** [0.002]	-0.00879** [0.004]	-0.114*** [0.005]	-0.0567*** [0.004]
Mother is working	0.00299*** [0.000]	0.000925 [0.001]	-0.000101 [0.001]	0.0155*** [0.002]	0.00591*** [0.001]
Mother is HH Head	-0.000972 [0.001]	-0.0000246 [0.001]	-0.00123 [0.002]	0.00701** [0.003]	-0.00259 [0.002]
Mother is married	-0.00134** [0.001]	-0.00610*** [0.001]	-0.00988*** [0.002]	-0.00849*** [0.002]	-0.0106*** [0.002]
Child is female	-0.00451*** [0.000]	-0.00217*** [0.001]	0.0208*** [0.001]	-0.0399*** [0.001]	-0.0220*** [0.001]
Multiple birth	0.0970*** [0.004]	0.0607*** [0.005]	0.391*** [0.010]	0.155*** [0.009]	0.148*** [0.008]
Birth order [reference: First born child]					
Second child	-0.00904*** [0.001]	-0.00495*** [0.001]	-0.0265*** [0.002]	0.0290*** [0.002]	0.00545*** [0.002]
Third child	-0.0108*** [0.001]	-0.00429*** [0.001]	-0.0342*** [0.002]	0.0456*** [0.003]	0.0137*** [0.002]
Fourth child	-0.0118*** [0.001]	-0.00484*** [0.001]	-0.0350*** [0.003]	0.0574*** [0.003]	0.0194*** [0.003]
Fifth or later born child	-0.0115*** [0.001]	-0.00392*** [0.001]	-0.0417*** [0.003]	0.0835*** [0.003]	0.0276*** [0.003]
Birth spacing ≤ 18 months	0.0218*** [0.001]	0.0234*** [0.002]	0.0240*** [0.004]	0.0462*** [0.004]	0.0333*** [0.003]
Wealth quintiles [reference: poorest 20%]					
Second quintile	0.00102 [0.001]	-0.00174* [0.001]	-0.00848*** [0.002]	-0.0172*** [0.003]	-0.0120*** [0.002]
Third quintile	0.000917 [0.001]	-0.00246** [0.001]	-0.0121*** [0.003]	-0.0371*** [0.003]	-0.0302*** [0.002]

<i>Fourth quintile</i>	0.000564 [0.001]	-0.00496*** [0.001]	-0.0170*** [0.003]	-0.0602*** [0.003]	-0.0511*** [0.003]
<i>Fifth quintile</i>	0.0000414 [0.001]	-0.00619*** [0.001]	-0.0206*** [0.003]	-0.0977*** [0.003]	-0.0790*** [0.003]
Month of birth	0.0000126 [0.000]	-0.0000336 [0.000]	-0.000283* [0.000]	-0.00372*** [0.000]	-0.00290*** [0.000]
N	752635	574675	400426	501484	512424
Results are from the main results regressions of table 2 & 3 (model 3) and are adjusted for the full set of control variables and include PSU fixed effects. ***, ** and * denote significance at the 1%, 5%, and 10% levels, respectively. Std. Errors in square brackets and are clustered at PSU level.					

For peer review only

Table A6: Associations between the exposure variables before and after the introduction of the WHO recommendation (Year 2002)

Neonatal Mortality	Before 2002 (1990-2002)			After 2002 (2003-2013)		
	(1)	(2)	(3)	(1)	(2)	(3)
<i>at least 1 ANC visit</i>	-0.0130*** [0.001]	-0.0129*** [0.001]	-0.00983*** [0.001]	-0.00798*** [0.001]	-0.00755*** [0.001]	-0.00534*** [0.001]
<i>at least 4 ANC visits & skilled ANC provider</i>			-0.00702*** [0.001]			-0.00453*** [0.001]
N	320338	320338	320338	432297	432297	432297
Infant Mortality	Before 2002 (1990-2002)			After 2002 (2003-2013)		
	(1)	(2)	(3)	(1)	(2)	(3)
<i>at least 1 ANC visit</i>	-0.0152*** [0.001]	-0.0126*** [0.001]	-0.00936*** [0.002]	-0.00987*** [0.001]	-0.00827*** [0.001]	-0.00733*** [0.001]
<i>at least 4 ANC visits & skilled ANC provider</i>			-0.00730*** [0.001]			-0.00189** [0.001]
N	244729	244729	244729	329946	329946	329946
Low Birthweight	Before 2002 (1990-2002)			After 2002 (2003-2013)		
	(1)	(2)	(3)	(1)	(2)	(3)
<i>at least 1 ANC visit</i>	-0.0404*** [0.005]	-0.0352*** [0.005]	-0.0181*** [0.006]	-0.0455*** [0.006]	-0.0416*** [0.006]	-0.0202*** [0.006]
<i>at least 4 ANC visits & skilled ANC provider</i>			-0.0259*** [0.003]			-0.0299*** [0.002]
N	145426	145426	145426	255000	255000	255000
Stunting	Before 2002 (1990-2002)			After 2002 (2003-2013)		
	(1)	(2)	(3)	(1)	(2)	(3)
<i>at least 1 ANC visit</i>	-0.0753*** [0.004]	-0.0416*** [0.004]	-0.0333*** [0.004]	-0.0586*** [0.004]	-0.0383*** [0.004]	-0.0335*** [0.004]
<i>at least 4 ANC visits & skilled ANC provider</i>			-0.0187*** [0.003]			-0.00964*** [0.003]
N	217791	217791	217791	283693	283693	283693
Underweight	Before 2002 (1990-2002)			After 2002 (2003-2013)		
	(1)	(2)	(3)	(1)	(2)	(3)
<i>at least 1 ANC visit</i>	-0.0534*** [0.003]	-0.0299*** [0.003]	-0.0204*** [0.003]	-0.0463*** [0.003]	-0.0338*** [0.003]	-0.0257*** [0.004]
<i>at least 4 ANC visits & skilled ANC provider</i>			-0.0214*** [0.003]			-0.0163*** [0.002]
N	223491	223491	223491	288933	288933	288933


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

<i>Adjusted for confounding</i>	no	yes	yes	no	yes	yes
---------------------------------	----	-----	-----	----	-----	-----

All regressions include PSU fixed effects. ***, ** and * denote significance at the 1%, 5%, and 10% levels, respectively. Std. Errors in square brackets and are clustered at PSU level. Control variables include: Mother's age, marital status and educational achievement, whether she heads the HH, child's sex and birth order and spacing, month of birth, whether it was a multiple birth and HH wealth quintile.


For peer review only

Figure A1: Associations between mortality outcomes and ANC attendance by world region



Figures show marginal effects and 95% confidence intervals for binary indicator that the mother attended at least one ANC visit (region-wise regressions). The estimated model corresponds to the specification (2) from table 2 & 3. All regressions include PSU fixed effects and the full set of covariates.

Figure A2: Associations between nutritional outcomes and ANC attendance by world region



Figures show marginal effects and 95% confidence intervals for binary indicator that the mother attended at least one ANC visit (region-wise regressions). The estimated model corresponds to the specification (2) from table 2 & 3. All regressions include PSU fixed effects and the full set of covariates.

Peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Page #
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	2-3
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	4-5
Objectives	3	State specific objectives, including any prespecified hypotheses	5
Methods			
Study design	4	Present key elements of study design early in the paper	6
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	6
Participants	6	(a) <i>Cohort study</i> —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up <i>Case-control study</i> —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls <i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	6
		(b) <i>Cohort study</i> —For matched studies, give matching criteria and number of exposed and unexposed <i>Case-control study</i> —For matched studies, give matching criteria and the number of controls per case	
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	6-7
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	6
Bias	9	Describe any efforts to address potential sources of bias	7-8
Study size	10	Explain how the study size was arrived at	8
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why	7-8
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding	7-8
		(b) Describe any methods used to examine subgroups and interactions	8
		(c) Explain how missing data were addressed	6,8
		(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed <i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed <i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	8
		(e) Describe any sensitivity analyses	

Continued on next page

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Results		Page #
Participants	13* (a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram	8 Figure 1
Descriptive data	14* (a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders (b) Indicate number of participants with missing data for each variable of interest (c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	9 (table A1)
Outcome data	15* <i>Cohort study</i> —Report numbers of outcome events or summary measures over time <i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure <i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	9
Main results	16 (a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	9-10
Other analyses	17 Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	9-10
Discussion		
Key results	18 Summarise key results with reference to study objectives	11
Limitations	19 Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	12-13
Interpretation	20 Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	12-13
Generalisability	21 Discuss the generalisability (external validity) of the study results	
Other information		
Funding	22 Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	14

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.