

Solvent transcytosis			Inhibitors		Inhibitor transcytosis			One-way ANOVA <i>Non-parametric, Kruskal-Wallis</i>		One-way ANOVA <i>Parametric</i>	
Median \pm s.d.	<i>n</i>	CV	Inhibitor	Conc.	Median \pm s.d.	<i>n</i>	CV	Mean rank diff.	<i>P</i>	Mean diff.	<i>P</i>
$7.9 \pm 3 \times 10^4$	40	41%	Brefeldin A	$5 \mu\text{g mL}^{-1}$	$2.3 \pm 2.2 \times 10^4$	8	73%	27.36	0.003	4.4×10^4	0.0004
$7.9 \pm 3 \times 10^4$	40	41%	Wortmannin	100 nM	$8.1 \pm 2.9 \times 10^4$	8	36%	-5.14	>0.99	-4.5×10^3	0.99
$7.9 \pm 3 \times 10^4$	40	41%	Bafilomycin	0.5 μM	$1 \pm .19 \times 10^5$	3	21%	-13.58	>0.99	-1.9×10^4	0.67
$7.9 \pm 3 \times 10^4$	40	41%	Chloroquine	100 μM	$5.3 \pm 2.7 \times 10^4$	4	42%	4.93	>0.99	1.1×10^4	0.90
$7.9 \pm 3 \times 10^4$	40	41%	W-7	100 μM	$8.2 \pm 1.1 \times 10^4$	8	14%	-6.95	>0.99	6.6×10^3	0.95