

Kidney Androgen-Regulated Protein (KAP) Transgenic Mice Are Protected Against High-Fat Diet Induced Metabolic Syndrome

Beatriz Bardaji de Quixano¹, Josep A. Villena², Miguel Aranda¹, Gemma Brils¹, Antoni Cuevas¹, Théana Hespel¹, Haizea Lekuona¹, Cristina Suárez¹, Olga Tornavaca¹, Anna Meseguer^{1,3,4,5}

¹Fisiopatologia Renal. Centre d'Investigacions en Bioquímica i Biologia Molecular (CIBBIM). Institut de Recerca Vall d'Hebron (VHIR), Barcelona, Spain.

²Laboratory of Metabolism and Obesity. Vall d'Hebron Institut de Recerca (VHIR), Barcelona, Spain.

³Departament de Bioquímica i Biologia Molecular. Unitat de Bioquímica de Medicina. Universitat Autònoma de Barcelona. Bellaterra (Barcelona) Spain.

⁴Instituto Reina Sofía de Investigación Nefrológica. Fundación Renal Íñigo Álvarez de Toledo. Spain.

⁵Red de Investigación Renal (REDINREN), Instituto Carlos III-FEDER, Madrid, Spain.

Address for correspondence:

Anna Meseguer, Ph.D.

Fisiopatologia Renal. CIBBIM.VHIR.

Pg. Vall d'Hebron 119-129. 08035 Barcelona. SPAIN.

Tel: 34-934894061.

FAX: 34-934894015.

E-mail: ana.meseguer@vhir.org

Short Title:

KAP and metabolic syndrome




Figure S1. Hemodynamic analyses

Evening tail-cuff measurements of systolic (SAP) diastolic (DAP) and mean arterial pressure (MAP) in Tg and control littermate mice fed on standard control diet (chow) or high fat diet (HFD), at 6-month of age (left panel). Mean and SEM from one experiment (n=8 per group) are represented. Statistical analysis on AUC ** p<0.01 control vs Tg, ## p<0.01 chow vs HFD. Heart rate measurements in the four groups of mice (right panel). Error bars correspond to SEM.