

Table S6. Biomarker levels in the sub-group of donors without AKI but with histological diagnosis of ATI (Subclinical ATI)

Biomarker N=326	No ATI N=239	Mild ATI N=59	Severe ATI N=28	P-value[^]
NGAL (ng/mL)	26.1 (9.8, 75.8)	36.6 (9.4, 128.4)	79.5 (22.7, 209.5)	0.02
L-FABP (ng/mL)	9.0 (3.2, 34.8)	11.8 (2.8, 56.4)	25.6 (6.4, 64.0)	0.23
IL-18 (pg/mL)	41.0 (20.0, 100.2)	40.5 (18.9, 84.6)	59.9 (25.8, 98.7)	0.40
KIM-1 (ng/mL)	1.4 (0.7, 3.1)	1.5 (0.9, 3.0)	1.4 (0.9, 4.5)	0.42

This is the subgroup of donors that excludes those with terminal serum creatinine ≥ 1.5 mg/dl. AKI defined as SCr rise ≥ 0.3 mg/dl or $\geq 50\%$ during hospitalization. Values are median (IQR). NGAL, neutrophil gelatinase-associated lipocalin; L-FABP, liver-fatty acid binding protein; IL-18, interleukin-18; KIM-1, kidney injury molecule 1; ATI, Acute Tubular Injury. Severe ATI includes moderate and severe ATI. [^]Anova test