

Supplement

Figure S1: Distribution of mean consultation duration in each practice for face-to-face or telephone consultations conducted by a GP or nurse




Table S1: Characteristics of practices according to mean duration of consultation with a GP

| | Practice mean consultation duration (minutes) (N=316) | | | | | |
|---|---|---------------------|--------------------|--------------------|--------------------|----------------|
| | <5 | ≥5 and <8 | ≥8 and <10 | ≥10 and <12 | ≥12 and <15 | ≥15 |
| Number of practices (%) | 4 (1.3) | 96 (30.4) | 118 (37.3) | 83 (26.3) | 14 (4.4) | 1 (0.3) |
| Mean list size (SD) | 5,580.0 (713.0) | 10,149.3 (4,603.6) | 10,232.1 (4809.5) | 8,771.6 (4,590.9) | 7,715.2 (2,912.3) | 9,220.0 (-) |
| Training practice | | | | | | |
| Yes: N (%) | 0 (0.0) | 33 (34.4) | 53 (44.9) | 35 (42.2) | 5 (35.7) | 0 (0.0) |
| Rurality | | | | | | |
| Not rural (Urban >10K - less sparse):N (%) | 4 (100.0) | 82 (85.4) | 101 (85.6) | 68 (81.9) | 12 (85.7) | 0 (0.0) |
| Mean GP consultation rate (per 10,000 person years) (SD) | 46,877.5 (17,478.6) | 42,488.2 (16,711.8) | 37,671.8 (9,954.9) | 32,618.1 (9,227.8) | 28,463.9 (9,387.5) | 13,895.5 (-) |
| Mean nurse consultation rate (per 10,000 person years) (SD) | 12,777.7 (4,657.8) | 14,270.9 (7,893.9) | 13,566.1 (7,677.1) | 11,707.4 (6,888.7) | 12,279.2 (8,778.4) | 11,124.7 (-) |
| Number of FTE GPs | | | | | | |
| ≤2: N (%) | 2 (50.0) | 9 (9.4) | 16 (13.6) | 14 (16.9) | 3 (21.4) | 0 (0.0) |
| >2 and ≤4: N (%) | 2 (50.0) | 24 (25.0) | 22 (18.6) | 24 (28.9) | 2 (14.3) | 0 (0.0) |
| >4 and ≤6: N (%) | 0 (0.0) | 37 (38.5) | 32 (27.1) | 24 (28.9) | 7 (50.0) | 1 (100.0) |
| >6 and ≤8: N (%) | 0 (0.0) | 15 (15.6) | 25 (21.2) | 15 (18.1) | 0 (0.0) | 0 (0.0) |
| >8 and ≤19: N (%) | 0 (0.0) | 11 (11.5) | 22 (18.6) | 5 (6.0) | 2 (14.3) | 0 (0.0) |
| Number of FTE nurses | | | | | | |
| ≤2: N (%) | 4 (100.0) | 51 (53.1) | 69 (58.5) | 52 (62.7) | 12 (85.7) | 0 (0.0) |
| >2 and ≤4: N (%) | 0 (0.0) | 22 (22.9) | 22 (18.6) | 18 (21.7) | 2 (14.3) | 1 (100.0) |
| >4 and ≤6: N (%) | 0 (0.0) | 8 (8.3) | 10 (8.5) | 2 (2.4) | 0 (0.0) | 0 (0.0) |
| >6 and ≤8: N (%) | 0 (0.0) | 4 (4.2) | 2 (1.7) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| >8 and ≤19: N (%) | 0 (0.0) | 2 (2.1) | 2 (1.7) | 0 (0.0) | 0 (0.0) | 0 (0.0) |
| QOF performance | | | | | | |
| 1 st quintile (poorest performance): N (%) | 2 (50.0) | 15 (15.6) | 18 (15.3) | 12 (14.5) | 3 (21.4) | 0 (0.0) |
| 2 nd quintile: N (%) | 0 (0.0) | 22 (22.9) | 13 (11.0) | 13 (15.7) | 1 (7.1) | 0 (0.0) |
| 3 rd quintile: N (%) | 1 (25.0) | 23 (24.0) | 24 (20.3) | 8 (9.6) | 2 (14.3) | 1 (100.0) |
| 4 th quintile: N (%) | 1 (25.0) | 18 (18.8) | 31 (26.3) | 27 (32.5) | 5 (35.7) | 0 (0.0) |
| 5 th quintile (best performance): N (%) | 0 (0.0) | 18 (18.8) | 30 (25.4) | 22 (26.5) | 3 (21.4) | 0 (0.0) |

Table S2: Full model for consultations with a GP

| | Change in duration (seconds) | p-value | 95% confidence interval | |
|--|------------------------------|---------|-------------------------|---------|
| Female gender (Male = reference) | 8.29 | 0.000 | 6.03 | 10.55 |
| Ethnic group (White = reference) | | | | |
| Asian | 4.11 | 0.231 | -2.61 | 10.84 |
| Chinese | -6.32 | 0.607 | -30.41 | 17.76 |
| Black | -5.63 | 0.205 | -14.34 | 3.07 |
| Mixed/ Other | 4.35 | 0.283 | -3.59 | 12.29 |
| Unknown | -11.00 | 0.000 | -13.50 | -8.49 |
| Index of multiple deprivation (1 st quintile = reference) | | | | |
| 2 nd quintile | 1.14 | 0.535 | -2.47 | 4.76 |
| 3 rd quintile | -2.39 | 0.235 | -6.33 | 1.55 |
| 4 th quintile | -3.57 | 0.095 | -7.77 | 0.62 |
| 5 th quintile (most deprived) | -5.04 | 0.037 | -9.77 | -0.30 |
| Unknown | -11.27 | 0.060 | -23.02 | 0.48 |
| Smoking status (Non-smoker = reference) | | | | |
| Current smoker | -2.36 | 0.147 | -5.54 | 0.83 |
| Ex-smoker | 0.11 | 0.944 | -2.91 | 3.13 |
| Unknown | 18.66 | 0.000 | 14.58 | 22.75 |
| Age group (0-14 years = reference) | | | | |
| 15-24 | 55.71 | 0.000 | 50.12 | 61.30 |
| 25-44 | 83.69 | 0.000 | 78.72 | 88.65 |
| 45-64 | 89.83 | 0.000 | 84.77 | 94.89 |
| 65-74 | 65.84 | 0.000 | 60.25 | 71.42 |
| 75+ | 58.44 | 0.000 | 52.96 | 63.93 |
| Telephone consultation (Face-to-face = reference) | -308.70 | 0.000 | -311.64 | -305.76 |
| Practice list size (centred, per 1000) | -5.21 | 0.026 | -9.81 | -0.61 |
| Number of FTE GPs (<=2 = reference) | | | | |
| >2 and <=4 | 27.74 | 0.206 | -15.23 | 70.70 |
| >4 and <=6 | 56.90 | 0.016 | 10.62 | 103.18 |
| >6 and <=8 | 58.06 | 0.061 | -2.61 | 118.73 |
| >8 and <=19 | 120.02 | 0.002 | 43.58 | 196.47 |
| Unknown | 211.75 | 0.120 | -54.93 | 478.44 |
| Number of FTE nurses (<=2 = reference) | | | | |
| >2 and <=4 | -10.07 | 0.568 | -44.59 | 24.46 |
| >4 and <=6 | -36.07 | 0.239 | -96.11 | 23.97 |
| >6 and <=8 | -141.57 | 0.003 | -235.44 | -47.70 |
| >8 and <=19 | 54.83 | 0.354 | -61.14 | 170.81 |
| Unknown | -16.52 | 0.429 | -57.41 | 24.37 |
| Training practice (No = reference) | | | | |
| Yes | 45.61 | 0.001 | 18.26 | 72.96 |
| Unknown | - | | | |
| Nurse consultation rate (centred, per 1000 per 10,000 person years) | 0.40 | 0.658 | -1.36 | 2.15 |
| GP consultation rate (centred, per 1000 per 10,000 person years) | -3.83 | 0.000 | -4.80 | -2.85 |
| QOF performance (1 st quintile = reference) | | | | |
| 2 nd quintile | 3.50 | 0.869 | -38.00 | 44.99 |
| 3 rd quintile | -16.22 | 0.427 | -56.19 | 23.76 |
| 4 th quintile | 27.47 | 0.147 | -9.67 | 64.61 |
| 5 th quintile (best performance) | 18.21 | 0.343 | -19.40 | 55.83 |
| Unknown | -50.43 | 0.633 | -257.64 | 156.78 |
| Rural practice (Urban = reference) | 17.21 | 0.309 | -15.98 | 50.39 |
| Mean duration | 417.27 | 0.000 | 367.50 | 467.04 |

Table S3: Final model for consultations with a GP including a term for GP registrar role (post-hoc sensitivity analysis)

| | Change in duration (seconds) | p-value | 95% confidence interval | |
|--|------------------------------|---------|-------------------------|---------|
| Female gender (Male = reference) | 7.40 | 0.000 | 5.16 | 9.63 |
| Ethnic group (White = reference) | | | | |
| Asian | 4.54 | 0.181 | -2.11 | 11.19 |
| Chinese | -7.76 | 0.523 | -31.58 | 16.07 |
| Black | -7.18 | 0.102 | -15.79 | 1.43 |
| Mixed/ Other | 3.82 | 0.340 | -4.03 | 11.68 |
| Unknown | -12.32 | 0.000 | -14.80 | -9.85 |
| Index of multiple deprivation (1 st quintile = reference) | | | | |
| 2 nd quintile | 0.52 | 0.776 | -3.06 | 4.10 |
| 3 rd quintile | -3.37 | 0.090 | -7.27 | 0.53 |
| 4 th quintile | -4.88 | 0.021 | -9.03 | -0.74 |
| 5 th quintile (most deprived) | -6.62 | 0.006 | -11.30 | -1.94 |
| Unknown | -8.24 | 0.165 | -19.87 | 3.39 |
| Smoking status (Non-smoker = reference) | | | | |
| Current smoker | -3.11 | 0.053 | -6.26 | 0.04 |
| Ex-smoker | -0.10 | 0.945 | -3.09 | 2.88 |
| Unknown | 15.56 | 0.000 | 11.52 | 19.60 |
| Age group (0-14 years = reference) | | | | |
| 15-24 | 52.35 | 0.000 | 46.82 | 57.88 |
| 25-44 | 81.88 | 0.000 | 76.97 | 86.79 |
| 45-64 | 92.15 | 0.000 | 87.14 | 97.15 |
| 65-74 | 71.73 | 0.000 | 66.21 | 77.26 |
| 75+ | 64.95 | 0.000 | 59.52 | 70.38 |
| Telephone consultation (Face-to-face = reference) | -295.45 | 0.000 | -298.37 | -292.53 |
| Training practice (No = reference) | | | | |
| Yes | 21.92 | 0.073 | -2.07 | 45.90 |
| Unknown | 133.11 | 0.107 | -28.55 | 294.77 |
| GP registrar conducting appointment (No=reference) | 245.04 | 0.000 | 241.49 | 248.59 |
| GP consultation rate (centred, per 1000 per 10,000 person years) | -3.33 | 0.000 | -4.24 | -2.41 |
| Mean duration | 460.60 | 0.000 | 443.91 | 477.30 |

Table S4: Characteristics of practices according to mean duration of consultation with a nurse

| | Practice mean consultation duration (minutes) (N=307) | | | | | |
|--|---|------------------------|------------------------|------------------------|-----------------------|------------------------|
| | <5 | ≥5 and <8 | ≥8 and <10 | ≥10 and <12 | ≥12 and <15 | ≥15 |
| Number of practices (%) | 3 (1.0) | 59 (19.2) | 103 (33.6) | 71 (23.1) | 54 (17.6) | 17 (5.5) |
| Mean list size (SD) | 8,189.0 (6,263.3) | 9,936.8 (4,101.1) | 10,029.6 (4,506.8) | 10,945.5 (5,387.0) | 8,591.0 (3,987.7) | 7,286.6 (3,564.8) |
| Training practice | | | | | | |
| Yes: N (%) | 2 (66.7) | 24 (40.7) | 43 (41.8) | 37 (52.1) | 15 (27.8) | 5 (29.4) |
| Rurality | | | | | | |
| Not rural (Urban >10K - less sparse):N (%) | 3 (100.0) | 50 (84.8) | 86 (83.5) | 59 (83.1) | 47 (87.0) | 15 (88.2) |
| Mean GP consultation rate (per 10,000 person years) (SD) | 34,558.4 (7,497.0) | 39,556.1 (17,065.8) | 39,333.7 (13,086.5) | 37,073.1 (11,166.8) | 34,315.5 (8,097.0) | 35,907.5 (13,077.5) |
| Mean nurse consultation rate (per 10,000 person years) (SD) | 15,359.3 (18,373.0) | 17,422.4 (7,482.5) | 15,404.9 (7,757.8) | 12,162.1 (5,803.8) | 9,868.6 (4,282.7) | 7,029.6 (3,980.1) |
| Number of FTE GPs | | | | | | |
| ≤2: N (%) | 1 (33.3) | 5 (8.5) | 10 (9.7) | 6 (8.5) | 12 (22.2) | 5 (29.4) |
| >2 and ≤4: N (%) | 1 (33.3) | 14 (23.7) | 28 (27.2) | 10 (14.1) | 16 (29.6) | 4 (23.5) |
| >4 and ≤6: N (%) | 0 (0.0) | 22 (37.3) | 27 (26.2) | 31 (43.4) | 13 (24.1) | 6 (35.3) |
| >6 and ≤8: N (%) | 1 (33.3) | 10 (17.0) | 22 (21.4) | 11 (15.5) | 9 (16.7) | 2 (11.8) |
| >8 and ≤19: N (%) | 0 (0.0) | 7 (11.9) | 16 (15.5) | 13 (18.3) | 4 (7.4) | 0 (0.0) |
| Number of FTE nurses | | | | | | |
| ≤2: N (%) | 2 (66.7) | 35 (59.3) | 51 (49.5) | 42 (59.2) | 38 (70.4) | 12 (70.6) |
| >2 and ≤4: N (%) | 1 (33.3) | 12 (20.3) | 28 (27.2) | 15 (21.2) | 7 (13.0) | 2 (11.8) |
| >4 and ≤6: N (%) | 0 (0.0) | 3 (5.1) | 11 (10.7) | 5 (7.0) | 1 (19.9) | 0 (0.0) |
| >6 and ≤8: N (%) | 0 (0.0) | 1 (1.7) | 0 (0.0) | 4 (5.6) | 1 (1.9) | 0 (0.0) |
| >8 and ≤19: N (%) | 0 (0.0) | 1 (1.7) | 2 (1.9) | 1 (1.4) | 0 (0.0) | 0 (0.0) |
| QOF performance | | | | | | |
| 1 st quintile (poorest performance): N (%) | 1 (33.3) | 9 (15.3) | 15 (14.6) | 13 (18.3) | 7 (13.0) | 2 (11.8) |
| 2 nd quintile: N (%) | 0 (0.0) | 5 (8.5) | 17 (16.5) | 13 (18.3) | 10 (18.5) | 1 (5.9) |
| 3 rd quintile: N (%) | 0 (0.0) | 9 (15.3) | 24 (23.3) | 13 (18.3) | 7 (13.0) | 6 (35.3) |
| 4 th quintile: N (%) | 1 (33.3) | 17 (28.8) | 28 (27.2) | 20 (28.2) | 11 (20.4) | 4 (23.5) |
| 5 th quintile (best performance): N (%) | 1 (33.3) | 17 (28.8) | 19 (18.5) | 12 (16.9) | 19 (35.2) | 4 (23.5) |

Table S5: Full model for consultations with a nurse

| | Change in duration (seconds) | p-value | 95% confidence interval | |
|--|------------------------------|---------|-------------------------|---------|
| Female gender (Male = reference) | -11.15 | 0.000 | -15.34 | -6.96 |
| Ethnic group (White = reference) | | | | |
| Asian | 0.18 | 0.979 | -12.99 | 13.35 |
| Chinese | -17.82 | 0.445 | -63.57 | 27.93 |
| Black | -1.67 | 0.849 | -18.84 | 15.50 |
| Mixed/ Other | -5.52 | 0.489 | -21.18 | 10.13 |
| Unknown | -2.52 | 0.279 | -7.09 | 2.04 |
| Index of multiple deprivation (1 st quintile = reference) | | | | |
| 2 nd quintile | 7.49 | 0.026 | 0.92 | 14.06 |
| 3 rd quintile | -0.54 | 0.883 | -7.70 | 6.62 |
| 4 th quintile | 5.38 | 0.167 | -2.24 | 13.01 |
| 5 th quintile (most deprived) | 7.92 | 0.071 | -0.67 | 16.51 |
| Unknown | -25.47 | 0.011 | -45.10 | -5.84 |
| Smoking status (Non-smoker = reference) | | | | |
| Current smoker | 26.40 | 0.000 | 20.49 | 32.31 |
| Ex-smoker | 15.00 | 0.000 | 9.59 | 20.42 |
| Unknown | 21.12 | 0.000 | 13.23 | 29.01 |
| Age group (0-14 years = reference) | | | | |
| 15-24 | 52.88 | 0.000 | 42.16 | 63.60 |
| 25-44 | 72.19 | 0.000 | 62.83 | 81.55 |
| 45-64 | 113.49 | 0.000 | 103.97 | 123.00 |
| 65-74 | 73.96 | 0.000 | 63.81 | 84.10 |
| 75+ | 75.54 | 0.000 | 65.43 | 85.65 |
| Telephone consultation (Face-to-face = reference) | -279.51 | 0.000 | -288.35 | -270.66 |
| Practice list size (centred, per 1000) | -6.09 | 0.065 | -12.56 | 0.38 |
| Number of FTE GPs (<=2 = reference) | | | | |
| >2 and <=4 | -63.99 | 0.045 | -126.58 | -1.39 |
| >4 and <=6 | -49.06 | 0.150 | -115.82 | 17.70 |
| >6 and <=8 | -33.73 | 0.442 | -119.68 | 52.22 |
| >8 and <=19 | -29.10 | 0.596 | -136.61 | 78.42 |
| Unknown | -106.02 | 0.625 | -531.17 | 319.12 |
| Number of FTE nurses (<=2 = reference) | | | | |
| >2 and <=4 | 20.31 | 0.408 | -27.79 | 68.41 |
| >4 and <=6 | 49.99 | 0.240 | -33.47 | 133.46 |
| >6 and <=8 | 113.35 | 0.088 | -16.77 | 243.48 |
| >8 and <=19 | 137.25 | 0.096 | -24.40 | 298.89 |
| Unknown | -7.81 | 0.791 | -65.58 | 49.97 |
| Training practice (No = reference) | | | | |
| Yes | -11.09 | 0.570 | -49.31 | 27.14 |
| Unknown | - | | | |
| Nurse consultation rate, centred, per 1000 per 10,000 person years) | -10.23 | 0.000 | -12.84 | -7.62 |
| GP consultation rate, centred, per 1000 per 10,000 person years) | -0.49 | 0.488 | -1.88 | 0.90 |
| QOF performance (1 st quintile = reference) | | | | |
| 2 nd quintile | 28.82 | 0.345 | -30.93 | 88.56 |
| 3 rd quintile | 6.08 | 0.835 | -50.99 | 63.14 |
| 4 th quintile | 9.11 | 0.738 | -44.24 | 62.47 |
| 5 th quintile (best performance) | 31.22 | 0.254 | -22.44 | 84.89 |
| Unknown | -81.78 | 0.575 | -367.98 | 204.42 |
| Rural practice (Urban = reference) | 13.93 | 0.561 | -33.09 | 60.96 |
| Mean duration | 543.41 | 0.000 | 470.92 | 615.90 |