

Low resistance to chytridiomycosis in direct-developing amphibians

**Andréa F. C. Mesquita¹, Carolina Lambertini², Mariana Lyra¹,
Leo R. Malagoli¹, Timothy Y. James³, Luís Felipe Toledo²,
Célio F. B. Haddad¹, C. Guilherme Becker*⁴**

¹ Universidade Estadual Paulista (UNESP), Instituto de Biociências, Departamento de Zoologia, and Centro de Aquicultura (CAUNESP), 13506-900 Rio Claro, SP, Brazil

² Departamento de Biologia Animal, I.B., Universidade Estadual de Campinas, 13083-862 Campinas, SP, Brazil

³ Department of Ecology and Evolutionary Biology, University of Michigan, 48109, Ann Arbor, MI, U.S.A.

⁴ Department of Biological Sciences, The University of Alabama, 35487 Tuscaloosa, AL, U.S.A.

Corresponding author:

C. Guilherme Becker

*cgbecker1@ua.edu

Supplementary Table S1

Table S1. GLMM testing the effect of host life history on gain in *Bd* load during the experiment. Day of swab included as fixed effect and species included as random effect.

Parameter	d.f. (den)	t	P
Intercept	22.73	1.23	0.231
Life history	3.925	-4.3	0.013*
Day of swab	53.55	1.99	0.051

Random effect of species accounted for 47.32%

Supplementary Photo S1



Photo S1. Specimen of the endemic direct-developing frog *Holoaden bradei* (Craugastoridae), last seen in the wild in 1978 and presumably extinct due to chytridiomycosis¹⁸. Die-offs were never seen at the high elevation quotas of Parque Nacional do Itatiaia, Brazil's Atlantic forest.