

Plant-endophytes interaction influences the secondary metabolism in *Echinacea purpurea* (L.)

Moench: an *in vitro* model.

Valentina Maggini^{1,2,3}, Marinella De Leo⁴, Alessio Mengoni¹, Eugenia Rosaria Gallo^{2,3}, Elisangela Miceli¹, Rose Vanessa Bandeira Reidel⁴, Sauro Biffi⁵, Luisa Pistelli⁴, Renato Fani¹, Fabio Firenzuoli³ and Patrizia Bogani¹

¹Department of Biology, University of Florence, Via Madonna del Piano 6, 50019 Sesto Fiorentino, Italy; ²Department of Experimental and Clinical Medicine, University of Florence, Largo Brambilla 3, 50134 Florence, Italy; ³Referring Center for Phytotherapy, Tuscany Region, Careggi University Hospital, Largo Brambilla 3, 50134 Florence, Italy; ⁴Department of Pharmacy, University of Pisa, Via Bonanno 33, 56126 Pisa, Italy; ⁵Botanical Garden Casola Valsenio, Via del Corso 6, 48010 Ravenna, Italy

Supplementary Figure 1. HPLC-PDA/UV(detected at 260 nm) alkamide profiles from *n*-hexane extracts of control and infected *E. purpurea* stem/leaves (SL). For peak data, see Supplementary Table 3. Identified compounds are listed in Table 1.

