

Plant-endophytes interaction influences the secondary metabolism in *Echinacea purpurea* (L.)

Moench: an *in vitro* model.

Valentina Maggini^{1,2,3}, Marinella De Leo⁴, Alessio Mengoni¹, Eugenia Rosaria Gallo^{2,3}, Elisangela Miceli¹, Rose Vanessa Bandeira Reidel⁴, Sauro Biffi⁵, Luisa Pistelli⁴, Renato Fani¹, Fabio Firenzuoli³ and Patrizia Bogani¹

¹Department of Biology, University of Florence, Via Madonna del Piano 6, 50019 Sesto Fiorentino, Italy; ²Department of Experimental and Clinical Medicine, University of Florence, Largo Brambilla 3, 50134 Florence, Italy; ³Referring Center for Phytotherapy, Tuscany Region, Careggi University Hospital, Largo Brambilla 3, 50134 Florence, Italy; ⁴Department of Pharmacy, University of Pisa, Via Bonanno 33, 56126 Pisa, Italy; ⁵Botanical Garden Casola Valsenio, Via del Corso 6, 48010 Ravenna, Italy

Supplementary Table 5. Sample preparation for HPLC analysis of the alkamides in the extracts from R and SL samples (CSL: stem/leaves extract from control plants; ISL: stem/leaves extract from infected plants; CR: root extract from control plants; IR: root extract from infected plants).

Sample	Fresh weight (g)	Dry weight (g)	<i>n</i> -hexane extract (mg)
C R	2.53	0.28	7.8
I R	1.90	0.24	2.4
C SL	5.90	1.26	17.4
I SL	4.12	0.92	11.2