

S6 Table. Comparison of dialysis patients and non-dialysis CKD patients

	Dialysis (n=299)	Non-dialysis (n=732)	P value
Age (years)	62(34-80)	56(35-70)	<0.0001
Gender, male (%)	180 (60)	477(65)	0.13
Diabetes mellitus, n (%)	58 (19)	211 (29)	0.001
CVD, n (%)	136 (46)	234 (32)	<0.0001
SGA>1, n (%)	131 (44)	189(26)	<0.0001
% HGS (n=289/697)	70 (42-108)	91 (56-119)	<0.0001
BMI (kg/m ²)	24 (19 -30)	25 (20 -31)	0.003
LBMI (kg/m ² ; n=280/610)	17.0 (13.8-20.5)	17.2 (13.9-20.5)	0.14
FBMI (kg/m ² ; n=280/610)	6.9 (3.8-11.3)	7.4 (4.4-11.5)	0.02
S-Albumin (g/L)	34 (28-40)	35 (27-41)	0.008
hsCRP (mg/L)	4.1 (0.4-32)	3.1 (0.5-24)	0.09

Data presented as median (10th - 90th percentile), number or percentage.

Abbreviations: CVD, cardiovascular disease; SGA, subjective global assessment; % HGS, handgrip strength as percentage of the controls; BMI, body mass index; LBMI, lean body mass index; FBMI, fat body mass index; S-Albumin, serum-albumin; hs CRP, high sensitivity C-reactive protein