

S7 Table. Comparison of CKD patients with % HGS \geq 75 and % HGS $<$ 75

	% HGS \geq 75 (n=596)	% HGS $<$75 (n=389)	P value
Age (years)	53(32-70)	64(44-79)	<0.0001
Gender, male (%)	398 (67)	233 (60)	0.03
Diabetes mellitus, n (%)	111 (19)	138 (35)	<0.0001
CVD, n (%)	135 (23)	216 (56)	<0.0001
Dialysis, n (%)	119(20)	170(44)	<0.0001
SGA$>$1, n (%)	105 (18)	192(49)	<0.0001
BMI (kg/m²)	24.9(20.5-30.6)	23.8(19.0-30.3)	0.0002
LBMI (kg/m²; n=528/346)	17.5 (14.5-20.9)	16.4 (13.6-19.8)	<0.0001
FBMI (kg/m²; n=528/346)	7.1 (4.1-11.5)	7.4 (4.1-11.2)	0.61
S-Albumin (g/L)	36(29-41)	33(26-39)	<0.0001
hsCRP (mg/L)	2.3(0.4-16.2)	6.3(0.8-35)	<0.0001

Data presented as median (10th - 90th percentile), number or percentage.

Abbreviations: % HGS, handgrip strength as percentage of the controls; CVD, cardiovascular disease; SGA, subjective global assessment; BMI, body mass index; LBMI, lean body mass index; FBMI, fat body mass index; S-Albumin, serum-albumin; hsCRP, high sensitivity C-reactive protein