

S12 Table. Comparison of CKD patients with inflammation (hsCRP \geq 10 mg/L) and without inflammation (hsCRP <10 mg/L).

	hsCRP<10 mg/L (n=773)	hsCRP \geq 10 mg/L (n=258)	P value
Age (years)	56(32-74)	61(43-75)	<0.0001
Gender, male (%)	486 (63)	171 (66)	0.32
Diabetes mellitus, n (%)	186 (24)	83 (32)	0.01
CVD, n (%)	224 (29)	146 (57)	<0.0001
Dialysis, n (%)	85(33)	214(28)	0.11
SGA>1, n (%)	189 (24)	131(51)	<0.0001
%HGS (n=744/241)	89(56-119)	70(42-103)	<0.0001
BMI (kg/m²)	24.6(19.9-30.4)	24.4(19.5-31.2)	1.00
LBMI (kg/m²; n=668/222)	17.3 (13.8-20.6)	16.8 (14.1-20.2)	0.08
FBMI (kg/m²; n=668/222)	7.1 (4.1-11.0)	7.6 (4.1-12.4)	0.10
S-Albumin (g/L)	36(29-41)	32(25-37)	<0.0001

Data presented as median (10th - 90th percentile), number or percentage.

Abbreviations: hsCRP, high sensitivity C-reactive protein; CVD, cardiovascular disease; SGA, subjective global assessment; % HGS, handgrip strength as percentage of the controls; BMI, body mass index; LBMI, lean body mass index; FBMI, fat body mass index; S-Albumin, serum-albumin