

SUPPLEMENTAL MATERIAL

Table S1. Independent all-cause mortality risk predictors in non-diabetic patients with stable CAD.

	Adjusted HR	95% CI	P-Value
IFG vs. no IFG	1·22	1·14 – 1·31	<0·001
BMI > 30 kg/m ²	1·19	1·02 – 1·24	0·01
Low HDL [#]	1·08	0·99 – 1·20	0·09
Elevated BP [@]	1·02	0·95 – 1·10	0·55
TG > 150 mg/dL	1·03	0·96 – 1·11	0·40
BMI > 30 kg/m ²	1·19	1·02 – 1·24	0·01
HTN	1·09	1·01 – 1·17	0·03
Past MI	1·46	1·34 – 1·60	<0·001
NYHA ≥ 3	1·19	1·01 – 1·39	0·03
Creatinine > 1·5 mg/dL	1·60	1·36 – 1·88	<0·001
Age > 60 years	1·08	1·07 – 1·09	<0·001
Male sex	1·21	1·07 – 1·35	0·001
COPD	1·27	1·05 – 1·54	0·01
Past CVA	1·19	0·86 – 1·65	0·30

BMI = body mass index; BP = blood pressure; COPD = chronic obstructive pulmonary disease; CVA = cerebral vascular accident; HDL = high-density lipoprotein; HTN = hypertension; IFG = impaired fasting glucose; MI = myocardial infarction; NYHA = New York Heart Association; TG = triglycerides.

[#] Low HDL defined as HDL < 40 mg/dL in males and HDL < 50 mg/dL in females.

[@] Systolic blood-pressure > 130 (mmHg) or/and diastolic blood-pressure > 85 (mmHg)