

Supplementary Table 3 Maternal health and lifestyle factors, sociodemographic characteristics, and sick leave categorization according to the country of residence among pregnant women and new mothers.

Independent variables	Total population, n=6686	Pregnant women, n=3486	New mothers, n=3200	Pregnant women vs new mothers
Health and lifestyle factors	n (%)	n (%)	n (%)	p-value
Alcohol in pregnancy*				
Yes	1069 (16.0)	494 (14.2)	575 (18.0)	<0.001
No	5562 (83.2)	2958 (84.9)	2604 (81.4)	
Smoking in pregnancy*				
Yes	539 (8.1)	268 (7.7)	271 (8.5)	0.243
No	6133 (91.7)	3210 (92.1)	2923 (91.3)	
Pregnant after infertility treatment				
Yes	476 (7.1)	269 (7.7)	207 (6.5)	0.047
No	6210 (92.9)	3217 (92.3)	2993 (93.5)	
Folic acid use*				
Yes	6151 (92.6)	3199 (92.4)	2952 (92.9)	0.449
No	489 (7.4)	263 (7.6)	226 (7.1)	
Medication for chronic indications				
Yes	1371 (20.5)	720 (20.7)	651 (20.3)	0.754
No	5315 (79.5)	2766 (79.4)	2549 (79.7)	
Medication for acute indications				<0.001
Yes	5254 (78.6)	2610 (74.9)	2644 (82.6)	
No	1432 (21.4)	876 (25.1)	556 (17.4)	
Number of acute conditions				
Mean (SD), range 0-10	4.2 (1.6)	4.2 (1.6)	4.1 (1.6)	<0.001
Chronic conditions**				
No chronic conditions	5279 (79.0)	2762 (79.2)	2517 (78.7)	0.828
Somatic conditions	1167 (17.5)	599 (17.2)	568 (17.8)	
Mental conditions	240 (3.6)	125 (3.6)	115 (3.6)	
Country of residence				
Sick leave policy***				
High	1179 (17.6)	674 (19.3)	505 (15.8)	<0.001
Medium	3128 (46.8)	1536 (44.1)	1592 (49.8)	
Low	2379 (35.6)	1276 (36.6)	1103 (34.5)	
Region				
Northern Europe	2351 (35.2)	1349 (38.7)	1002 (31.3)	<0.001
Eastern Europe	1956 (29.3)	861 (24.7)	1095 (34.2)	
Western Europe	2379 (35.6)	1276 (36.3)	1103 (34.5)	
Sociodemographic and maternal characteristics				
Maternal age (years)				
≤20	168 (2.5)	112 (3.2)	56 (1.8)	<0.001
21-30	3704 (55.4)	1974 (56.6)	1730 (54.1)	
31-40	2698 (40.4)	1351 (38.8)	1347 (42.1)	
≥41	116 (1.7)	49 (1.4)	67 (2.1)	
Marital status				
Married/cohabitant	6375 (95.4)	3333 (95.6)	3042 (95.1)	0.287
Single/divorced/other	311 (4.7)	153 (4.4)	158 (4.9)	

Employment*				
Student	695 (10.4)	342 (9.8)	353 (11.0)	0.014
Healthcare personnel	1088 (16.3)	607 (17.4)	481 (15.0)	
Employed in other sectors	4893 (73.2)	2530 (72.6)	2363 (73.8)	
Highest level of education				
Primary school	191 (2.9)	96 (2.8)	95 (2.9)	0.858
High-school	1736 (26.0)	916 (26.3)	820 (25.6)	
University or collage	3985 (59.6)	2077 (59.6)	1908 (59.6)	
Other education	774 (11.6)	397 (11.4)	377 (11.8)	
Primiparity				
Yes	3603 (53.9)	2085 (59.8)	1518 (47.4)	<0.001
No	3083 (46.1)	1401 (40.2)	1682 (52.6)	
Pregnancy planned*				
Yes	4657 (69.7)	2443 (70.1)	2214 (69.2)	0.076
No, but expected	1491 (22.3)	788 (22.6)	703 (22.0)	
No	519 (7.8)	246 (7.1)	273 (8.5)	

*Total numbers do not add up due to missing values: Folic acid n=46 (0.7%), employment n=10 (0.2%), alcohol in pregnancy n=55 (0.8%), smoking in pregnancy n=14 (0.2%), and pregnancy planned n=19 (0.3%).

** Chronic disorders were categorized as no conditions, somatic conditions only (allergy, asthma, diabetes, hypothyroidism, cardiovascular diseases, and rheumatic disorders), and any mood disorders (anxiety and/or depression), i.e. women with both somatic and mood disorders were categorized in the latter group.

*** **High:** Norway, **Medium:** Croatia, Finland, Poland, Russia, Serbia, Slovenia, and Sweden, and **Low:** Italy, France, the United Kingdom, and Switzerland.