

BMJ Open

The STOP-AB Trial protocol: Safety of discontinuing patient antibiotic treatment when physicians no longer consider it necessary.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2016-015814
Article Type:	Protocol
Date Submitted by the Author:	02-Jan-2017
Complete List of Authors:	Llor, Carl; Primary healthcare centre Barcelona-2B (Via Roma) Moragas, Ana; Universitat Rovira i Virgili, Primary Healthcare Centre Jaume I Bayona, Carolina; Primary Healthcare Centre La Marina Cots, Josep M.; Universitat de Barcelona, Primary Healthcare Centre La Marina Molero, José M. ; Primary Healthcare Centre San Andrés Ribas, Joana; Primary Healthcare Centre Hospital Son Llätzer Fóthy, Julio Francisco ; Primary Healthcare Centre Coll d'en Rabassa Gutiérrez, Isabel ; Primary Healthcare Centre Rondilla I Sánchez, Coro ; Primary Care Centre Virxe da Peregrina Ortega, Jesús; Primary Healthcare Centre Rincón de Soto Arranz, Javier; Primary Healthcare Centre Arquitecte Bennàssar Botanes, Jenifer ; Primary Healthcare Centre Cornellà de Llobregat Robles, Purificación; Primary Care Centre Can Vidalet
Primary Subject Heading:	General practice / Family practice
Secondary Subject Heading:	Infectious diseases
Keywords:	PRIMARY CARE, Clinical trials < THERAPEUTICS, Respiratory infections < THORACIC MEDICINE, Public health < INFECTIOUS DISEASES

SCHOLARONE™
Manuscripts

Title

The STOP-AB Trial protocol: Safety of discontinuing patient antibiotic treatment when physicians no longer consider it necessary.

Authors

Carl Llor¹, Ana Moragas², Carolina Bayona³, Josep M. Cots⁴, José M. Molero⁵, Joana Ribas⁶, Julio Francisco Fóthy⁷, Isabel Gutiérrez⁸, Coro Sánchez⁹, Jesús Ortega¹⁰, Javier Arranz¹¹, Jenifer Botanes¹², Purificación Robles¹³

¹Primary Healthcare Centre Barcelona-2B (Via Roma), Spain

²Universitat Rovira i Virgili. Primary Healthcare Centre Jaume I, Tarragona, Spain

³Primary Healthcare Centre La Marina, Barcelona, Spain

⁴Universitat of Barcelona. Primary Healthcare Centre La Marina, Barcelona, Spain

⁵Primary Healthcare Centre San Andrés, Madrid, Spain

⁶Primary Healthcare Centre Hospital Son Llàtzer, Palma, Spain

⁷Primary Healthcare Centre Coll d'en Rabassa, Palma, Spain

⁸Primary Healthcare Centre Rondilla I, Valladolid, Spain

⁹Primary Care Centre Virxe da Peregrina, Pontevedra, Spain

¹⁰Primary Healthcare Centre Rincón de Soto, Rincón de Soto, Spain

¹¹Primary Healthcare Centre Arquitecte Bennàssar, Palma, Spain

¹²Primary Healthcare Centre Cornellà de Llobregat, Barcelona, Spain

¹³Primary Care Centre Can Vidalet, Esplugues de Llobregat, Spain

Address for correspondence

Carl Llor

Primary Healthcare Centre Barcelona-2B (Via Roma)

1
2
3 c. Manso 19, 3rd floor

4
5 08015 Barcelona, Catalonia, Spain

6
7 Email: carles.llor@urv.cat
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

ABSTRACT

Introduction: Since 2011 the Spanish Society of Family Medicine has been recommending GPs to ask their patients to stop taking antibiotics when they suspect a viral infection. However, there is no evidence that discontinuing antibiotic therapy is safe. The main objective of this study is to determine whether discontinuation of antibiotic therapy when a GP no longer considers it necessary makes any difference in terms of the number of days with severe symptoms.

Methods and analysis: This is a multicentre, open-label, randomised controlled clinical trial. The study will be conducted in twenty primary care centres in Spain. We will include patients from 18 to 75 years of age with uncomplicated acute respiratory tract infections (RTIs) who have previously taken any dose of antibiotic but physicians no longer consider it necessary. The patients will be randomly assigned to the usual strategy of continuing antibiotic treatment (control group) or discontinuing antibiotic therapy (intervention group). A sample size of 215 patients per group was calculated on the basis of a reduction of one day in the duration of severe symptoms as a clinically relevant outcome. The primary outcome will be duration of severe symptoms, i.e. symptoms scored 5 or 6 by means of a symptom diary. Secondary outcomes will include: antibiotics taken, adverse events, patient satisfaction, and complications within the first 3 months. A post-trial observational study by means of a qualitative analysis is planned to be carried out after the clinical trial to know the percentage of the use of the strategy of discontinuing antibiotic treatment and the pros and cons of its use.

Ethics and dissemination: The study was approved by the Ethical Board of *Fundació Jordi Gol i Gurina* (reference number: 16/093). The findings of this trial will be disseminated through research conferences and peer-reviewed journals.

Trial registration number: NCT02900820.

Strengths and limitations of this study

- Some patients might not complete the symptom diaries. However, strategies to improve diary return rates will be implemented and non-respondents will be telephoned.
- Most patients will be diagnosed of rhinosinusitis, pharyngitis and bronchitis, limiting the inferences for patients with influenza and those with exacerbations of chronic bronchitis and mild-to-moderate COPD.
- It could be argued that the open nature of the study may cause a placebo effect favouring antibiotics. However, this effect will be minimised by the similar structured information all patients will receive about the self-limiting nature of respiratory tract infections and the advice about non-antibiotic medication use.
- Another possible limitation of this study is the fact that microbiological studies will not be taken into account. In primary care the response to treatment is mainly judged with clinical rather than microbiological criteria.
- The open design will allow us to study the perceptions of patients in a situation similarly to usual practice. The strengths of our study are its pragmatic design and that our study, as far as we know, is the first trial to assess if discontinuation of antibiotic therapy when a GP no longer considers it necessary makes any difference in terms of the number of days with severe symptoms in a randomised fashion.

BACKGROUND

Acute respiratory tract infections (RTIs) are among the most common reasons for a healthcare encounter in Western countries, accounting for about 15% of all visits [1]. Most episodes are caused by viruses, and in otherwise healthy adults these infections are typically self-limited and do not require a visit to a physician or a prescription medication.

Nevertheless, many patients with uncomplicated, self-limited RTIs seek care in the primary care offices. A recent study found that 72% of the primary care visits due to an acute respiratory tract infection did not seem to require an office visit [2], subsequently increasing healthcare costs and often leading to inappropriate antibiotic prescription [3,4]. Very importantly, inappropriate overuse of antibiotic medications can be very detrimental and may lead to antibiotic resistance [5,6], which has a tremendous impact on the economy [7], and patients often experience adverse effects such as diarrhoea, thrush, nausea, urticaria, and rash, which give rise to further office visits and time off work. Despite being uncommon, patients might also experience serious complications such as anaphylaxis and *Clostridium difficile* infection [8]. In addition, medicalisation of a self-limited condition makes it more likely for patients to visit a healthcare provider the next time they have a similar episode [9].

General practitioners (GP) have always been told to continue an antibiotic regimen once the patient has initiated it in order to prevent the patient from acquiring resistant organisms. This might be true for confirmed bacterial infections. In fact, giving the right antibiotic at an adequate dose, along with good compliance with the daily regimen by the patient, i.e. taking the correct dose at the appropriate intervals, may be more important for treatment success than taking an antibiotic for a long period of time [10]. Prescribing an adequate dose of an antibiotic improves its clinical efficacy for a serious bacterial infection, with longer courses of antibiotics being associated with the greatest risk of antimicrobial resistance at both an individual and community level [11]. Some studies on carriage of antibiotic-resistant pneumococcal strains have shown that a high dose of antibiotic for a shorter duration results in less bacterial resistance than a lower dose for a longer duration [12,13]. Ideally, antibiotics should be dosed according to their pharmacokinetic and pharmacodynamic qualities to achieve the best clinical outcomes for the patient, as well as limiting the spread of antimicrobial resistance [14].

GPs very often visit patients with suspected viral infections, mainly of the upper and lower airways, for which antibiotic treatment makes no difference in terms of clinical outcomes.

1
2
3 Conversely, continuing an antibiotic regimen when this is not indicated might prompt the
4 acquisition of resistant bugs and cause adverse events [15]. Therefore, stopping the antibiotic
5 regimen once a previously suspected viral infection has been confirmed should be common
6 practice. Since 2011 the Spanish Society of Family Medicine has been recommending GPs to
7 ask their patients to stop taking antibiotics when they suspect a viral infection [16]. Despite
8 this recommendation, GPs are reluctant and feel unsafe to discontinue an antibiotic once the
9 patient has already started it. In fact, there is no evidence that discontinuing antibiotic therapy
10 for these conditions is safe and studies demonstrating that this practice should be carried out
11 are lacking. In Europe, the prescription for antibiotics differs between countries, and
12 antibiotics are prescribed significantly more often in southern compared to northern Europe,
13 with Spain being one of the countries with the highest consumption [17].

20
21 Discontinuing an antibiotic regimen could be carried out in the following situations:

22
23
24
25 *Patients diagnosed with infectious diseases for which antibiotics are not necessary, i.e. an*
26 *antibiotic course for an episode of common cold or influenza.*

27
28
29 According to the World Health Organization, 80% of the RTIs in the community have a viral
30 origin. Recent systematic reviews have suggested that antibiotics make no difference to the
31 course of common cold and influenza when compared to placebo [18,19]. Despite having a
32 viral aetiology, 2.2% of 12,373 episodes of common cold were treated with antibiotics in a
33 large study including 340 GPs carried out in Spain [20]. This unnecessary antibiotic
34 prescription might be mainly explained by uncertainty in the diagnosis and by GPs'
35 perceptions regarding patient expectations for a prescription [21,22]. Data from the GRACE
36 (Genomics to combat resistance against antibiotics in community-acquired lower respiratory
37 tract infections in Europe) study clearly show that antibiotics are expected by 45% of the
38 patients with lower RTIs [23].

39
40
41
42
43
44
45
46
47 *Patients diagnosed with clinical conditions for which antibiotics might be necessary but*
48 *according to the history and clinical examination the GP considers that antibiotics are not*
49 *needed because of suspected viral origin or patients feel that the drug has not worked as*
50 *expected and feel they need a clinical reassessment, i.e. an antibiotic regimen for a suspected*
51 *rhinosinusitis.*

1
2
3 Despite the fact that antibiotics might be required for some patients with uncomplicated acute
4 pharyngitis, acute rhinosinusitis, and acute bronchitis, most of these infections are self-
5 limiting and recent systematic reviews have suggested that antibiotics only slightly modify
6 the course of most of these infections [24-27]. Even among patients with acute exacerbations
7 of chronic obstructive pulmonary disease without purulent sputum, the benefit of antibiotics
8 is controversial, both in the community and the hospital settings [28-30]. However, antibiotic
9 prescription is very high in these four infectious diseases, and over 60% of adults presenting
10 with RTIs receive an antibiotic; in addition, these antibiotics are increasingly broad-spectrum
11 antimicrobial agents [31,32]. Even in Holland, the country with the lowest antibiotic
12 consumption in Europe, 46% of the antibiotics prescribed for RTIs are not needed while in
13 only 4% of the situations in which antibiotics are not prescribed they should have been
14 recommended according to the Dutch guidelines [33].
15
16
17
18
19
20
21
22
23
24

25 *Patients who have taken some doses of an antibiotic not prescribed by a health professional*
26 *for an infectious disease for which antibiotics are not necessary, i.e. from leftovers found in*
27 *the household or an antibiotic bought without prescription at the pharmacy.*
28
29

30 The sale of antibiotics and other antimicrobial medicines without prescription remains
31 widespread, with many countries lacking standard treatment guidelines and thereby
32 increasing the potential for overuse of antimicrobial medicines by the public and medical
33 professionals [34]. This practice is common outside Northern Europe and North America.
34 The percentage of non-prescription access to antimicrobials is often underestimated, and also
35 depends on the methodology used for making estimations. In 2016, the European
36 Commission published a questionnaire-based study (Eurobarometer) which was carried out in
37 28 European countries, including 1,053 respondents in Spain. This study described that 6% of
38 users reported to have obtained antibiotics in the previous year without prescription or stated
39 that they used the leftovers from a previous course [35]. However, when more reliable
40 methods are used to know how many individuals obtain antibiotics from sources other than
41 their GPs these percentages clearly increase [36]. Self-medication with antimicrobials is also
42 widespread, occurring among the population in the same countries where over-the-counter
43 sale is available [37].
44
45
46
47
48
49
50
51
52
53
54

55 **STUDY AIMS**

56
57
58
59
60

1
2
3 The main objective of the present trial is to know if discontinuation of antibiotic therapy
4 when a GP no longer considers it necessary makes any difference in terms of the number of
5 days with severe symptoms.
6

7
8 Secondary objectives:
9

- 10 1. Assessment of the incidence of adverse effects of medication
 - 11 2. Assessment of the antibiotic consumption
 - 12 3. Assessment of the satisfaction with health care and belief of the effectiveness of
13 antibiotics by the patients included
 - 14 4. Assessment of the number of complications observed within the first 3 months
- 15
16
17
18
19
20
21

22 **METHODS AND ANALYSIS**

23 **Study design**

24
25 This will be a multicentre, open-label, randomised controlled clinical trial comparing two
26 therapy strategies for uncomplicated acute RTIs. Patients will be randomly assigned to either
27 the usual strategy of continuing antibiotic treatment (control group) or discontinuing
28 antibiotic therapy (intervention group). The randomisation will be stratified by condition of
29 interest - pharyngitis, rhinosinusitis (including common cold), acute bronchitis, acute
30 exacerbations of chronic bronchitis or mild-to-moderate chronic obstructive pulmonary
31 disease, and influenza -. General practitioners (GP) will randomise patients centrally using an
32 electronic online platform. An open-label study has been designed given the type of the two
33 strategies used and considering that an effect on their beliefs is expected. Neither patients nor
34 health professionals will be blinded.
35
36
37
38
39
40
41
42
43
44

45 **Eligibility criteria**

46
47 Any patient from 18 to 75 years of age attending the GP consultation with an uncomplicated
48 RTI who has previously taken any dose of antibiotic due to any of the following 3 clinical
49 scenarios and accepts to participate in the clinical trial will be included:
50

- 51 • Patients diagnosed with clinical conditions for which antibiotics are not necessary
- 52
53
54
55
56
57
58
59
60

- Patients diagnosed with a clinical condition for which antibiotics might be necessary but according to the history and clinical examination the GP considers that antibiotics are not needed to be taken or the patients feel that the antibiotic regimen has not worked as expected and feel they need clinical reassessment
- Patients who have taken some doses of an antibiotic (from leftovers found in the household or obtained at the pharmacy without any medical prescription) for a clinical condition for which antibiotics are not necessary

Exclusion criteria

- Subjects under 18 and over 75 years of age
- Confirmed bacterial infection
- Patients requiring hospital admission
- Severe impairment of signs (impairment of consciousness, respiratory rate > 30 respirations per minute, heart rate > 125 beats per minute, systolic blood pressure < 90 mm Hg, diastolic blood pressure < 60 mm Hg, temperature > 40°C, oxygen saturation < 92%)
- Problems to comply with treatment at home – sociopathy or psychiatric problems, drug or alcohol addiction, or within an inadequate family setting –
- Lack of tolerance to oral treatment, such as the presence of nausea and vomiting, gastrectomy, post-surgery and/or diarrhoea
- Significant comorbidity including severe renal failure, hepatic cirrhosis, severe heart failure, immunosuppression – chronic HIV infection, transplantation, neutropenic, or patients receiving immunosuppressive drugs or corticosteroids –
- Terminal disease
- Patients admitted to a long-term residence
- Difficulty to attend the programmed visits
- Refusal to participate in the study

Outcome measures

1
2
3 The primary outcome measure will be the duration of severe symptoms. Each symptom will
4 be scored using a six-point Likert scale and symptoms scoring 5 or 6 will be considered as
5 severe. We will include common symptoms such as fever, discomfort or general pain, cough,
6 difficulty in sleeping, and changes in everyday life in all patients, and specific symptoms
7 according to the condition (Table 1) [38].
8
9

10 11 Secondary endpoints

- 12 • Adverse events of medication, reported by the patients
- 13 • Antibiotic consumption. Patients will be asked about the antibiotic use in the two arms
- 14 • Satisfaction with health care by means a questionnaire with a Likert scale to patients
- 15 • Belief in the effectiveness of antibiotics by means a questionnaire with a Likert scale to
- 16 patients
- 17 • Complications related to the RTIs will be registered during the first 3 months after
- 18 randomisation. These complications will be prospectively recorded by the GPs by means
- 19 of a standardised questionnaire and will be reported in a maximum of 48 hours to the
- 20 study clinical coordinator, who will report to the Safety and Data Monitoring Committee,
- 21 whose aim is to assess the safety of the two strategies
- 22
23
24
25
26
27
28
29
30
31
32
33

34 35 **Sample size calculation**

36 Excellence guidelines for treatment of RTIs include estimates of average duration of the
37 illness (before and after seeing a doctor) of one week for acute sore throat, one and a half
38 weeks for rhinosinusitis, including common cold, and three weeks for acute cough or
39 bronchitis [39]. In a recent study on the effectiveness of delayed prescribing strategies carried
40 out in Spain the mean duration of severe symptoms in uncomplicated acute RTIs including
41 pharyngitis, rhinosinusitis, acute bronchitis, and acute exacerbation of chronic bronchitis,
42 among patients treated immediately with antibiotics was 3.6 days (standard deviation of 3.3)
43 [38]. Considering a reduction of one day in the duration of severe symptoms as a clinically
44 relevant outcome, with a bilateral approximation, a sample of 215 patients per group will be
45 able to detect this difference with an alpha error of 5% and a power of 80% (beta =0.2) and
46 considering 20% of losses.
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 In terms of the number of researchers to be included in this clinical trial, we anticipate that
4 each researcher will include 10-12 patients in two years. Thus, we are planning to invite
5 approximately 40 researchers from 20 different healthcare centres.
6
7
8
9

10 **Data collection and ascertainment of visits**

11
12 The patients will be randomised to one of the 2 treatment strategies. Baseline data will be
13 collected in the clinic by the physician or with the help of the nursing staff. To standardize
14 data collection, all of the participating healthcare professionals will be trained by the
15 coordinating centre. The patients will receive information on the study and, if they are
16 interested in participating, they will be provided with an informed consent form to read and
17 sign. A maximum length of 15-20 minutes is expected for the interview and the introduction
18 of the data. The study scheme and the visit program will be explained to the patient (Table 2).
19
20
21
22
23

24 After randomisation, information on the strategy to which they have been allocated will be
25 given to the participants, and they will be informed as to the appropriate measures to take in
26 case of worsening or no improvement of their condition. In addition, they will be given a
27 diary with a validated questionnaire of symptoms for each condition, which they should
28 complete while symptoms related to the respiratory condition are present. The degree of
29 satisfaction or concern with different aspects of the therapy as well as the use of antibiotics
30 will also be recorded in this diary.
31
32
33
34
35

36 Patients will be interviewed by telephone 2 or 3 days after their inclusion in the study. At this
37 first follow-up visit, a worsening of the clinical situation of the patient will be evaluated to
38 determine whether a change in the antibiotic treatment is necessary among patients in the
39 intervention group or the antibiotic regimen should be continued in patients in the control
40 group. We will provide the patients with the opportunity to ask questions and discuss any
41 problems with diary completion. In addition, compliance and possible secondary effects of
42 the treatment will be evaluated. The second follow-up visit will be scheduled at day 14-28
43 after inclusion and will depend on the infection involved at patient recruitment as established
44 in the NICE guideline [39]. On this visit, the clinical evolution of the signs and symptoms
45 will be evaluated and the need to change or continue the antibiotic treatment, respectively, in
46 the case of worsening in the clinical manifestations will be determined. Symptom diaries will
47 be collected on this visit. Participants who do not return their diary will be telephoned by the
48 trial team in order to provide a reminder and enquire whether they need any assistance in
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 returning their diary or given the option of giving the information required over the phone
4 (minimum data set). Possible secondary effects of the treatment will also be assessed.
5
6 Patients will be asked to revisit again if symptoms continue or they present recurrence of the
7 infection. On the last follow-up visit, scheduled at day 90, the recurrences or medical visits or
8 hospital admissions due to the basal infectious disease will be reported.
9
10

11 12 13 **Data analysis procedure**

14
15 Characteristics of the study population will be described using frequencies for categorical
16 variables and mean and standard deviation (SD) for quantitative variables. To compare the
17 two strategies studied, we will use the Fisher's exact test for categorical variables and the t
18 test for continuous variables. To compare the duration of symptoms across strategies, we will
19 use the t test for continuous variables for each symptom, and a linear regression model per
20 symptom, with symptom duration as the dependent variable, and both antibiotic consumption
21 and symptom duration prior to the visit as the independent variables. The same approach will
22 be performed to compare the severity of symptoms between the two strategies, but the
23 dependent variable for each model will be symptom severity. The Chi-square test will be
24 used to compare antibiotic consumption, satisfaction, percentage of adverse effects, and the
25 appearance of complications between the intervention and control groups. The level of
26 significance will be 5% ($\alpha=0.05$).
27
28
29
30
31
32
33
34

35
36 Criteria for withdrawals will be the presence of a serious adverse event, an unsatisfactory
37 therapeutic effect, protocol violation or withdrawal of informed consent. Patients will be able
38 to choose to interrupt the medication any time during the course of the study. However, they
39 will be followed in the same way as the other patients. Analyses will follow the intention-to-
40 treat principle in such a way that any event in any patient will be included in the group to
41 which the patient was randomised.
42
43
44
45
46
47

48 **ETHICS AND DISSEMINATION**

49
50 Each patient should give their written consent to participate in the study after being informed
51 in intelligible language for him/her on the nature, scope and possible consequences of the
52 trial. After consent is submitted, the patients will be randomised. Data confidentiality will be
53 ensured at all times, as stated in the researcher's commitment sheet, as will compliance with
54
55
56
57
58
59
60

1
2
3 the current legislation regarding the protection of personal data. This is a clinical trial based
4 on the outpatient setting, and neither patients nor researchers will receive any monetary
5 compensation. From an ethical point of view, this is to certify that the objective of the study
6 is relevant for primary care, the power of the study may be considered as reasonable, this is
7 an original study, the risks which the participants may incur justify the study being carried
8 out with a totally favourable benefit/risk quotient, and we ensure the external validity of the
9 study to the primary care reality, with the inclusion and exclusion criteria clearly described.
10
11
12
13
14 Vulnerable populations will not be participating in this study.
15

16
17 The trial has been registered with the National Institutes of Health (NIH) trial registry
18 (NCT02900820). This protocol was presented in the last Conference of the Spanish Society
19 of Family Medicine held in La Coruña in June 2016. A range of dissemination activities are
20 planned at national and international conferences. We will publish the final report in an open
21 access peer-review journal. A summary of the findings will be sent to the participating
22 practices on completion of the STOP-AB study.
23
24
25
26
27

28 **DISCUSSION**

29
30 Overprescribing of antibiotics for RTIs is considerable in the Western countries. The results
31 of this study will be applicable to patients diagnosed with clinical conditions for which
32 antibiotics are not or might not be necessary. Few GPs in Spain ask patients to discontinue
33 antibiotic therapy because of fear of possible worsening of symptoms and the appearance of
34 complications even in cases of suspected viral infections. In other situations, GPs do not wish
35 to contradict the decision made by other doctors because it might look unfair. If the results of
36 this study do not find any difference between the two groups there will be no reason to
37 withhold the discontinuation of an antibiotic regimen when a GP no longer considers it
38 necessary. We believe that there is no risk – and every advantage – in stopping a course of an
39 antibiotic immediately after a bacterial infection has been excluded or is unlikely, as Gilbert
40 recently suggested [15]. The most obvious circumstances in which it is appropriate to stop
41 antibiotics are when the antibiotics are initiated without certainty of what infection is being
42 treated, if any treatable bacterial infection is present at all, and for infections that are almost
43 always self-limiting, e.g. episodes of acute bronchitis. Patient expectation often plays a role
44 in the decision to start antibiotic treatment in these cases.
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 A post-trial implementation observational clinical study is planned to be carried out after the
4 clinical trial aimed at knowing the percentage of the use of the strategy of discontinuing
5 antibiotic treatment by GPs, and the pros and cons of its use by means of face to face or
6 telephone interviews to patients and GPs. These interviews will be audio recorded,
7 transcribed and analysed using standard qualitative techniques such as framework analysis.
8 This qualitative-based study will allow us to better identify the aspects of the physician-
9 patient interview that should be highlighted when discontinuing an ongoing antibiotic
10 regimen earlier than the agreed duration.
11
12
13
14
15

16 The impact of our trial is likely to be important since the evidence available in this field is
17 still limited and adequately designed and conducted clinical trials are warranted to clarify the
18 effectiveness of discontinuing antibiotic therapy in these cases. Similarly, the inclusion of
19 most of the uncomplicated RTIs will increase the validity of the results. These characteristics
20 make this study innovative and relevant to this field.
21
22
23
24
25

26 27 **CONTRIBUTORS**

28 CL, AM, JMC and CB were involved in the conception and design of the study. JFF, JMM,
29 JR, IG, JA, JO, CS, JB and PR have made substantial contributions to conception and design.
30 AM, JFF, JMM, JR, IG, JA, JO, CS, JB and PR will be involved in trial conduct and
31 recruitment. CL drafted the manuscript and all authors read and approved the final
32 manuscript.
33
34
35
36
37
38
39

40 41 **FUNDING**

42 This work is supported by the Catalan Society of Family Medicine, grant number FAP1601.
43 Neither the funding society nor the sponsor – Spanish Society of Family and Community
44 Medicine – has any role in the design of the study.
45
46
47
48

49 50 **ETHICAL APPROVAL**

51 The study was approved by the Primary Care Research and Ethics Committee of the *Jordi*
52 *Gol i Gurina* Foundation, Barcelona, Spain (reference number, 16/093).
53
54
55
56
57
58
59
60

References

1. Hak E, Rovers MM, Kuyvenhoven MM, Schellevis FG, Verheij TJM. Incidence of GP-diagnosed respiratory tract infections according to age, gender and high-risk co-morbidity: the Second Dutch National Survey of General Practice. *Fam Pract* 2006;23:291–4.
2. Renati S, Linder J. Necessity of office visits for acute respiratory tract infections in primary care. *Fam Pract* 2016;33:312–7.
3. Linder JA, Stafford RS. Antibiotic treatment of adults with sore throat by community primary care physicians: a national survey, 1989-1999. *JAMA* 2001;286:1181–6.
4. Grijalva CJ, Nuorti JP, Griffin MR. Antibiotic prescription rates for acute respiratory tract infections in US ambulatory settings. *JAMA* 2009;302:758–66.
5. Goossens H, Ferech M, Vander Stichele R, Elseviers M, ESAC Project Group. Outpatient antibiotic use in Europe and association with resistance: a cross-national database study. *Lancet* 2005;365:579–87.
6. Costelloe C, Metcalfe C, Lovering A, Mant D, Hay AD. Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: systematic review and meta-analysis. *BMJ* 2010;340:c2096.
7. Smith R, Coast J. The true cost of antimicrobial resistance. *BMJ* 2013;346:f1493.
8. Armbruster S, Goldkind L. A 5-year retrospective review of experience with *Clostridium difficile*-associated diarrhea. *Mil Med* 2012;177:456–9.
9. Little P, Gould C, Williamson I, Warner G, Gantley M, Kinmonth AL. Reattendance and complications in a randomised trial of prescribing strategies for sore throat: the medicalising effect of prescribing antibiotics. *BMJ* 1997;315:350–2.
10. Ball P, Baquero F, Cars O, File T, Garau J, Klugman K, et al; on behalf of the Consensus Group on Resistance and Prescribing in Respiratory Tract Infection. Antibiotic therapy of community respiratory tract infections: strategies for optimal outcomes and minimized resistance emergence. *J Antimicrob Chemother* 2002;49:31–40.
11. Bell BG, Schellevis F, Stobberingh E, Goossens H, Pringle M. A systematic review and meta-analysis of the effects of antibiotic consumption on antibiotic resistance. *BMC Infect Dis* 2014;14:13.

12. Guillemot D, Carbon C, Balkau B, Geslin P, Lecoœur H, Vauzelle-Kervroëdan F, et al. Low dosage and long treatment duration of beta-lactam: risk factors for carriage of penicillin-resistant *Streptococcus pneumoniae*. JAMA 1998;279:365–70.
13. Schrag SJ, Peña C, Fernández J, Sánchez J, Gómez V, Pérez E, et al. Effect of short-course, high-dose amoxicillin therapy on resistant pneumococcal carriage: a randomized trial. JAMA 2001;286:49–56.
14. Adembri C, Novelli A. Pharmacokinetic and pharmacodynamic parameters of antimicrobials: potential for providing dosing regimens that are less vulnerable to resistance. Clin Pharmacokinet 2009;48:517–28.
15. Gilbert GL. Knowing when to stop antibiotic therapy. Med J Aust 2015;202:121–2.
16. Grupo de Enfermedades Infecciosas. Sociedad Española de Medicina Familiar y Comunitaria. 10 mandamientos – Uso antibióticos. Decálogo para un uso más racional de los antibióticos para profesionales sanitarios, 2011. Available at: https://www.semfyec.es/es/grupos_trabajo/infecciosas/noticias_grupo/detalle/dia_mundial_resistencias_2011/ [accessed 2 January 2017]
17. Versporten A, Bolokhovets G, Ghazaryan L, Abilova V, Pyshnik G, Spasojevic T, et al; WHO/Europe-ESAC Project Group. Antibiotic use in eastern Europe: a cross-national database study in coordination with the WHO Regional Office for Europe. Lancet Infect Dis 2014;14:381–7.
18. Arroll B. Antibiotics for upper respiratory tract infections: an overview of Cochrane reviews. Respir Med 2005;99:255–61.
19. Kenealy T, Arroll B. Antibiotics for the common cold and acute purulent rhinitis. Cochrane Database Syst Rev 2013;(6):CD000247.
20. Llor C, Hernández S, Cots JM, Bjerrum L, González B, García G, et al; Grupo de Trabajo Happy Audit España. Los médicos que disponen de pruebas rápidas disminuyen significativamente la prescripción de antibióticos en el resfriado común. Rev Esp Quimioter 2013;26:12–20.
21. Whaley LE, Businger AC, Dempsey PP, Linder JA. Visit complexity, diagnostic uncertainty, and antibiotic prescribing for acute cough in primary care: a retrospective study. BMC Fam Pract 2013;14:120.

- 1
- 2
- 3 22. Cockburn J, Pit S. Prescribing behaviour in clinical practice: patients' expectations and
- 4 doctors' perceptions of patients' expectations—A questionnaire study. *Br Med J*
- 5 1997;315:520–3.
- 6
- 7
- 8 23. Coenen S, Francis N, Kelly M, Hood K, Nuttall J, Little P; on behalf of the GRACE
- 9 Project Group. Are patient views about antibiotics related to clinician perceptions,
- 10 management and outcome? A multi-country study in outpatients with acute cough. *PLoS*
- 11 *One* 2013;8:e76691.
- 12
- 13
- 14
- 15 24. Spinks A, Glasziou PP, Del Mar CB. Antibiotics for sore throat. *Cochrane Database Syst*
- 16 *Rev* 2013;(11):CD000023.
- 17
- 18
- 19 25. Young J, De Sutter A, Merenstein D, van Essen GA, Kaiser L, Varonen H, *et al.*
- 20 Antibiotics for adults with clinically diagnosed acute rhinosinusitis: a meta-analysis of
- 21 individual patient data. *Lancet* 2008;371:908–14.
- 22
- 23
- 24 26. Lemiengre MB, van Driel ML, Merenstein D, Young J, De Sutter AI. Antibiotics for
- 25 clinically diagnosed acute rhinosinusitis in adults. *Cochrane Database Syst Rev*
- 26 2012;10:CD006089.
- 27
- 28
- 29 27. Smith SM, Fahey T, Smucny J, Becker LA. Antibiotics for acute bronchitis. *Cochrane*
- 30 *Database Syst Rev* 2014;(3):CD000245.
- 31
- 32
- 33 28. Llor C, Moragas A, Hernández S, Bayona C, Miravittles M. Efficacy of antibiotic therapy
- 34 for acute exacerbations of mild to moderate chronic obstructive pulmonary disease. *Am J*
- 35 *Respir Crit Care Med* 2012;186:716–23.
- 36
- 37
- 38 29. Soler N, Esperatti M, Ewig S, Huerta A, Agustí C, Torres A. Sputum purulence-guided
- 39 antibiotic use in hospitalised patients with exacerbations of COPD. *Eur Respir J*
- 40 2012;40:1344–53.
- 41
- 42
- 43
- 44 30. Vollenweider DJ, Jarrett H, Steurer-Stey CA, Garcia-Aymerich J, Puhan MA. Antibiotics
- 45 for exacerbations of chronic obstructive pulmonary disease. *Cochrane Database Syst Rev*
- 46 2012;12:CD010257.
- 47
- 48
- 49 31. Barnett ML, Linder JA. Antibiotic prescribing to adults with sore throat in the United
- 50 States, 1997-2010. *JAMA Intern Med* 2014;174:138–40.
- 51
- 52
- 53 32. Barnett ML, Linder JA. Antibiotic prescribing for adults with acute bronchitis in the
- 54 United States, 1996-2010. *JAMA* 2014;311:2020–2.
- 55
- 56
- 57
- 58
- 59
- 60

- 1
2
3 33. Dekker AR, Verheij TJM, van der Velden AW. Inappropriate antibiotic prescription for
4 respiratory tract indications: most prominent in adult patients. *Fam Pract* 2015;32:401–7.
5
6
7 34. Morgan DJ, Okeke IN, Laxminarayan R, Perencevich EN, Weisenberg S.
8 Nonprescription antimicrobial use worldwide: a systematic review. *Lancet Infect Dis*
9 2011;11:692–701.
10
11 35. Special Eurobarometer 445, report. Antimicrobial resistance, June 2016. ISBN 978-92-
12 79-58818-1; doi:10.2875/760366. Available at: [http://ec.europa.eu/dgs/health_food-](http://ec.europa.eu/dgs/health_food-safety/amr/docs/eb445_amr_generalreport_en.pdf)
13 [safety/amr/docs/eb445_amr_generalreport_en.pdf](http://ec.europa.eu/dgs/health_food-safety/amr/docs/eb445_amr_generalreport_en.pdf) [accessed 2 January 2017]
14
15
16
17 36. Llor C, Moragas A. Pros and cons of the over-the-counter sales of antimicrobials. *WMJ*
18 2015;61:64–8.
19
20
21 37. Grigoryan L, Haaijer-Ruskamp FM, Burgerhof JGM, Burgerhof JG, Mechtler R,
22 Deschepper R, et al. Self-medication with antimicrobial drugs in Europe. *Emerg Infect*
23 *Dis* 2006;12:452–9.
24
25
26 38. de la Poza Abad M, Mas Dalmau G, Moreno Bakedano M, González González AI,
27 Canellas Criado Y, Hernández Anadón S, et al; Delayed Antibiotic Prescription (DAP)
28 Group. Prescription strategies in acute uncomplicated respiratory infections: a
29 randomized clinical trial. *JAMA Intern Med* 2016;176:21–9.
30
31
32
33 39. NICE Short Clinical Guidelines Technical Team. Respiratory tract infections—antibiotic
34 prescribing. Prescribing of antibiotics for self-limiting respiratory tract infections in
35 adults and children in primary care. National Institute for Health and Care Excellence,
36 2008.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Symptom diary

For each symptom choose the degree of affectation in the last 24 hours, according to the following Table of codes:

Code	Explanation
0	Normal or does not upset me / does not hurt
1	Affectation is insignificant or affects me slightly / light pain
2	Mild or slightly significant affectation or affects me slightly or very little / moderate pain
3	Moderately significant affectation or affects me moderately / considerable pain
4	Significant affectation or affects me considerably / intense pain
5	Highly significant affectation or affects me highly significantly / very intense pain
6	Maximum affectation or affects me a great deal / unbearable pain

Pharyngitis: 9 symptoms were recorded daily: feeling of fever, headache, general discomfort or pain, cough, sore throat, difficulty swallowing (solids or liquids), runny nose, difficulty sleeping and difficulty with activities of daily living. Each symptom will be scored following a 0-6 scale.

Rhinosinusitis: 11 symptoms were recorded daily: feeling of fever, general discomfort or pain, headache, sudden pain in the face, pain in the face when touching, runny nose, nasal mucus colour, cough, sore throat, difficulty sleeping, and difficulty with activities of daily living. Each symptom (save for nasal mucus colour) will be scored from 0 to 6. In case of presence of nasal mucus colour, the colour will be assessed: transparent (T), light yellow clear (Y), green (V) or other (O).

Acute bronchitis: 9 symptoms were recorded daily: feeling of fever, general discomfort or pain, cough, expectoration or phlegm (mucus when coughing), shortness of breath (suffocation, fatigue), pain when breathing (chest pain), chest breath sounds, difficulty sleeping and difficulty with activities of daily living. Each symptom (save for expectoration

1
2
3 or phlegm) will be scored from 0 to 6. In case of expectoration or phlegm, their colour will be
4 assessed: transparent (T), light yellow clear (Y), green (V) or other (O).
5

6 Exacerbation of chronic bronchitis or mild to moderate chronic obstructive pulmonary
7 disease: 9 symptoms were recorded daily: feeling of fever, general discomfort or pain, cough,
8 expectoration or phlegm (mucus when coughing), shortness of breath (suffocation, fatigue),
9 pain when breathing (chest pain), chest breath sounds, difficulty sleeping and difficulty with
10 daily life activities. Each symptom (except for expectoration or phlegm) will be scored from
11 0 to 6. In case of expectoration or phlegm, their colour will be assessed: transparent (T), light
12 yellow clear (Y), green (V) or other (O).
13
14
15
16
17

18 Influenza: 15 symptoms were recorded daily: feeling of fever, runny nose, sore throat,
19 headache, cough, shortness of breath (suffocation, fatigue), muscle pain, sweat or chills,
20 diarrhoea, nausea or vomits, abdominal pain, dizziness, general discomfort or pain, difficulty
21 sleeping, and difficulty with activities of daily living. Each symptom will be scored from 0 to
22 6.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 2. Visit schedule during the clinical trial

	Visit 1 Day 0	Visit 2* Day 2-3	Visit 3 Day 14-28**	Visit 4* Day 90
Medical history and physical examination	x		x	
Informed consent form	x			
Randomisation	x			
Giving out of symptom diaries	x			
Evaluation of clinical evolution		x	x	
Assessment of adverse events		x	x	
Collection of symptom diaries			x	
Reporting of recurrences, medical visits or hospital admissions due to the basal infection		x	x	x

*Phone call

**Depending on the infection (14 days for influenza and sore throat; 28 days for acute bronchitis, acute rhinosinusitis, and acute exacerbations of chronic bronchitis or mild-to-moderate chronic obstructive pulmonary disease). A further visit should be scheduled if symptoms persist on visit 3.



SPIRIT 2013 Checklist: Recommended items to address in a clinical trial protocol and related documents*

Section/item	Item No	Description	Addressed on page number
Administrative information			
Title	1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	__1__
Trial registration	2a	Trial identifier and registry name. If not yet registered, name of intended registry	__3,12__
	2b	All items from the World Health Organization Trial Registration Data Set	_____
Protocol version	3	Date and version identifier	__cover letter__
Funding	4	Sources and types of financial, material, and other support	__13__
Roles and responsibilities	5a	Names, affiliations, and roles of protocol contributors	__1,13__
	5b	Name and contact information for the trial sponsor	__14__
	5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities	__14__
	5d	Composition, roles, and responsibilities of the coordinating centre, steering committee, endpoint adjudication committee, data management team, and other individuals or groups overseeing the trial, if applicable (see Item 21a for data monitoring committee)	__N/A__

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47**Introduction**

Background and rationale	6a	Description of research question and justification for undertaking the trial, including summary of relevant studies (published and unpublished) examining benefits and harms for each intervention	_____5-7_____
	6b	Explanation for choice of comparators	_____5-7_____
Objectives	7	Specific objectives or hypotheses	_____8_____
Trial design	8	Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group), allocation ratio, and framework (eg, superiority, equivalence, noninferiority, exploratory)	_____8_____

Methods: Participants, interventions, and outcomes

Study setting	9	Description of study settings (eg, community clinic, academic hospital) and list of countries where data will be collected. Reference to where list of study sites can be obtained	_____8_____
Eligibility criteria	10	Inclusion and exclusion criteria for participants. If applicable, eligibility criteria for study centres and individuals who will perform the interventions (eg, surgeons, psychotherapists)	_____8,9_____
Interventions	11a	Interventions for each group with sufficient detail to allow replication, including how and when they will be administered	_____8-11_____
	11b	Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose change in response to harms, participant request, or improving/worsening disease)	_____11_____
	11c	Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence (eg, drug tablet return, laboratory tests)	_____10,11_____
	11d	Relevant concomitant care and interventions that are permitted or prohibited during the trial	_____N/A_____
Outcomes	12	Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg, median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen efficacy and harm outcomes is strongly recommended	_____9_____
Participant timeline	13	Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see Figure)	_____10,11,table 1_____

1				
2				
3	Sample size	14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	___9,10___
4				
5	Recruitment	15	Strategies for achieving adequate participant enrolment to reach target sample size	___10___
6				
7				

8 **Methods: Assignment of interventions (for controlled trials)**

9 Allocation:

10				
11				
12	Sequence generation	16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	___8___
13				
14				
15				
16				
17	Allocation concealment mechanism	16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	___N/A___
18				
19				
20				
21	Implementation	16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	___8___
22				
23				
24	Blinding (masking)	17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	___8___
25				
26				
27		17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	___N/A___
28				
29				
30				

31 **Methods: Data collection, management, and analysis**

32				
33	Data collection methods	18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known. Reference to where data collection forms can be found, if not in the protocol	___10,11___
34				
35				
36				
37				
38		18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols	___10,11___
39				
40				
41				
42				
43				
44				
45				
46				
47				

1				
2				
3	Data management	19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	_____11_____
4				
5				
6				
7	Statistical methods	20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	_____11_____
8				
9				
10		20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	_____N/A_____
11				
12		20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any statistical methods to handle missing data (eg, multiple imputation)	_____11_____
13				
14				
15	Methods: Monitoring			
16				
17	Data monitoring	21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent from the sponsor and competing interests; and reference to where further details about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not needed	_____10,11_____
18				
19				
20				
21				
22		21b	Description of any interim analyses and stopping guidelines, including who will have access to these interim results and make the final decision to terminate the trial	_____N/A_____
23				
24				
25	Harms	22	Plans for collecting, assessing, reporting, and managing solicited and spontaneously reported adverse events and other unintended effects of trial interventions or trial conduct	_____10,11_____
26				
27				
28	Auditing	23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent from investigators and the sponsor	_____11_____
29				
30				
31				
32	Ethics and dissemination			
33				
34	Research ethics approval	24	Plans for seeking research ethics committee/institutional review board (REC/IRB) approval	_____3,13_____
35				
36				
37	Protocol amendments	25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes, analyses) to relevant parties (eg, investigators, REC/IRBs, trial participants, trial registries, journals, regulators)	_____N/A_____
38				
39				
40				
41				
42				
43				
44				
45				
46				
47				

1				
2				
3	Consent or assent	26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	___ 8 ___
4				
5				
6		26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	___ N/A ___
7				
8	Confidentiality	27	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	___ 8 ___
9				
10				
11	Declaration of interests	28	Financial and other competing interests for principal investigators for the overall trial and each study site	___ Not applicable ___
12				
13				
14	Access to data	29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	___ 14 ___
15				
16				
17	Ancillary and post-trial care	30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	___ N/A ___
18				
19				
20	Dissemination policy	31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	___ 11,12 ___
21				
22				
23				
24				
25		31b	Authorship eligibility guidelines and any intended use of professional writers	___ N/A ___
26				
27		31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	___ N/A ___
28				
29	Appendices			
30				
31	Informed consent materials	32	Model consent form and other related documentation given to participants and authorised surrogates	___ N/A ___
32				
33				
34	Biological specimens	33	Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies, if applicable	___ N/A ___
35				
36				

37 *It is strongly recommended that this checklist be read in conjunction with the SPIRIT 2013 Explanation & Elaboration for important clarification on the items.
 38 Amendments to the protocol should be tracked and dated. The SPIRIT checklist is copyrighted by the SPIRIT Group under the Creative Commons
 39 "[Attribution-NonCommercial-NoDerivs 3.0 Unported](https://creativecommons.org/licenses/by-nc-nd/3.0/)" license.
 40

BMJ Open

The STOP-AB Trial protocol: Efficacy and safety of discontinuing patient antibiotic treatment when physicians no longer consider it necessary.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2016-015814.R1
Article Type:	Protocol
Date Submitted by the Author:	06-Mar-2017
Complete List of Authors:	Llor, Carl; Primary healthcare centre Barcelona-2B (Via Roma) Moragas, Ana; Universitat Rovira i Virgili, Primary Healthcare Centre Jaume I Bayona, Carolina; Primary Healthcare Centre La Marina Cots, Josep M.; Universitat de Barcelona, Primary Healthcare Centre La Marina Molero, José M. ; Primary Healthcare Centre San Andrés Ribas, Joana; Primary Healthcare Centre Hospital Son Llàtzer Fóthy, Julio Francisco ; Primary Healthcare Centre Coll d'en Rabassa Gutiérrez, Isabel ; Primary Healthcare Centre Rondilla I Sánchez, Coro ; Primary Care Centre Virxe da Peregrina Ortega, Jesús; Primary Healthcare Centre Rincón de Soto Arranz, Javier; Primary Healthcare Centre Arquitecte Bennàssar Botanes, Jenifer ; Primary Healthcare Centre Cornellà de Llobregat Robles, Purificación; Primary Care Centre Can Vidalet
Primary Subject Heading:	General practice / Family practice
Secondary Subject Heading:	Infectious diseases
Keywords:	PRIMARY CARE, Clinical trials < THERAPEUTICS, Respiratory infections < THORACIC MEDICINE, Public health < INFECTIOUS DISEASES

SCHOLARONE™
Manuscripts

Title

The STOP-AB trial protocol: Efficacy and safety of discontinuing patient antibiotic treatment when physicians no longer consider it necessary.

Authors

Carl Llor¹, Ana Moragas², Carolina Bayona³, Josep M. Cots⁴, José M. Molero⁵, Joana Ribas⁶, Julio Francisco Fóthy⁷, Isabel Gutiérrez⁸, Coro Sánchez⁹, Jesús Ortega¹⁰, Javier Arranz¹¹, Jenifer Botanes¹², Purificación Robles¹³

¹Primary Healthcare Centre Barcelona-2B (Via Roma), Spain

²Universitat Rovira i Virgili. Primary Healthcare Centre Jaume I, Tarragona, Spain

³Primary Healthcare Centre La Marina, Barcelona, Spain

⁴Universitat of Barcelona. Primary Healthcare Centre La Marina, Barcelona, Spain

⁵Primary Healthcare Centre San Andrés, Madrid, Spain

⁶Primary Healthcare Centre Hospital Son Llàtzer, Palma, Spain

⁷Primary Healthcare Centre Coll d'en Rabassa, Palma, Spain

⁸Primary Healthcare Centre Rondilla I, Valladolid, Spain

⁹Primary Care Centre Virxe da Peregrina, Pontevedra, Spain

¹⁰Primary Healthcare Centre Rincón de Soto, Rincón de Soto, Spain

¹¹Primary Healthcare Centre Arquitecte Bennàssar, Palma, Spain

¹²Primary Healthcare Centre Cornellà de Llobregat, Barcelona, Spain

¹³Primary Care Centre Can Vidalet, Esplugues de Llobregat, Spain

Address for correspondence

Carl Llor

Primary Healthcare Centre Barcelona-2B (Via Roma)

1
2
3 c. Manso 19, 3rd floor

4
5 08015 Barcelona, Catalonia, Spain

6
7 Email: carles.llor@urv.cat
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

ABSTRACT

Introduction: Since 2011 the Spanish Society of Family Medicine has recommended general practitioners (GPs) to ask their patients to stop taking antibiotics when they suspect a viral infection. However, this practice is seldom used because there is no evidence that discontinuing antibiotic therapy is safe. The main objective of this study is to determine whether discontinuation of antibiotic therapy when a GP no longer considers it necessary makes any difference in terms of the number of days with severe symptoms.

Methods and analysis: This is a multicentre, open-label, randomised controlled clinical trial. The study will be conducted in ten primary care centres in Spain. We will include patients from 18 to 75 years of age with uncomplicated acute respiratory tract infections – acute rhinosinusitis, acute sore throat, influenza, or acute bronchitis – who have previously taken any dose of antibiotic for less than 3 days, but physicians no longer consider it necessary. The patients will be randomly assigned to the usual strategy of continuing antibiotic treatment or to discontinuing antibiotic therapy. A sample size of 240 patients per group was calculated on the basis of a reduction of one day in the duration of severe symptoms being a clinically relevant outcome. The primary outcome will be duration of severe symptoms, i.e. symptoms scored 5 or 6 by means of validated symptom diaries. Secondary outcomes will include: antibiotics taken, adverse events, patient satisfaction, and complications within the first 3 months.

Ethics and dissemination: The study was approved by the Ethical Board of *Fundació Jordi Gol i Gurina* (reference number: 16/093). The findings of this trial will be disseminated through research conferences and peer-reviewed journals.

Trial registration number: NCT02900820.

Strengths and limitations of this study

- The open design will allow us to study the perceptions of patients in a situation similar to usual practice. Apart from its pragmatic design our study will be the first trial to assess if discontinuation of antibiotic therapy when a GP no longer considers it necessary makes any difference in terms of the number of days with severe symptoms.
- Some patients might not complete the symptom diaries. However, strategies to improve diary return rates will be implemented and non-respondents will be telephoned.
- The open nature of the study may cause a placebo effect favouring antibiotics. However, this effect will be minimised by the similar structured information all patients will receive about the self-limiting nature of respiratory tract infections and advice about the use of non-antibiotic medications.
- This trial might be underpowered for the detection of differences between the two groups in terms of adverse events and complications within the first three months, since these outcomes are considered secondary endpoints in this study.
- Another possible limitation of this study is the fact that microbiological studies will not be taken into account. In primary care the response to treatment in respiratory tract infections is mainly judged by clinical rather than microbiological criteria.

BACKGROUND

Acute respiratory tract infections (RTIs) are among the most common reasons for a healthcare encounter in Western countries, accounting for about 15% of all visits [1]. Most episodes are caused by viruses, and in otherwise healthy adults these infections are typically self-limiting and do not require a visit to a physician or a prescription medication.

Nevertheless, many patients with uncomplicated, self-limited RTIs seek care in primary care offices. A recent study found that 72% of primary care visits due to an acute respiratory tract infection did not seem to require an office visit [2], which subsequently increases healthcare costs and often leads to inappropriate antibiotic prescription [3,4]. Very importantly, inappropriate overuse of antibiotic medications can be very detrimental and may lead to antibiotic resistance [5,6], which has a tremendous impact on the economy [7], and patients often experience adverse effects such as diarrhoea, thrush, nausea, urticaria, and rash, which give rise to further office visits and time off work. Despite being uncommon, patients might also experience serious complications such as anaphylaxis and *Clostridium difficile* infection [8]. In addition, medicalisation of a self-limited condition makes it more likely for patients to visit a healthcare provider the next time they have a similar episode [9].

General practitioners (GP) have generally been told to continue an antibiotic regimen once the patient has initiated it in order to prevent the patient from acquiring resistant organisms. However, this dogma of completing an antibiotic regimen once initiated might not be associated with less antimicrobial resistance. Some studies have shown that short-course regimens can be as effective as longer courses of therapy, resulting in less emergence of antibiotic resistance, which is consistent with what we know about natural selection, the driver of antibiotic resistance [10-12]. The use of shorter therapies, defined as the taking of an antibiotic for 5 days or less, is commonly used by women with acute uncomplicated urinary tract infections. However, in other infections, such as community-acquired pneumonia, acute exacerbations of chronic obstructive pulmonary disease, acute bacterial sinusitis, cellulitis, and pyelonephritis, shorter regimens of 3 to 6 days, which have also been shown to be as effective as longer therapies, are seldom used by GPs [13].

This often-heard statement of completing an antibiotic course once initiated and the perceived need to treat beyond resolution of symptoms is usually driven by a desire to prevent worse outcomes and relapses. Giving the right antibiotic at an adequate dose, along with good compliance with the daily regimen by the patient (i.e. taking the correct dose at the

1
2
3 appropriate intervals), has been said to be the most important strategy for treatment success in
4 bacterial infections [14]. However, GPs very often see patients with suspected viral infections
5 of the upper and lower airways, for which antibiotic treatment makes little or no difference in
6 terms of clinical outcomes and, conversely, can cause some side effects and might also
7 prompt the acquisition of resistant organisms [15]. Despite this, GPs are reluctant and feel
8 that it is unsafe to discontinue an antibiotic once the patient has already started it, mainly due
9 to the ambiguity about the appropriateness of discontinuing medication felt by GPs, and in
10 part because the clinical guidelines do not encourage discontinuation of medication, as they
11 offer GPs a weak rationale for discontinuation [16]. Moreover, when it comes to acute
12 infections, nobody wants to be seen as having withheld or discontinued antibiotic treatment
13 from a patient who subsequently deteriorates, especially if the patient winds up in hospital
14 [17].

15
16 Since 2011 the Spanish Society of Family Medicine has recommended GPs to ask their
17 patients to stop taking antibiotics given by other clinicians when they suspect a viral infection
18 [18]. Despite this recommendation, this strategy is seldom used in routine clinical practice.
19 There is limited evidence about the efficacy and safety of continuing an antibiotic regimen if
20 not needed, but concomitantly, evidence as to whether discontinuing antibiotic therapy for
21 these conditions is safe is lacking, and studies demonstrating the safety of this practice should
22 be carried out.

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37 Discontinuing an antibiotic regimen could be undertaken in one of the following two
38 situations:

- 39
40
41 *1. Patients diagnosed with infectious diseases for which antibiotics – prescribed by other*
42 *physicians – are not necessary, i.e. an antibiotic course for suspected viral infections.*

43
44 According to the World Health Organization, 80% of the RTIs in the community have a viral
45 origin. Most of these infections are self-limiting, and recent systematic reviews have
46 suggested that antibiotics only slightly modify the course of most of these infections. A
47 recent systematic review suggested that antibiotics do not improve the duration of symptoms
48 in patients with common cold compared to those receiving placebo [19]. Antibiotics are
49 associated with modest benefits in sore throat; they reduce soreness and fever, the duration of
50 symptoms, and the incidence of suppurative and non-suppurative complications compared to
51 placebo mainly in patients with infection caused by group A β -haemolytic streptococcus [20].

1
2
3 This modest effect of antibiotics has also been observed in acute rhinosinusitis in which
4 antibiotics can shorten the time to cure, but only five participants per 100 cure faster at any
5 time point between one and two weeks if they receive antibiotics instead of placebo [21]. In
6 acute bronchitis, no difference has been shown between antibiotic and placebo groups in
7 terms of the percentage of patients described as achieving clinical improvement at follow-up
8 [22]. However, the participants in all these clinical trials had a significantly greater risk of
9 adverse effects with antibiotics than with placebo [23].
10
11
12
13

14 Despite the low percentage of bacterial infections, antibiotic prescription is very high in these
15 infectious diseases, with over 60% of adults presenting with acute rhinosinusitis, acute
16 bronchitis and acute sore throat receiving an antibiotic in Spain [24]. Antibiotic prescription
17 in our country has increased over the last years and Spain now constitutes one of the leading
18 countries in the world when it comes to the percentage of antibiotics prescribed [25,26]. This
19 unnecessary antibiotic prescription in RTIs has also been seen in other affluent countries; in
20 addition, these antibiotics are increasingly broad-spectrum antimicrobial agents [27,28]. Even
21 in Holland, the country with the lowest antibiotic consumption in Europe, 46% of the
22 antibiotics prescribed for RTIs are not needed while in only 4% of the situations in which
23 antibiotics are not prescribed they should have been recommended according to the Dutch
24 guidelines [29]. This unnecessary antibiotic prescription might mainly be explained by
25 uncertainty in the diagnosis and by GPs' perceptions regarding patient expectations for a
26 prescription [30,31]. Data from the GRACE (Genomics to combat resistance against
27 antibiotics in community-acquired lower respiratory tract infections in Europe) study clearly
28 show that antibiotics are expected by 45% of the patients with lower RTIs [32].
29
30
31
32
33
34
35
36
37
38
39
40

41 *2. Patients who have taken several doses of an antibiotic not prescribed by a health*
42 *professional for an infectious disease for which antibiotics are not necessary, i.e. from*
43 *leftovers found in the household or an antibiotic bought without prescription at the*
44 *pharmacy.*
45
46
47

48 The sale of antibiotics and other antimicrobial medicines without a prescription remains
49 widespread, with many countries lacking standard treatment guidelines, thereby increasing
50 the potential for overuse of antimicrobial medicines by the public and medical professionals
51 [33]. This practice is common outside Northern Europe and North America. The percentage
52 of non-prescription access to antimicrobials is often underestimated and also depends on the
53
54
55
56
57
58
59
60

1
2
3 methodology used for making estimations. In 2016, the European Commission published a
4 questionnaire-based study (Eurobarometer) which was carried out in 28 European countries,
5 including 1,053 respondents in Spain. This study described that 6% of users reported having
6 obtained antibiotics in the previous year without a prescription or stated that they had used
7 the leftovers from a previous course [34]. However, when more reliable methods are used to
8 know how many individuals obtain antibiotics from sources other than their GPs, these
9 percentages clearly increase [35]. Self-medication with antimicrobials is also widespread,
10 occurring among the population in the same countries in which over-the-counter sales are
11 available [36].
12
13
14
15
16
17
18
19

20 **STUDY AIMS**

21
22 The main objective of the present trial is to know if the discontinuation of antibiotic therapy
23 when a GP no longer considers it necessary makes any difference in terms of the number of
24 days with severe symptoms.
25

26
27 Secondary objectives:

- 28 1. Assessment of the incidence of adverse effects of medication.
- 29 2. Assessment of antibiotic consumption.
- 30 3. Assessment of the satisfaction with healthcare and belief in the effectiveness of
31 antibiotics by the patients included.
- 32 4. Assessment of the number of complications observed within the first 3 months.
33
34
35
36
37
38
39
40

41 **METHODS AND ANALYSIS**

42 **Study design**

43
44 This will be a multicentre, open-label, randomised controlled clinical trial comparing two
45 therapeutic strategies for uncomplicated acute RTIs. Patients will be randomly assigned to
46 either the usual strategy of continuing antibiotic treatment or discontinuing antibiotic therapy.
47 The randomisation will be stratified by condition of interest – acute sore throat, rhinosinusitis
48 (including common cold), acute bronchitis, and influenza -. GPs will randomise patients
49 using a centralised electronic online platform. An open-label study has been designed given
50
51
52
53
54
55
56
57
58
59
60

1
2
3 the type of the two strategies used and considering that an effect on their beliefs is expected.
4 Neither patients nor health professionals will be blinded.
5
6
7

8 **Eligibility criteria**

9
10 Any patient from 18 to 75 years of age attending the GP consultation with an uncomplicated
11 RTI who has previously taken any dose of antibiotic for less than three days due to either of
12 the 2 following clinical scenarios and accepts to participate in the clinical trial will be
13 included:
14
15

- 16 • Patients diagnosed with clinical conditions for which antibiotics prescribed by other
17 health professionals are not necessary.
- 18 • Patients who have taken several doses of an antibiotic (from leftovers found in the
19 household or obtained at the pharmacy without a medical prescription) for a clinical
20 condition for which an antibiotic is not necessary.
21
22
23
24
25

26 **Exclusion criteria**

- 27 • Subjects under 18 or over 75 years of age.
- 28 • Confirmed bacterial infection.
- 29 • Patients requiring hospital admission.
- 30 • Severe impairment of clinical signs of infection (impairment of consciousness, respiratory
31 rate > 30 respirations per minute, heart rate > 125 beats per minute, systolic blood
32 pressure < 90 mm Hg, diastolic blood pressure < 60 mm Hg, temperature > 40°C, oxygen
33 saturation < 92%).
- 34 • Problems to comply with treatment at home, such as sociopathy or psychiatric problems,
35 or drug or alcohol addiction.
- 36 • Lack of tolerance to oral treatment, such as the presence of nausea and vomiting,
37 gastroectomy, post-surgery and/or diarrhoea.
- 38 • Significant comorbidity including severe renal failure, hepatic cirrhosis, severe heart
39 failure, immunosuppression – chronic HIV infection, transplantation, neutropenia, or
40 patients receiving immunosuppressive drugs or corticosteroids.
- 41 • Terminal disease, defined as a life expectancy less than 12 months.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- Patients admitted to a long-term residence.
- Patients who state that they are unable to see their doctor at the practice office.
- Refusal to participate in the study.

Outcome measures

The primary outcome measure will be the duration of severe symptoms. Each symptom will be scored using a six-point Likert scale and symptoms scoring 5 or 6 will be considered as severe. We will include common symptoms such as fever, discomfort or general pain, cough, difficulty in sleeping, and changes in everyday life in all patients, and specific symptoms according to the condition (Table 1) [37].

Secondary endpoints

- Adverse events of the medication given for this infection reported by the patients.
- Antibiotic consumption. Patients will be asked about antibiotic use in the two study arms.
- Patient satisfaction with healthcare by means of a questionnaire with a Likert scale.
- Patient belief in the effectiveness of antibiotics by means of a questionnaire with a Likert scale.
- Complications related to the RTIs will be registered during the first 3 months after randomisation. As in the recent study of Gulliford et al, we will consider the cases of pneumonia, empyema, peritonsillar abscess, mastoiditis, otitis media, bacterial meningitis, and intracranial abscess [38]. These complications will be prospectively recorded by the GPs by means of a standardised questionnaire and will be reported within a maximum of 48 hours to the study clinical coordinator, who will report to the Safety and Data Monitoring Committee which will assess the safety of the two strategies.

Sample size calculation

Excellence guidelines for treatment of RTIs include estimates of average duration of the illness (before and after seeing a doctor) of one week for acute sore throat, one and a half weeks for rhinosinusitis, including common cold, and three weeks for acute cough or bronchitis [39]. In a recent study on the effectiveness of delayed prescribing strategies carried

1
2
3 out in Spain, the mean duration of severe symptoms in uncomplicated acute RTIs including
4 acute sore throat, rhinosinusitis, and acute bronchitis, among patients not treated with
5 antibiotics was 4.7 days (standard deviation of 3.6) [37]. Considering a reduction of one day
6 in the duration of severe symptoms as a clinically relevant outcome, with a bilateral
7 approximation, a sample of 240 patients per group will be able to detect this difference with
8 an alpha error of 5% and a power of 80% (beta =0.2), considering 15% of losses (based on
9 the percentage of patients who did not return symptom diaries in a previous study) [37]. In
10 terms of the number of investigators to be included in this clinical trial, we anticipate that
11 over a two-year period each GP will include 32 patients. Thus, we are planning to invite 15
12 GPs from 10 different healthcare centres.
13
14
15
16
17
18
19
20
21

22 **Data collection and ascertainment of visits**

23 The patients will be randomised to one of the 2 treatment strategies. Baseline data will be
24 collected in the clinic by the physician or with the help of the nursing staff. To standardize
25 data collection, all of the participating healthcare professionals will be trained by the
26 coordinating centre. Only experienced GPs (those working for more than 15 years) and those
27 who feel comfortable with the design of the study will participate in the study. This will be
28 achieved by the administration of a questionnaire with clinical questions, recommendations
29 of local guidelines and vignettes to check if they are confident and comfortable with the
30 strategy of stopping an already commenced antibiotic course. The patients will receive
31 information on the study and, if they are interested in participating, they will be provided
32 with an informed consent form to read and sign. A maximum length of 15-20 minutes is
33 expected for the interview and the introduction of the data. The study scheme and the visit
34 programme will be explained to the patient (Table 2).
35
36
37
38
39
40
41
42

43 On the basal visit, GPs will collect information about the type of diagnoses, prior time
44 elapsed with symptoms, number of days taking an antibiotic, and type of antibiotic taken.
45 Patients who made the decision to take an antibiotic by themselves (either purchased at the
46 pharmacy or taken from leftovers stored at home) and are assigned to the usual strategy of
47 continuing antibiotic treatment will be provided with a medical prescription of the same
48 antibiotic until completing the recommended therapy duration according to local guidelines,
49 even if the antibiotic is not first-line treatment.
50
51
52
53
54
55
56
57
58
59
60

1
2
3 After randomisation, information on the strategy to which they have been allocated will be
4 given to the participants, and they will be informed as to the appropriate measures to take in
5 case their condition worsens or there is no improvement. In addition, they will be given a
6 diary with a validated questionnaire of symptoms for each condition, which they should
7 complete while symptoms related to the respiratory condition are present. The degree of
8 satisfaction or concern with different aspects of the therapy as well as the use of antibiotics
9 will also be recorded in this diary.
10
11
12
13

14 Patients will be interviewed by telephone 2 or 3 days after their inclusion in the study. At this
15 first follow-up visit, a worsening of the clinical situation of the patient will be evaluated to
16 determine whether antibiotic treatment is necessary among patients in the group assigned to
17 discontinuation (first-line antibiotics will be recommended in this case) or whether the
18 antibiotic regimen should be continued or changed to the first-line drug in patients in the
19 group allocated to continuation. We will provide the patients with the opportunity to ask
20 questions and discuss any problems with diary completion. In addition, compliance and
21 possible secondary effects of the treatment will be evaluated. The second follow-up visit will
22 be scheduled at 14-28 days after inclusion and will depend on the infection involved at
23 patient recruitment as established in the NICE guidelines [39]. On this visit, the clinical
24 evolution of the signs and symptoms will be evaluated and the need to change or continue the
25 antibiotic treatment, respectively, in the case of worsening of the clinical manifestations will
26 be determined. The symptom diaries will be collected on this visit. Participants who do not
27 return their diary will be telephoned by the trial team in order to provide a reminder and
28 enquire whether they need any assistance in returning their diary. Possible secondary effects
29 of the treatment will also be assessed. Patients will be asked to revisit if symptoms continue
30 or they present recurrence of the infection. On the last follow-up visit, scheduled at day 90,
31 recurrences, medical visits or hospital admissions due to the basal infectious disease will be
32 reported.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 **Data analysis procedure**

49

50 Characteristics of the study population will be described using frequencies for categorical
51 variables and mean and standard deviation (SD) for quantitative variables. To compare the
52 two strategies studied, we will use the t test for continuous variables. To compare the
53 duration of symptoms across strategies, we will use the t test for continuous variables for
54
55
56
57
58
59
60

1
2
3 each symptom, and a linear regression model per symptom, with symptom duration as the
4 dependent variable, and both antibiotic consumption and symptom duration prior to the visit
5 as the independent variables. The same approach will be performed to compare the severity
6 of symptoms between the two strategies, but the dependent variable for each model will be
7 symptom severity. The Chi-square test will be used to compare antibiotic consumption,
8 satisfaction, percentage of adverse effects, and the appearance of complications between the
9 two groups. The level of significance will be 5% ($\alpha=0.05$).
10
11
12
13

14 Criteria for withdrawals will be evidence of protocol violation, withdrawal of informed
15 consent or a serious adverse event, defined as any untoward medical circumstance that results
16 in death, is life-threatening, requires inpatient hospitalisation or prolongs existing
17 hospitalisation, results in persistent or significant disability/incapacity, or requires
18 intervention to prevent permanent impairment or damage. Patients will be able to choose to
19 interrupt the medication at any time during the course of the study. However, they will be
20 followed in the same way as the other patients. Analyses will follow the intention-to-treat
21 principle.
22
23
24
25
26
27
28
29

30 **ETHICS AND DISSEMINATION**

31
32 Each patient should provide written consent to participate in the study after being informed in
33 intelligible language for him/her on the nature, scope and possible consequences of the trial.
34 After consent is submitted, the patients will be randomised. Data confidentiality will be
35 ensured at all times, as stated in the researcher's commitment sheet, as will compliance with
36 the current legislation regarding the protection of personal data. This is a clinical trial based
37 on the outpatient setting, and neither patients nor researchers will receive any monetary
38 compensation. From an ethical point of view, this is to certify that the objective of the study
39 is relevant for primary care, the power of the study may be considered as reasonable, this is
40 an original study, the risks which the participants may incur justify the study being carried
41 out with a totally favourable benefit/risk quotient, and we ensure the external validity of the
42 study to the primary care reality, with clearly described inclusion and exclusion criteria.
43
44
45
46
47
48
49
50 Vulnerable populations will not participate in this study.
51

52 The trial has been registered with the National Institutes of Health (NIH) trial registry
53 (NCT02900820). This protocol was presented in the last Conference of the Spanish Society
54 of Family Medicine held in La Coruña in June 2016. A range of dissemination activities are
55
56
57
58
59
60

1
2
3 planned at national and international conferences. We will publish the final report in an open
4 access peer-review journal. A summary of the findings will be sent to the participating
5 practices on completion of the STOP-AB study.
6
7
8
9

10 **DISCUSSION**

11
12 Overprescribing of antibiotics for RTIs is considerable in Western countries. The results of
13 this study will be applicable to patients diagnosed with clinical conditions for which
14 antibiotics are not or might not be necessary. Few GPs ask patients to discontinue antibiotic
15 therapy because of fear of possible worsening of symptoms and the appearance of
16 complications even in cases of suspected viral infections. Guidelines usually provide
17 dominating triggers for prescribing and weak priming for discontinuation. In other situations,
18 GPs do not wish to contradict the decision made by other doctors because it might look
19 unfair. If the results of this study do not find any difference between the two groups, there
20 will be no reason to continue an antibiotic regimen when a GP no longer considers it
21 necessary. We believe that there is no risk – and every advantage – in stopping a course of an
22 antibiotic immediately after a bacterial infection has been excluded or is unlikely, as Gilbert
23 recently suggested [15]. The most obvious circumstances in which it is appropriate to stop
24 antibiotics are when the antibiotics are initiated without certainty of what infection is being
25 treated, if any treatable bacterial infection is present at all, and for infections that are almost
26 always self-limiting, e.g. episodes of acute bronchitis. Patient expectation often plays a role
27 in the decision to start antibiotic treatment in these cases.
28
29
30
31
32
33
34
35
36
37

38
39 The impact of our trial is likely to be important since the evidence available in this field is
40 still limited, and adequately designed and conducted clinical trials are warranted to clarify the
41 efficacy and safety of discontinuing antibiotic therapy in these cases. Similarly, the inclusion
42 of most of the uncomplicated RTIs and the fact that only patients with less than three days of
43 antibiotic therapy will be invited to participate will increase the validity of the results. These
44 characteristics make this study innovative and relevant to this field.
45
46
47
48
49
50

51 **CONTRIBUTORS**

52
53 CL, AM, JMC and CB were involved in the conception and design of the study. JFF, JMM,
54 JR, IG, JA, JO, CS, JB and PR have made substantial contributions to conception and design.
55
56
57
58
59
60

1
2
3 AM, JFF, JMM, JR, IG, JA, JO, CS, JB and PR will be involved in carrying out the trial and
4 recruitment. CL drafted the manuscript and all authors read and approved the final
5 manuscript.
6
7
8
9

10 **FUNDING**

11
12 This work is supported by the Catalan Society of Family Medicine, grant number FAP1601.
13 Neither the funding society nor the sponsor – Spanish Society of Family and Community
14 Medicine – had any role in the design of the study.
15
16
17

18 **ETHICAL APPROVAL**

19
20 The study was approved by the Primary Care Research and Ethics Committee of the *Jordi*
21 *Gol i Gurina* Foundation, Barcelona, Spain (reference number, 16/093).
22
23
24
25

26 **References**

- 27
28
29
30 1. Hak E, Rovers MM, Kuyvenhoven MM, Schellevis FG, Verheij TJM. Incidence of GP-
31 diagnosed respiratory tract infections according to age, gender and high-risk co-
32 morbidity: the Second Dutch National Survey of General Practice. *Fam Pract*
33 2006;23:291–4.
34
35
36 2. Renati S, Linder J. Necessity of office visits for acute respiratory tract infections in
37 primary care. *Fam Pract* 2016;33:312–7.
38
39
40 3. Linder JA, Stafford RS. Antibiotic treatment of adults with sore throat by community
41 primary care physicians: a national survey, 1989-1999. *JAMA* 2001;286:1181–6.
42
43
44 4. Grijalva CJ, Nuorti JP, Griffin MR. Antibiotic prescription rates for acute respiratory tract
45 infections in US ambulatory settings. *JAMA* 2009;302:758–66.
46
47
48 5. Goossens H, Ferech M, Vander Stichele R, Elseviers M, ESAC Project Group. Outpatient
49 antibiotic use in Europe and association with resistance: a cross-national database study.
50 *Lancet* 2005;365:579–87.
51
52
53
54
55
56
57
58
59
60

6. Costelloe C, Metcalfe C, Lovering A, Mant D, Hay AD. Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: systematic review and meta-analysis. *BMJ* 2010;340:c2096.
7. Smith R, Coast J. The true cost of antimicrobial resistance. *BMJ* 2013;346:f1493.
8. Armbruster S, Goldkind L. A 5-year retrospective review of experience with *Clostridium difficile*-associated diarrhea. *Mil Med* 2012;177:456–9.
9. Little P, Gould C, Williamson I, Warner G, Gantley M, Kinmonth AL. Reattendance and complications in a randomised trial of prescribing strategies for sore throat: the medicalising effect of prescribing antibiotics. *BMJ* 1997;315:350–2.
10. Singh N, Rogers P, Atwood CW, Wagener MM, Yu VL. Short-course empiric antibiotic therapy for patients with pulmonary infiltrates in the intensive care unit. A proposed solution for indiscriminate antibiotic prescription. *Am J Respir Crit Care Med* 2000;162(2 Pt 1):505–11.
11. Bell BG, Schellevis F, Stobberingh E, Goossens H, Pringle M. A systematic review and meta-analysis of the effects of antibiotic consumption on antibiotic resistance. *BMC Infect Dis* 2014;14:13.
12. Schrag SJ, Peña C, Fernández J, Sánchez J, Gómez V, Pérez E, et al. Effect of short-course, high-dose amoxicillin therapy on resistant pneumococcal carriage: a randomized trial. *JAMA* 2001;286:49–56.
13. Spellberg B. The new antibiotic mantra - "shorter is better". *JAMA Intern Med* 2016;176:1254–5.
14. Ball P, Baquero F, Cars O, File T, Garau J, Klugman K, et al; on behalf of the Consensus Group on Resistance and Prescribing in Respiratory Tract Infection. Antibiotic therapy of community respiratory tract infections: strategies for optimal outcomes and minimized resistance emergence. *J Antimicrob Chemother* 2002;49:31–40.
15. Gilbert GL. Knowing when to stop antibiotic therapy. *Med J Aust* 2015;202:121–2.
16. Nixon MS, Vendelø MT. General practitioners' decisions about discontinuation of medication: an explorative study. *J Health Organ Manag.* 2016;30:565–80.
17. Kumar S, Little P, Britten N. Why do general practitioners prescribe antibiotics for sore throat? Grounded theory interview study. *BMJ.* 2003;326:138.

18. Grupo de Enfermedades Infecciosas. Sociedad Española de Medicina Familiar y Comunitaria. 10 mandamientos – Uso antibióticos. Decálogo para un uso más racional de los antibióticos para profesionales sanitarios, 2011. Available at: https://www.semfyc.es/es/grupos_trabajo/infecciosas/noticias_grupo/detalle/dia_mundial_resistencias_2011/ [accessed 4 March 2017]
19. Kenealy T, Arroll B. Antibiotics for the common cold and acute purulent rhinitis. *Cochrane Database Syst Rev* 2013;(6):CD000247.
20. Spinks A, Glasziou PP, Del Mar CB. Antibiotics for sore throat. *Cochrane Database Syst Rev* 2013;(11):CD000023.
21. Lemiengre MB, van Driel ML, Merenstein D, Young J, De Sutter AI. Antibiotics for clinically diagnosed acute rhinosinusitis in adults. *Cochrane Database Syst Rev* 2012;(10):CD006089.
22. Smith SM, Fahey T, Smucny J, Becker LA. Antibiotics for acute bronchitis. *Cochrane Database Syst Rev* 2014;(3):CD000245.
23. Arroll B. Antibiotics for upper respiratory tract infections: an overview of Cochrane reviews. *Respir Med* 2005;99:255–61.
24. Llor C, Cots JM, Hernández S, Ortega J, Arranz J, Monedero MJ, et al; Happy Audit Study Group. Effectiveness of two types of intervention on antibiotic prescribing in respiratory tract infections in Primary Care in Spain. *Happy Audit Study. Aten Primaria*. 2014;46:492–500
25. Versporten A, Bolokhovets G, Ghazaryan L, Abilova V, Pyshnik G, Spasojevic T, et al; WHO/Europe-ESAC Project Group. Antibiotic use in eastern Europe: a cross-national database study in coordination with the WHO Regional Office for Europe. *Lancet Infect Dis* 2014;14:381–7.
26. Laxminarayan R, Sridhar D, Blaser M, Wang M, Woolhouse M. Achieving global targets for antimicrobial resistance. *Science* 2016;353:874–5.
27. Barnett ML, Linder JA. Antibiotic prescribing to adults with sore throat in the United States, 1997-2010. *JAMA Intern Med* 2014;174:138–40.
28. Barnett ML, Linder JA. Antibiotic prescribing for adults with acute bronchitis in the United States, 1996-2010. *JAMA* 2014;311:2020–2.

- 1
2
3 29. Dekker AR, Verheij TJM, van der Velden AW. Inappropriate antibiotic prescription for
4 respiratory tract indications: most prominent in adult patients. *Fam Pract* 2015;32:401–7.
5
- 6
7 30. Whaley LE, Businger AC, Dempsey PP, Linder JA. Visit complexity, diagnostic
8 uncertainty, and antibiotic prescribing for acute cough in primary care: a retrospective
9 study. *BMC Fam Pract* 2013;14:120.
10
- 11
12 31. Cockburn J, Pit S. Prescribing behaviour in clinical practice: patients' expectations and
13 doctors' perceptions of patients' expectations—A questionnaire study. *Br Med J*
14 1997;315:520–3.
15
- 16
17 32. Coenen S, Francis N, Kelly M, Hood K, Nuttall J, Little P; on behalf of the GRACE
18 Project Group. Are patient views about antibiotics related to clinician perceptions,
19 management and outcome? A multi-country study in outpatients with acute cough. *PLoS*
20 *One* 2013;8:e76691.
21
- 22
23 33. Morgan DJ, Okeke IN, Laxminarayan R, Perencevich EN, Weisenberg S.
24 Nonprescription antimicrobial use worldwide: a systematic review. *Lancet Infect Dis*
25 2011;11:692–701.
26
- 27
28 34. Special Eurobarometer 445, report. Antimicrobial resistance, June 2016. ISBN 978-92-
29 79-58818-1; doi:10.2875/760366. Available at: [http://ec.europa.eu/dgs/health_food-](http://ec.europa.eu/dgs/health_food-safety/amr/docs/eb445_amr_generalreport_en.pdf)
30 [safety/amr/docs/eb445_amr_generalreport_en.pdf](http://ec.europa.eu/dgs/health_food-safety/amr/docs/eb445_amr_generalreport_en.pdf) [accessed 4 March 2017]
31
- 32
33 35. Llor C, Moragas A. Pros and cons of the over-the-counter sales of antimicrobials. *WMJ*
34 2015;61:64–8.
35
- 36
37 36. Grigoryan L, Haaiker-Ruskamp FM, Burgerhof JGM, Burgerhof JG, Mechtler R,
38 Deschepper R, et al. Self-medication with antimicrobial drugs in Europe. *Emerg Infect*
39 *Dis* 2006;12:452–9.
40
- 41
42 37. de la Poza Abad M, Mas Dalmau G, Moreno Bakedano M, González González AI,
43 Canellas Criado Y, Hernández Anadón S, et al; Delayed Antibiotic Prescription (DAP)
44 Group. Prescription strategies in acute uncomplicated respiratory infections: a
45 randomized clinical trial. *JAMA Intern Med* 2016;176:21–9.
46
- 47
48 38. Gulliford MC, Moore MV, Little P, Hay AD, Fox R, Prevost AT, et al. Safety of reduced
49 antibiotic prescribing for self-limiting respiratory tract infections in primary care: cohort
50 study using electronic health records. *BMJ* 2016;354:i3410.
51
- 52
53
54
55
56
57
58
59

1
2
3 39. NICE Short Clinical Guidelines Technical Team. Respiratory tract infections—antibiotic
4 prescribing. Prescribing of antibiotics for self-limiting respiratory tract infections in
5 adults and children in primary care. National Institute for Health and Care Excellence,
6 2008.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 1. Symptom diary

For each symptom choose the degree of affectation in the last 24 hours, according to the following Table of Codes:

Code	Explanation
0	Normal / does not hurt
1	Affectation is insignificant / mild pain
2	Mild or slightly significant affectation / moderate pain
3	Moderately significant affectation / considerable pain
4	Significant affectation / intense pain
5	Highly significant affectation / very intense pain
6	Maximum affectation / unbearable pain

Acute sore throat: 9 symptoms will be recorded daily: feeling of fever, headache, general discomfort or pain, cough, sore throat, difficulty swallowing (solids or liquids), runny nose, difficulty sleeping and difficulty in carrying out daily life activities. Each symptom will be scored following a 0-6 scale.

Rhinosinusitis: 11 symptoms will be recorded daily: feeling of fever, general discomfort or pain, headache, sudden pain in the face, pain in the face on touching, runny nose, nasal mucus colour, cough, sore throat, difficulty sleeping, and difficulty in carrying out daily life activities. Each symptom (save for nasal mucus colour) will be scored from 0 to 6. In the case of the presence of nasal mucus colour, the colour will be assessed: transparent (T), light yellow clear (Y), green (V) or other (O).

Acute bronchitis: 9 symptoms will be recorded daily: feeling of fever, general discomfort or pain, cough, expectoration or phlegm (mucus when coughing), shortness of breath (suffocation, fatigue), pain when breathing (chest pain), chest breathing sounds, difficulty sleeping and difficulty in carrying out daily life activities. Each symptom (save for expectoration or phlegm) will be scored from 0 to 6. In the case of expectoration or phlegm, the colour will be assessed: transparent (T), light yellow clear (Y), green (V) or other (O).

1
2
3 Influenza: 15 symptoms will be recorded daily: feeling of fever, runny nose, sore throat,
4 headache, cough, shortness of breath (suffocation, fatigue), muscle pain, sweat or chills,
5 diarrhoea, nausea or vomiting, abdominal pain, dizziness, general discomfort or pain,
6 difficulty sleeping, and difficulty in carrying out daily life activities. Each symptom will be
7 scored from 0 to 6.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 2. Visit schedule during the clinical trial

	Visit 1 Day 0	Visit 2* Day 2-3	Visit 3 Day 14-28**	Visit 4* Day 90
Medical history and physical examination	x		x	
Informed consent form	x			
Randomisation	x			
Giving out of symptom diaries	x			
Evaluation of clinical evolution		x	x	
Assessment of adverse events		x	x	
Collection of symptom diaries			x	
Reporting of recurrences, medical visits or hospital admissions due to the basal infection		x	x	x

*Phone call

**Depending on the infection (14 days for influenza and sore throat; 28 days for acute bronchitis, and acute rhinosinusitis). A further visit should be scheduled if symptoms persist on visit 3.



STANDARD PROTOCOL ITEMS: RECOMMENDATIONS FOR INTERVENTIONAL TRIALS

SPIRIT 2013 Checklist: Recommended items to address in a clinical trial protocol and related documents*

Section/item	Item No	Description	Addressed on page number
Administrative information			
Title	1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	__1__
Trial registration	2a	Trial identifier and registry name. If not yet registered, name of intended registry	__3,13__
	2b	All items from the World Health Organization Trial Registration Data Set	_____
Protocol version	3	Date and version identifier	__cover letter__
Funding	4	Sources and types of financial, material, and other support	__15__
Roles and responsibilities	5a	Names, affiliations, and roles of protocol contributors	__1,14,15__
	5b	Name and contact information for the trial sponsor	__15__
	5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities	__15__
	5d	Composition, roles, and responsibilities of the coordinating centre, steering committee, endpoint adjudication committee, data management team, and other individuals or groups overseeing the trial, if applicable (see Item 21a for data monitoring committee)	__N/A__

1 **Introduction**

2

3 Background and rationale 6a Description of research question and justification for undertaking the trial, including summary of relevant studies (published and unpublished) examining benefits and harms for each intervention _____ 5-8 _____

4

5

6 6b Explanation for choice of comparators _____ 5-8 _____

7

8 Objectives 7 Specific objectives or hypotheses _____ 8 _____

9

10 Trial design 8 Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group), allocation ratio, and framework (eg, superiority, equivalence, noninferiority, exploratory) _____ 8,9 _____

11

12

13

14 **Methods: Participants, interventions, and outcomes**

15

16 Study setting 9 Description of study settings (eg, community clinic, academic hospital) and list of countries where data will be collected. Reference to where list of study sites can be obtained _____ 9 _____

17

18

19 Eligibility criteria 10 Inclusion and exclusion criteria for participants. If applicable, eligibility criteria for study centres and individuals who will perform the interventions (eg, surgeons, psychotherapists) _____ 9,10 _____

20

21

22 Interventions 11a Interventions for each group with sufficient detail to allow replication, including how and when they will be administered _____ 8-12 _____

23

24

25 11b Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose change in response to harms, participant request, or improving/worsening disease) _____ 12 _____

26

27

28 11c Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence (eg, drug tablet return, laboratory tests) _____ 11,12 _____

29

30

31 11d Relevant concomitant care and interventions that are permitted or prohibited during the trial _____ N/A _____

32

33

34 Outcomes 12 Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg, median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen efficacy and harm outcomes is strongly recommended _____ 10 _____

35

36

37

38

39

40 Participant timeline 13 Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see Figure) _____ 11,12,table 1_ _____

41

42

43

44

45

46

1 Sample size 14 Estimated number of participants needed to achieve study objectives and how it was determined, including _____10,11_____

2 clinical and statistical assumptions supporting any sample size calculations

3

4 Recruitment 15 Strategies for achieving adequate participant enrolment to reach target sample size _____11_____

5

6 **Methods: Assignment of interventions (for controlled trials)**

7

8 Allocation:

9

10 Sequence 16a Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any _____8_____

11 generation factors for stratification. To reduce predictability of a random sequence, details of any planned restriction

12 (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants

13 or assign interventions

14

15

16 Allocation 16b Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, _____N/A_____

17 concealment opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned

18 mechanism

19

20 Implementation 16c Who will generate the allocation sequence, who will enrol participants, and who will assign participants to _____8_____

21 interventions

22

23

24 Blinding (masking) 17a Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome _____8,9_____

25 assessors, data analysts), and how

26

27 17b If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's _____N/A_____

28 allocated intervention during the trial

29

30

31 **Methods: Data collection, management, and analysis**

32

33 Data collection 18a Plans for assessment and collection of outcome, baseline, and other trial data, including any related _____11,12_____

34 methods processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of

35 study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known.

36 Reference to where data collection forms can be found, if not in the protocol

37

38

39 18b Plans to promote participant retention and complete follow-up, including list of any outcome data to be _____11,12_____

40 collected for participants who discontinue or deviate from intervention protocols

41

42

1	Data management	19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	_____12_____
2				
3				
4				
5	Statistical methods	20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	_____12_____
6				
7				
8		20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	_____N/A_____
9				
10		20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any statistical methods to handle missing data (eg, multiple imputation)	_____12_____
11				
12				
13				
14	Methods: Monitoring			
15				
16	Data monitoring	21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent from the sponsor and competing interests; and reference to where further details about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not needed	_____11,12_____
17				
18				
19				
20				
21				
22		21b	Description of any interim analyses and stopping guidelines, including who will have access to these interim results and make the final decision to terminate the trial	_____N/A_____
23				
24				
25	Harms	22	Plans for collecting, assessing, reporting, and managing solicited and spontaneously reported adverse events and other unintended effects of trial interventions or trial conduct	_____11,12_____
26				
27				
28	Auditing	23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent from investigators and the sponsor	_____12_____
29				
30				
31				
32	Ethics and dissemination			
33				
34	Research ethics approval	24	Plans for seeking research ethics committee/institutional review board (REC/IRB) approval	_____3,13_____
35				
36				
37	Protocol amendments	25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes, analyses) to relevant parties (eg, investigators, REC/IRBs, trial participants, trial registries, journals, regulators)	_____N/A_____
38				
39				
40				
41				
42				
43				
44				
45				
46				

1	Consent or assent	26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	___9___
2				
3				
4		26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	___N/A___
5				
6				
7	Confidentiality	27	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	___9___
8				
9				
10	Declaration of interests	28	Financial and other competing interests for principal investigators for the overall trial and each study site	___Not applicable___
11				
12				
13	Access to data	29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	___15___
14				
15				
16	Ancillary and post-trial care	30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	___N/A___
17				
18				
19				
20	Dissemination policy	31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	___13,14___
21				
22				
23				
24		31b	Authorship eligibility guidelines and any intended use of professional writers	___N/A___
25				
26		31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	___N/A___
27				
28				
29	Appendices			
30				
31	Informed consent materials	32	Model consent form and other related documentation given to participants and authorised surrogates	___N/A___
32				
33				
34	Biological specimens	33	Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies, if applicable	___N/A___
35				
36				

37 *It is strongly recommended that this checklist be read in conjunction with the SPIRIT 2013 Explanation & Elaboration for important clarification on the items.
 38 Amendments to the protocol should be tracked and dated. The SPIRIT checklist is copyrighted by the SPIRIT Group under the Creative Commons
 39 "[Attribution-NonCommercial-NoDerivs 3.0 Unported](https://creativecommons.org/licenses/by-nc-nd/3.0/)" license.
 40
 41
 42

BMJ Open

The STOP-AB Trial protocol: Efficacy and safety of discontinuing patient antibiotic treatment when physicians no longer consider it necessary.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2016-015814.R2
Article Type:	Protocol
Date Submitted by the Author:	03-Apr-2017
Complete List of Authors:	Llor, Carl; Primary healthcare centre Barcelona-2B (Via Roma) Moragas, Ana; Universitat Rovira i Virgili, Primary Healthcare Centre Jaume I Bayona, Carolina; Primary Healthcare Centre La Marina Cots, Josep M.; Universitat de Barcelona, Primary Healthcare Centre La Marina Molero, José M. ; Primary Healthcare Centre San Andrés Ribas, Joana; Primary Healthcare Centre Hospital Son Llàtzer Fóthy, Julio Francisco ; Primary Healthcare Centre Coll d'en Rabassa Gutiérrez, Isabel ; Primary Healthcare Centre Rondilla I Sánchez, Coro ; Primary Care Centre Virxe da Peregrina Ortega, Jesús; Primary Healthcare Centre Rincón de Soto Arranz, Javier; Primary Healthcare Centre Arquitecte Bennàssar Botanes, Jenifer ; Primary Healthcare Centre Cornellà de Llobregat Robles, Purificación; Primary Care Centre Can Vidalet
Primary Subject Heading:	General practice / Family practice
Secondary Subject Heading:	Infectious diseases
Keywords:	PRIMARY CARE, Clinical trials < THERAPEUTICS, Respiratory infections < THORACIC MEDICINE, Public health < INFECTIOUS DISEASES

SCHOLARONE™
Manuscripts

Title

The STOP-AB trial protocol: Efficacy and safety of discontinuing patient antibiotic treatment when physicians no longer consider it necessary.

Authors

Carl Llor¹, Ana Moragas², Carolina Bayona³, Josep M. Cots⁴, José M. Molero⁵, Joana Ribas⁶, Julio Francisco Fóthy⁷, Isabel Gutiérrez⁸, Coro Sánchez⁹, Jesús Ortega¹⁰, Javier Arranz¹¹, Jenifer Botanes¹², Purificación Robles¹³

¹Primary Healthcare Centre Barcelona-2B (Via Roma), Spain

²Universitat Rovira i Virgili. Primary Healthcare Centre Jaume I, Tarragona, Spain

³Primary Healthcare Centre La Marina, Barcelona, Spain

⁴Universitat of Barcelona. Primary Healthcare Centre La Marina, Barcelona, Spain

⁵Primary Healthcare Centre San Andrés, Madrid, Spain

⁶Primary Healthcare Centre Hospital Son Llàtzer, Palma, Spain

⁷Primary Healthcare Centre Coll d'en Rabassa, Palma, Spain

⁸Primary Healthcare Centre Rondilla I, Valladolid, Spain

⁹Primary Care Centre Virxe da Peregrina, Pontevedra, Spain

¹⁰Primary Healthcare Centre Rincón de Soto, Rincón de Soto, Spain

¹¹Primary Healthcare Centre Arquitecte Bennàssar, Palma, Spain

¹²Primary Healthcare Centre Cornellà de Llobregat, Barcelona, Spain

¹³Primary Care Centre Can Vidalet, Esplugues de Llobregat, Spain

Address for correspondence

Carl Llor

Primary Healthcare Centre Barcelona-2B (Via Roma)

1
2
3 c. Manso 19, 3rd floor

4
5 08015 Barcelona, Catalonia, Spain

6
7 Email: carles.llor@gmail.com
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

ABSTRACT

Introduction: Since 2011 the Spanish Society of Family Medicine has recommended general practitioners (GPs) to ask their patients to stop taking antibiotics when they suspect a viral infection. However, this practice is seldom used because uncertainty about diagnosis, and fear of consequences of discontinuing antibiotic therapy, as well as perceived pressure to continue prescribing antibiotics and potential conflict with patients are more of a concern for GPs than antibiotic resistance. The main objective of this study is to determine whether discontinuation of antibiotic therapy when a GP no longer considers it necessary has any impact on the number of days with severe symptoms.

Methods and analysis: This is a multicentre, open-label, randomised controlled clinical trial. The study will be conducted in ten primary care centres in Spain. We will include patients from 18 to 75 years of age with uncomplicated acute respiratory tract infections – acute rhinosinusitis, acute sore throat, influenza, or acute bronchitis – who have previously taken any dose of antibiotic for less than 3 days, which physicians no longer consider necessary. The patients will be randomly assigned to the usual strategy of continuing antibiotic treatment or to discontinuing antibiotic therapy. A sample size of 240 patients per group was calculated on the basis of a reduction of one day in the duration of severe symptoms being a clinically relevant outcome. The primary outcome will be the duration of severe symptoms, i.e. symptoms scored 5 or 6 by means of validated symptom diaries. Secondary outcomes will include: antibiotics taken, adverse events, patient satisfaction, and complications within the first 3 months.

Ethics and dissemination: The study was approved by the Ethical Board of *Fundació Jordi Gol i Gurina* (reference number: 16/093). The findings of this trial will be disseminated through research conferences and peer-reviewed journals.

Trial registration number: NCT02900820.

Strengths and limitations of this study

- The open design of this study will allow us to study the perceptions of patients in a situation similar to that of usual practice. Apart from its pragmatic design our study will be the first trial to assess if discontinuation of antibiotic therapy when a GP no longer considers it necessary has any impact on the number of days with severe symptoms.
- Some patients might not complete the symptom diaries. However, strategies to improve diary return rates will be implemented and reminder telephone calls will be made to non-respondents.
- The open nature of the study may cause a placebo effect favouring antibiotics. However, this effect will be minimised by the similar structured information all patients will receive about the self-limiting nature of respiratory tract infections and advice about the use of non-antibiotic medications.
- This trial might be underpowered for the detection of differences between the two groups in terms of adverse events and complications within the first three months, since these outcomes are considered secondary endpoints in this study.
- Another possible limitation of this study is the fact that microbiological studies will not be taken into account. In primary care the response to treatment in respiratory tract infections is mainly judged by clinical rather than microbiological criteria.

BACKGROUND

Acute respiratory tract infections (RTIs) are among the most common reasons for a healthcare encounter in Western countries, accounting for about 15% of all visits [1]. Most episodes are caused by viruses, and in otherwise healthy adults these infections are typically self-limiting and do not require a visit to a physician or a prescription medication.

Nevertheless, many patients with uncomplicated, self-limited RTIs seek care in primary care offices. A recent study found that 72% of primary care visits due to an acute respiratory tract infection did not seem to require an office visit [2], which subsequently increases healthcare costs and often leads to inappropriate antibiotic prescription [3,4]. Very importantly, inappropriate overuse of antibiotic medications can be very detrimental and may lead to antibiotic resistance [5,6], which has a tremendous impact on the economy [7], and patients often experience adverse effects such as diarrhoea, thrush, nausea, urticaria, and rash, which give rise to further office visits and time off work. Despite being uncommon, patients might also experience serious complications such as anaphylaxis and *Clostridium difficile* infection [8]. In addition, medicalisation of a self-limited condition makes it more likely for patients to visit a healthcare provider the next time they have a similar episode [9].

General practitioners (GP) have generally been told to continue an antibiotic regimen once the patient has initiated it in order to prevent the patient from acquiring resistant organisms. However, this dogma of completing an antibiotic regimen once initiated might not be associated with less antimicrobial resistance. Some studies have shown that short-course regimens can be as effective as longer courses of therapy, resulting in less emergence of antibiotic resistance, which is consistent with what we know about natural selection, the driver of antibiotic resistance [10-12]. The use of shorter therapies, defined as the taking of an antibiotic for 5 days or less, is commonly used for uncomplicated urinary tract infections. Short course regimens – from 3 to 5 days – have also shown to be as effective as longer therapies in community-acquired pneumonia, acute bacterial sinusitis and acute exacerbations of chronic obstructive pulmonary disease but are seldom used by GPs [13]. When it comes to pneumonia, despite the efforts of infectious committees and guidelines developed by different societies, the duration of antibiotic use is still a major issue for which there is a lack of adherence both in primary and secondary care worldwide [14].

This often-heard statement of completing an antibiotic course once initiated and the perceived need to treat beyond resolution of symptoms is usually driven by a desire to prevent worse

1
2
3 outcomes and relapses. Giving the right antibiotic at an adequate dose, along with good
4 compliance with the daily regimen by the patient (i.e. taking the correct dose at the
5 appropriate intervals), has been said to be the most important strategy for treatment success in
6 bacterial infections [15]. However, GPs very often see patients with suspected viral infections
7 of the upper and lower airways, for which antibiotic treatment makes little or no difference in
8 terms of clinical outcomes and, conversely, can cause some side effects and might also
9 prompt the acquisition of resistant organisms [16].
10
11
12
13

14 Since 2011 the Spanish Society of Family Medicine has recommended GPs to ask their
15 patients to stop taking antibiotics given by other clinicians when they suspect a viral infection
16 [17]. Despite this recommendation, this strategy is seldom used in routine clinical practice.
17 GPs are reluctant and feel that it is unsafe to discontinue an antibiotic once the patient has
18 already started it, mainly due to the ambiguity about the appropriateness of discontinuing
19 medication felt by GPs, and in part because the clinical guidelines do not encourage
20 discontinuation of medication, as they offer GPs a weak rationale for discontinuation [18]. In
21 addition, some studies have also shown that other issues such as uncertainty about diagnosis,
22 ease of follow-up and fear of consequences of non-prescribing, as well as perceived pressure
23 to prescribe and potential conflict with patients which might lead to consequences for the
24 future doctor–patient relationship are more of a concern for GPs continuing to prescribe
25 antibiotics than antibiotic resistance [19]. When it comes to acute infections, GPs might feel
26 uncomfortable to discontinue antibiotic therapy from a patient who subsequently deteriorates,
27 especially if the patient needs to be admitted to hospital [20]. There is limited evidence about
28 the efficacy and safety of continuing an antibiotic regimen if not needed, but concomitantly,
29 evidence as to whether discontinuing antibiotic therapy for these conditions is safe is lacking,
30 and studies demonstrating the safety of this practice should be carried out.
31
32
33
34
35
36
37
38
39
40
41
42
43
44

45 Discontinuing an antibiotic regimen could be undertaken in one of the following two
46 situations:
47

- 48 1. *Patients diagnosed with infectious diseases for which antibiotics – prescribed by other*
49 *physicians – are not necessary, i.e. an antibiotic course for suspected viral infections.*
50
51

52 According to the World Health Organization, 80% of the RTIs in the community have a viral
53 origin. Most of these infections are self-limiting, and recent systematic reviews have
54 suggested that antibiotics only slightly modify the course of most of these infections. A
55
56
57
58
59
60

1
2
3 recent systematic review suggested that antibiotics do not improve the duration of symptoms
4 in patients with common cold compared to those receiving placebo [21]. Antibiotics are
5 associated with modest benefits in sore throat; they reduce soreness and fever, the duration of
6 symptoms, and the incidence of suppurative and non-suppurative complications compared to
7 placebo mainly in patients with infection caused by group A β -haemolytic streptococcus [22].
8 This modest effect of antibiotics has also been observed in acute rhinosinusitis in which
9 antibiotics can shorten the time to cure, but only five participants per 100 cure faster at any
10 time point between one and two weeks if they receive antibiotics instead of placebo [23]. In
11 acute bronchitis, no difference has been shown between antibiotic and placebo groups in
12 terms of the percentage of patients described as achieving clinical improvement at follow-up
13 [24]. However, the participants in all these clinical trials had a significantly greater risk of
14 adverse effects with antibiotics than with placebo [25].

15
16 Despite the low percentage of bacterial infections, antibiotic prescription is very high in these
17 infectious diseases, with over 60% of adults presenting with acute rhinosinusitis, acute
18 bronchitis and acute sore throat receiving an antibiotic in Spain [26]. Antibiotic prescription
19 in our country has increased over the last years and Spain now constitutes one of the leading
20 countries in the world when it comes to the percentage of antibiotics prescribed [27,28]. This
21 unnecessary antibiotic prescription in RTIs has also been seen in other affluent countries; in
22 addition, these antibiotics are increasingly broad-spectrum antimicrobial agents [29,30]. Even
23 in Holland, the country with the lowest antibiotic consumption in Europe, 46% of the
24 antibiotics prescribed for RTIs are not needed while in only 4% of the situations in which
25 antibiotics are not prescribed they should have been recommended according to the Dutch
26 guidelines [31]. This unnecessary antibiotic prescription might mainly be explained by
27 uncertainty in the diagnosis and by GPs' perceptions regarding patient expectations for a
28 prescription [32,33]. Data from the GRACE (Genomics to combat resistance against
29 antibiotics in community-acquired lower respiratory tract infections in Europe) study clearly
30 show that antibiotics are expected by 45% of the patients with lower RTIs [34].

31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50 2. *Patients who have taken several doses of an antibiotic not prescribed by a health*
51 *professional for an infectious disease for which antibiotics are not necessary, i.e. from*
52 *leftovers found in the household or an antibiotic bought without prescription at the*
53 *pharmacy.*
54
55

1
2
3 The sale of antibiotics and other antimicrobial medicines without a prescription remains
4 widespread, with many countries lacking standard treatment guidelines, thereby increasing
5 the potential for overuse of antimicrobial medicines by the public and medical professionals
6 [35]. This practice is common outside Northern Europe and North America. The percentage
7 of non-prescription access to antimicrobials is often underestimated and also depends on the
8 methodology used for making estimations. In 2016, the European Commission published a
9 questionnaire-based study (Eurobarometer) which was carried out in 28 European countries,
10 including 1,053 respondents in Spain. This study described that 6% of users reported having
11 obtained antibiotics in the previous year without a prescription or stated that they had used
12 the leftovers from a previous course [36]. However, when more reliable methods are used to
13 know how many individuals obtain antibiotics from sources other than their GPs, these
14 percentages clearly increase [37]. Self-medication with antimicrobials is also widespread,
15 occurring among the population in the same countries in which over-the-counter sales are
16 available [38].
17
18
19
20
21
22
23
24
25
26
27

28 **STUDY AIMS**

29
30 The main objective of the present trial is to know if the discontinuation of antibiotic therapy
31 when a GP no longer considers it necessary has any impact on the number of days with
32 severe symptoms.
33

34
35 Secondary objectives:

- 36 1. Assessment of the incidence of adverse effects.
- 37 2. Assessment of antibiotic consumption.
- 38 3. Assessment of the satisfaction with healthcare and belief in the effectiveness of
39 antibiotics by the patients included.
- 40 4. Assessment of the number of complications observed within the first 3 months.
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49

50 **METHODS AND ANALYSIS**

51 **Study design**

52
53 This will be a multicentre, open-label, randomised controlled clinical trial comparing two
54 therapeutic strategies for uncomplicated acute RTIs. Patients will be randomly assigned to
55
56
57
58
59
60

1
2
3 either the usual strategy of continuing antibiotic treatment or discontinuing antibiotic therapy.
4 The randomisation will be stratified by condition of interest – acute sore throat, rhinosinusitis
5 (including common cold), acute bronchitis, and influenza -. GPs will randomise patients
6 using a centralised electronic online platform. An open-label study has been designed taking
7 into account the type of the two strategies used and considering that an effect on GPs' and
8 patients' beliefs is expected. Neither patients nor health professionals will be blinded.
9
10
11
12
13
14

15 **Eligibility criteria**

16
17 Any patient from 18 to 75 years of age attending the GP consultation with an uncomplicated
18 RTI who has previously taken any dose of antibiotic for less than three days due to either of
19 the 2 following clinical scenarios and accepts to participate in the clinical trial will be
20 included:
21
22

- 23
24 • Patients diagnosed with clinical conditions for which antibiotics prescribed by other
25 health professionals are not necessary.
26
27
- 28 • Patients who have taken several doses of an antibiotic (from leftovers found in the
29 household or obtained at the pharmacy without a medical prescription) for a clinical
30 condition for which an antibiotic is not necessary.
31
32

33 **Exclusion criteria**

- 34
35 • Subjects under 18 or over 75 years of age.
36
37
- 38 • Confirmed bacterial infection.
39
40
- 41 • Patients requiring hospital admission.
42
43
- 44 • Severe impairment of clinical signs of infection (impairment of consciousness, respiratory
45 rate > 30 respirations per minute, heart rate > 125 beats per minute, systolic blood
46 pressure < 90 mm Hg, diastolic blood pressure < 60 mm Hg, temperature > 40°C, oxygen
47 saturation < 92%).
48
49
- 50 • Problems to comply with treatment at home, such as sociopathy or psychiatric problems,
51 or drug or alcohol addiction.
52
53
- 54 • Lack of tolerance to oral treatment, such as the presence of nausea and vomiting,
55 gastrectomy, post-surgery and/or diarrhoea.
56
57
58
59
60

- Significant comorbidity including severe renal failure, hepatic cirrhosis, severe heart failure, immunosuppression – chronic HIV infection, transplantation, neutropenia, or patients receiving immunosuppressive drugs or corticosteroids.
- Terminal disease, defined as a life expectancy less than 12 months.
- Patients admitted to a long-term residence.
- Patients who state that they are unable to see their doctor at the practice office.
- Refusal to participate in the study.

Outcome measures

The primary outcome measure will be the duration of severe symptoms. Each symptom will be scored using a six-point Likert scale and symptoms scoring 5 or 6 will be considered as severe. We will include common symptoms such as feeling of fever, discomfort or general malaise, cough, difficulty in sleeping, and changes in everyday life in all patients, and specific symptoms according to the condition (Table 1) [39].

Secondary endpoints

- Adverse events in the two study groups.
- Antibiotic consumption. This information will be collected in the symptom diaries and also by the pharmacy units of the different healthcare systems.
- Patient satisfaction with healthcare by means of a questionnaire with a Likert scale.
- Patient belief in the effectiveness of antibiotics by means of a questionnaire with a Likert scale.
- Complications related to the RTIs will be registered during the first 3 months after randomisation. As in the recent study of Gulliford et al, we will consider the cases of pneumonia, empyema, peritonsillar abscess, mastoiditis, otitis media, bacterial meningitis, and intracranial abscess [40]. These complications will be prospectively recorded by the GPs by means of a standardised questionnaire and will be reported within a maximum of 48 hours to the study clinical coordinator, who will report to the Safety and Data Monitoring Committee which will assess the safety of the two strategies.

Sample size calculation

Excellence guidelines for the treatment of RTIs include estimates of average duration of the illness (before and after seeing a doctor) of one week for acute sore throat, one and a half weeks for rhinosinusitis, including common cold, and three weeks for acute cough or bronchitis [39]. In a recent study on the effectiveness of delayed prescribing strategies carried out in Spain, the mean duration of severe symptoms in uncomplicated acute RTIs including acute sore throat, rhinosinusitis, and acute bronchitis, among patients not treated with antibiotics was 4.7 days (standard deviation of 3.6) [39]. Considering a reduction of one day in the duration of severe symptoms as a clinically relevant outcome, with a bilateral approximation, a sample of 240 patients per group will be able to detect this difference with an alpha error of 5% and a power of 80% (beta =0.2), considering 15% of losses (based on the percentage of patients who did not return symptom diaries in a previous study) [39]. In terms of the number of investigators to be included in this clinical trial, we anticipate that over a two-year period each GP will include 32 patients. Thus, we are planning to invite 15 GPs from 10 different healthcare centres.

Data collection and ascertainment of visits

The patients will be randomised to one of the 2 treatment strategies. Baseline data will be collected in the clinic by the physician or with the help of the nursing staff. To standardize data collection, all of the participating healthcare professionals will be trained by the coordinating centre. Only experienced GPs (those who have worked for 15 to 25 years) and those who feel comfortable with the design of the study will participate in the study. This will be achieved by the administration of a questionnaire with clinical questions, recommendations of local guidelines and vignettes to check if they are confident and comfortable with the strategy of stopping an already commenced antibiotic course. The patients will receive information on the study and, if they are interested in participating, they will be provided with an informed consent form to read and sign. A maximum length of 15-20 minutes is expected for the interview and the introduction of the data. The study scheme and the visit programme will be explained to the patient (Table 2). GPs will fill out a screening log with all the patients who meet all the inclusion criteria and none of the exclusion criteria regardless of whether the patients consent to participate or not. This will

1
2
3 allow us to evaluate the percentage of patients who accept to participate in the trial and
4 determine the reasons why they do not wish to participate if they refuse.
5

6 On the baseline visit, GPs will collect information about the type of diagnoses, prior time
7 elapsed with symptoms, number of days taking an antibiotic, and type of antibiotic taken.
8 Patients who made the decision to take an antibiotic by themselves (either purchased at the
9 pharmacy or taken from leftovers stored at home) and are assigned to the usual strategy of
10 continuing antibiotic treatment will be provided with a medical prescription of the same
11 antibiotic until completing the recommended therapy duration according to local guidelines,
12 even if the antibiotic is not first-line treatment.
13
14
15
16
17

18 After randomisation, information on the strategy to which they have been allocated will be
19 given to the participants, and they will be informed as to the appropriate measures to take in
20 case their condition worsens or there is no improvement. In addition, they will be given a
21 diary with a validated questionnaire of symptoms for each condition, which they should
22 complete while symptoms related to the respiratory condition are present. The degree of
23 satisfaction or concern with different aspects of the therapy as well as the use of antibiotics
24 will also be recorded in this diary.
25
26
27
28
29

30 Patients will be interviewed by telephone 2 or 3 days after their inclusion in the study. At this
31 first follow-up visit, a worsening of the clinical situation of the patient will be evaluated to
32 determine whether antibiotic treatment is necessary among patients in the group assigned to
33 discontinuation (first-line antibiotics will be recommended in this case) or whether the
34 antibiotic regimen should be continued or changed to the first-line drug in patients in the
35 group allocated to continuation. We will provide the patients with the opportunity to ask
36 questions and discuss any problems with diary completion. In addition, compliance and
37 possible secondary effects of the treatment will be evaluated. The second follow-up visit will
38 be scheduled at 14-28 days after inclusion and will depend on the infection involved at
39 patient recruitment as established in the NICE guidelines [41]. On this visit, the clinical
40 evolution of the signs and symptoms will be evaluated and the need to change or continue the
41 antibiotic treatment, respectively, will be determined in the case of worsening of the clinical
42 manifestations. The symptom diaries will be collected on this visit. Participants who do not
43 return their diary will be telephoned by the trial team as a reminder and to enquire whether
44 they need any assistance in returning their diary. Possible secondary effects of the treatment
45 will also be assessed. Patients will be asked to revisit if symptoms continue or they present
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 recurrence of the infection. On the last follow-up visit, scheduled at day 90, recurrences,
4 medical visits or hospital admissions due to the baseline infectious disease will be reported.
5
6
7

8 **Data analysis procedure**

9
10 The characteristics of the study population will be described using frequencies for categorical
11 variables and mean and standard deviation (SD) for quantitative variables. A bivariate
12 analysis of the baseline data will be performed between patients returning and not returning
13 symptoms diaries for assessing if the latter population differ from the patients including in the
14 study analysis. To compare the two strategies studied, we will use the t test for continuous
15 variables. To compare the duration of symptoms across strategies, we will use the t test for
16 continuous variables for each symptom, and a linear regression model per symptom, with
17 symptom duration as the dependent variable, and both antibiotic consumption and symptom
18 duration prior to the visit as the independent variables. The same approach will be made to
19 compare the severity of symptoms between the two strategies, but the dependent variable for
20 each model will be symptom severity. The Chi-square test will be used to compare antibiotic
21 consumption, satisfaction, percentage of adverse effects, and the appearance of complications
22 between the two groups. The level of significance will be 5% ($\alpha=0.05$).
23
24
25
26
27
28
29
30
31

32 Criteria for withdrawals will be evidence of protocol violation, withdrawal of informed
33 consent or a serious adverse event, defined as any untoward medical circumstance that results
34 in death, is life-threatening, requires inpatient hospitalisation or prolongs existing
35 hospitalisation, results in persistent or significant disability/incapacity, or requires
36 intervention to prevent permanent impairment or damage. Patients may choose to interrupt
37 the medication at any time during the course of the study. However, they will be followed in
38 the same way as the other patients. Analyses will follow the intention-to-treat principle.
39
40
41
42
43
44
45

46 **ETHICS AND DISSEMINATION**

47
48 Each patient should provide written consent to participate in the study after being informed in
49 intelligible language for him/her on the nature, scope and possible consequences of the trial.
50 After consent is submitted, the patients will be randomised. Data confidentiality will be
51 ensured at all times, as stated in the researcher's commitment sheet, as will compliance with
52 the current legislation regarding the protection of personal data. This is a clinical trial based
53
54
55
56
57
58
59
60

1
2
3 on the outpatient setting, and neither patients nor researchers will receive any monetary
4 compensation. From an ethical point of view, this is to certify that the objective of the study
5 is relevant for primary care, the power of the study may be considered as reasonable, this is
6 an original study, the risks which the participants may incur justify the study being carried
7 out with a totally favourable benefit/risk quotient, and we ensure the external validity of the
8 study to the primary care reality, with clearly described inclusion and exclusion criteria.

9
10
11
12 Vulnerable populations will not participate in this study.

13
14
15 The trial has been registered with the National Institutes of Health (NIH) trial registry
16 (NCT02900820). This protocol was presented in the last Conference of the Spanish Society
17 of Family Medicine held in La Coruña in June 2016. A range of dissemination activities are
18 planned at national and international conferences. We will publish the final report in an open
19 access peer-review journal. A summary of the findings will be sent to the participating
20 practices on completion of the STOP-AB study.

21 22 23 24 25 26 27 **DISCUSSION**

28
29 Overprescribing of antibiotics for RTIs is considerable in Western countries. The results of
30 this study will be applicable to patients diagnosed with clinical conditions for which
31 antibiotics are not or might not be necessary. Few GPs ask patients to discontinue antibiotic
32 therapy because of fear of possible worsening of symptoms and the appearance of
33 complications even in cases of suspected viral infections. Guidelines usually provide
34 dominating triggers for prescribing and weak priming for discontinuation. In other situations,
35 GPs do not wish to contradict the decision made by other doctors because it might look
36 unfair. If the results of this study do not find any difference between the two groups, there
37 will be no reason to continue an antibiotic regimen when a GP no longer considers it
38 necessary. We believe that there is no risk – and every advantage – in stopping a course of an
39 antibiotic immediately after a bacterial infection has been excluded or is unlikely, as Gilbert
40 recently suggested [16]. The most obvious circumstances in which it is appropriate to stop
41 antibiotics are when the antibiotics are initiated without certainty of what infection is being
42 treated, if any treatable bacterial infection is present at all, and for infections that are almost
43 always self-limiting, e.g. episodes of acute bronchitis. Patient expectation often plays a role
44 in the decision to start antibiotic treatment in these cases.

1
2
3 The impact of our trial is likely to be important since the evidence available in this field is
4 still limited, and adequately designed and conducted clinical trials are warranted to clarify the
5 efficacy and safety of discontinuing antibiotic therapy in these cases. Similarly, the inclusion
6 of most of the uncomplicated RTIs and the fact that only patients with less than three days of
7 antibiotic therapy will be invited to participate will increase the validity of the results. These
8 characteristics make this study innovative and relevant to this field.
9
10
11
12
13
14

15 **CONTRIBUTORS**

16
17 CL, AM, JMC and CB were involved in the conception and design of the study. JFF, JMM,
18 JR, IG, JA, JO, CS, JB and PR have made substantial contributions to conception and design.
19 AM, JFF, JMM, JR, IG, JA, JO, CS, JB and PR will be involved in carrying out the trial and
20 recruitment. CL drafted the manuscript and all authors read and approved the final
21 manuscript.
22
23
24
25
26
27

28 **FUNDING**

29
30 This work is supported by the Catalan Society of Family Medicine, grant number FAP1601.
31 Neither the funding society nor the sponsor – Spanish Society of Family and Community
32 Medicine – had any role in the design of the study.
33
34
35
36
37

38 **COMPETING INTERESTS**

39
40 CL reports receiving research grants from the European Commission (Sixth and Seventh
41 Programme Frameworks), Catalan Society of Family Medicine, and Instituto de Salud Carlos
42 III (Spanish Ministry of Health). The other authors declare no competing interests.
43
44
45
46

47 **ETHICAL APPROVAL**

48
49 The study was approved by the Primary Care Research and Ethics Committee of the *Jordi*
50 *Gol i Gurina* Foundation, Barcelona, Spain (reference number, 16/093).
51
52
53
54

55 **DATA SHARE STATEMENT**

All data created during this research will be openly available from the Spanish Society of Family Medicine.

REFERENCES

1. Hak E, Rovers MM, Kuyvenhoven MM, Schellevis FG, Verheij TJM. Incidence of GP-diagnosed respiratory tract infections according to age, gender and high-risk comorbidity: the Second Dutch National Survey of General Practice. *Fam Pract* 2006;23:291–4.
2. Renati S, Linder J. Necessity of office visits for acute respiratory tract infections in primary care. *Fam Pract* 2016;33:312–7.
3. Linder JA, Stafford RS. Antibiotic treatment of adults with sore throat by community primary care physicians: a national survey, 1989-1999. *JAMA* 2001;286:1181–6.
4. Grijalva CJ, Nuorti JP, Griffin MR. Antibiotic prescription rates for acute respiratory tract infections in US ambulatory settings. *JAMA* 2009;302:758–66.
5. Goossens H, Ferech M, Vander Stichele R, Elseviers M, ESAC Project Group. Outpatient antibiotic use in Europe and association with resistance: a cross-national database study. *Lancet* 2005;365:579–87.
6. Costelloe C, Metcalfe C, Lovering A, Mant D, Hay AD. Effect of antibiotic prescribing in primary care on antimicrobial resistance in individual patients: systematic review and meta-analysis. *BMJ* 2010;340:c2096.
7. Smith R, Coast J. The true cost of antimicrobial resistance. *BMJ* 2013;346:f1493.
8. Armbruster S, Goldkind L. A 5-year retrospective review of experience with *Clostridium difficile*-associated diarrhea. *Mil Med* 2012;177:456–9.
9. Little P, Gould C, Williamson I, Warner G, Gantley M, Kinmonth AL. Reattendance and complications in a randomised trial of prescribing strategies for sore throat: the medicalising effect of prescribing antibiotics. *BMJ* 1997;315:350–2.
10. Singh N, Rogers P, Atwood CW, Wagener MM, Yu VL. Short-course empiric antibiotic therapy for patients with pulmonary infiltrates in the intensive care unit. A proposed solution for indiscriminate antibiotic prescription. *Am J Respir Crit Care Med* 2000;162(2 Pt 1):505–11.

11. Bell BG, Schellevis F, Stobberingh E, Goossens H, Pringle M. A systematic review and meta-analysis of the effects of antibiotic consumption on antibiotic resistance. *BMC Infect Dis* 2014;14:13.
12. Schrag SJ, Peña C, Fernández J, Sánchez J, Gómez V, Pérez E, et al. Effect of short-course, high-dose amoxicillin therapy on resistant pneumococcal carriage: a randomized trial. *JAMA* 2001;286:49–56.
13. Spellberg B. The new antibiotic mantra - "shorter is better". *JAMA Intern Med* 2016;176:1254–5.
14. Viasus D, Vecino-Moreno M, De La Hoz JM, Carratalà J. Antibiotic stewardship in community-acquired pneumonia. *Expert Rev Anti Infect Ther* 2017;15:351–9.
15. Ball P, Baquero F, Cars O, File T, Garau J, Klugman K, et al; on behalf of the Consensus Group on Resistance and Prescribing in Respiratory Tract Infection. Antibiotic therapy of community respiratory tract infections: strategies for optimal outcomes and minimized resistance emergence. *J Antimicrob Chemother* 2002;49:31–40.
16. Gilbert GL. Knowing when to stop antibiotic therapy. *Med J Aust* 2015;202:121–2.
17. Grupo de Enfermedades Infecciosas. Sociedad Española de Medicina Familiar y Comunitaria. 10 mandamientos – Uso antibióticos. Decálogo para un uso más racional de los antibióticos para profesionales sanitarios, 2011. Available at: https://www.semfyec.es/es/grupos_trabajo/infecciosas/noticias_grupo/detalle/dia_mundial_resistencias_2011/ [accessed 2 April 2017]
18. Nixon MS, Vendelø MT. General practitioners' decisions about discontinuation of medication: an explorative study. *J Health Organ Manag* 2016;30:565–80.
19. Tonkin-Crine S, Yardley L, Little P. Antibiotic prescribing for acute respiratory tract infections in primary care: a systematic review and meta-ethnography. *J Antimicrob Chemother* 2011;66:2215–23.
20. Kumar S, Little P, Britten N. Why do general practitioners prescribe antibiotics for sore throat? Grounded theory interview study. *BMJ* 2003;326:138.
21. Kenealy T, Arroll B. Antibiotics for the common cold and acute purulent rhinitis. *Cochrane Database Syst Rev* 2013;(6):CD000247.

- 1
2
3 22. Spinks A, Glasziou PP, Del Mar CB. Antibiotics for sore throat. *Cochrane Database Syst*
4 *Rev* 2013;(11):CD000023.
5
- 6
7 23. Lemiengre MB, van Driel ML, Merenstein D, Young J, De Sutter AI. Antibiotics for
8 clinically diagnosed acute rhinosinusitis in adults. *Cochrane Database Syst Rev*
9 2012;(10):CD006089.
10
- 11
12 24. Smith SM, Fahey T, Smucny J, Becker LA. Antibiotics for acute bronchitis. *Cochrane*
13 *Database Syst Rev* 2014;(3):CD000245.
14
- 15
16 25. Arroll B. Antibiotics for upper respiratory tract infections: an overview of Cochrane
17 reviews. *Respir Med* 2005;99:255–61.
18
- 19
20 26. Llor C, Cots JM, Hernández S, Ortega J, Arranz J, Monedero MJ, et al; Happy Audit
21 Study Group. Effectiveness of two types of intervention on antibiotic prescribing in
22 respiratory tract infections in Primary Care in Spain. *Happy Audit Study. Aten Primaria*
23 2014;46:492–500.
24
- 25
26 27. Versporten A, Bolokhovets G, Ghazaryan L, Abilova V, Pyshnik G, Spasojevic T, et al;
27 WHO/Europe-ESAC Project Group. Antibiotic use in eastern Europe: a cross-national
28 database study in coordination with the WHO Regional Office for Europe. *Lancet Infect*
29 *Dis* 2014;14:381–7.
30
- 31
32 28. Laxminarayan R, Sridhar D, Blaser M, Wang M, Woolhouse M. Achieving global targets
33 for antimicrobial resistance. *Science* 2016;353:874–5.
34
- 35
36 29. Barnett ML, Linder JA. Antibiotic prescribing to adults with sore throat in the United
37 States, 1997-2010. *JAMA Intern Med* 2014;174:138–40.
38
- 39
40 30. Barnett ML, Linder JA. Antibiotic prescribing for adults with acute bronchitis in the
41 United States, 1996-2010. *JAMA* 2014;311:2020–2.
42
- 43
44 31. Dekker AR, Verheij TJM, van der Velden AW. Inappropriate antibiotic prescription for
45 respiratory tract indications: most prominent in adult patients. *Fam Pract* 2015;32:401–7.
46
- 47
48 32. Whaley LE, Businger AC, Dempsey PP, Linder JA. Visit complexity, diagnostic
49 uncertainty, and antibiotic prescribing for acute cough in primary care: a retrospective
50 study. *BMC Fam Pract* 2013;14:120.
51
52
53
54
55
56
57
58
59
60

- 1
2
3 33. Cockburn J, Pit S. Prescribing behaviour in clinical practice: patients' expectations and
4 doctors' perceptions of patients' expectations—A questionnaire study. *Br Med J*
5 1997;315:520–3.
6
7
8 34. Coenen S, Francis N, Kelly M, Hood K, Nuttall J, Little P; on behalf of the GRACE
9 Project Group. Are patient views about antibiotics related to clinician perceptions,
10 management and outcome? A multi-country study in outpatients with acute cough. *PLoS*
11 *One* 2013;8:e76691.
12
13
14
15 35. Morgan DJ, Okeke IN, Laxminarayan R, Perencevich EN, Weisenberg S.
16 Nonprescription antimicrobial use worldwide: a systematic review. *Lancet Infect Dis*
17 2011;11:692–701.
18
19
20 36. Special Eurobarometer 445, report. Antimicrobial resistance, June 2016. ISBN 978-92-
21 79-58818-1; doi:10.2875/760366. Available at: [http://ec.europa.eu/dgs/health_food-](http://ec.europa.eu/dgs/health_food-safety/amr/docs/eb445_amr_generalreport_en.pdf)
22 [safety/amr/docs/eb445_amr_generalreport_en.pdf](http://ec.europa.eu/dgs/health_food-safety/amr/docs/eb445_amr_generalreport_en.pdf) [accessed 2 April 2017]
23
24
25
26 37. Llor C, Moragas A. Pros and cons of the over-the-counter sales of antimicrobials. *WMJ*
27 2015;61:64–8.
28
29
30 38. Grigoryan L, Haaiker-Ruskamp FM, Burgerhof JGM, Burgerhof JG, Mechtler R,
31 Deschepper R, et al. Self-medication with antimicrobial drugs in Europe. *Emerg Infect*
32 *Dis* 2006;12:452–9.
33
34
35 39. de la Poza Abad M, Mas Dalmau G, Moreno Bakedano M, González González AI,
36 Canellas Criado Y, Hernández Anadón S, et al; Delayed Antibiotic Prescription (DAP)
37 Group. Prescription strategies in acute uncomplicated respiratory infections: a
38 randomized clinical trial. *JAMA Intern Med* 2016;176:21–9.
39
40
41
42 40. Gulliford MC, Moore MV, Little P, Hay AD, Fox R, Prevost AT, et al. Safety of reduced
43 antibiotic prescribing for self-limiting respiratory tract infections in primary care: cohort
44 study using electronic health records. *BMJ* 2016;354:i3410.
45
46
47 41. NICE Short Clinical Guidelines Technical Team. Respiratory tract infections—antibiotic
48 prescribing. Prescribing of antibiotics for self-limiting respiratory tract infections in
49 adults and children in primary care. National Institute for Health and Care Excellence,
50 2008.
51
52
53
54
55
56
57
58
59
60

Table 1. Symptom diary

For each symptom choose the degree of affectation in the last 24 hours, according to the following Table of Codes:

Code	Explanation
0	Normal / does not hurt
1	Affectation is insignificant / mild pain
2	Mild or slightly significant affectation / moderate pain
3	Moderately significant affectation / considerable pain
4	Significant affectation / intense pain
5	Highly significant affectation / very intense pain
6	Maximum affectation / unbearable pain

Acute sore throat. The following symptoms will be recorded daily: feeling of fever, headache, general discomfort or malaise, cough, sore throat, difficulty swallowing (solids or liquids), runny nose, ear pain, diarrhoea, nausea, vomiting, difficulty sleeping and difficulty in carrying out daily life activities. Each symptom will be scored following a 0-6 scale.

Rhinosinusitis. The following symptoms will be recorded daily: feeling of fever, general discomfort or malaise, headache, sudden pain in the face, pain in the face on touching, runny nose, nasal mucus colour, cough, sore throat, expectoration or phlegm (mucus when coughing), diarrhoea, nausea, vomiting, difficulty sleeping, and difficulty in carrying out daily life activities. Each symptom (save for nasal mucus colour) will be scored from 0 to 6. In the case of the presence of nasal mucus colour, the colour will be assessed: transparent (T), light yellow clear (Y), green (V) or other (O).

Acute bronchitis. The following symptoms will be recorded daily: feeling of fever, general discomfort or malaise, cough, expectoration or phlegm (mucus when coughing), shortness of breath (suffocation, fatigue), pain when breathing (chest pain), chest breathing sounds, diarrhoea, nausea, vomiting, difficulty sleeping and difficulty in carrying out daily life activities. Each symptom (save for expectoration or phlegm) will be scored from 0 to 6. In the

1
2
3 case of expectoration or phlegm, the colour will be assessed: transparent (T), light yellow
4 clear (Y), green (V) or other (O).
5

6 Influenza. The following symptoms will be recorded daily: feeling of fever, runny nose,
7 headache, general discomfort or malaise, cough, muscle pain, sore throat, sweat or chills,
8 shortness of breath, weakness or fatigue, dizziness, diarrhoea, nausea or vomiting, difficulty
9 sleeping, and difficulty in carrying out daily life activities. Each symptom will be scored from
10 0 to 6.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table 2. Visit schedule during the clinical trial

	Visit 1 Day 0	Visit 2* Day 2-3	Visit 3 Day 14-28**	Visit 4* Day 90
Medical history and physical examination	x		x	
Informed consent form	x			
Randomisation	x			
Giving out of symptom diaries	x			
Evaluation of clinical evolution		x	x	
Assessment of adverse events		x	x	
Collection of symptom diaries			x	
Reporting of recurrences, medical visits or hospital admissions due to the baseline infection		x	x	x

*Phone call

**Depending on the infection (14 days for influenza and sore throat; 28 days for acute bronchitis, and acute rhinosinusitis). A further visit should be scheduled if symptoms persist on visit 3.



STANDARD PROTOCOL ITEMS: RECOMMENDATIONS FOR INTERVENTIONAL TRIALS

SPIRIT 2013 Checklist: Recommended items to address in a clinical trial protocol and related documents*

Section/item	Item No	Description	Addressed on page number
Administrative information			
Title	1	Descriptive title identifying the study design, population, interventions, and, if applicable, trial acronym	__1__
Trial registration	2a	Trial identifier and registry name. If not yet registered, name of intended registry	__3,13__
	2b	All items from the World Health Organization Trial Registration Data Set	_____
Protocol version	3	Date and version identifier	__cover letter__
Funding	4	Sources and types of financial, material, and other support	__15__
Roles and responsibilities	5a	Names, affiliations, and roles of protocol contributors	__1,14,15__
	5b	Name and contact information for the trial sponsor	__15__
	5c	Role of study sponsor and funders, if any, in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication, including whether they will have ultimate authority over any of these activities	__15__
	5d	Composition, roles, and responsibilities of the coordinating centre, steering committee, endpoint adjudication committee, data management team, and other individuals or groups overseeing the trial, if applicable (see Item 21a for data monitoring committee)	__N/A__

1 **Introduction**

2

3 Background and rationale 6a Description of research question and justification for undertaking the trial, including summary of relevant studies (published and unpublished) examining benefits and harms for each intervention _____ 5-8 _____

4

5

6 6b Explanation for choice of comparators _____ 5-8 _____

7

8 Objectives 7 Specific objectives or hypotheses _____ 8 _____

9

10 Trial design 8 Description of trial design including type of trial (eg, parallel group, crossover, factorial, single group), allocation ratio, and framework (eg, superiority, equivalence, noninferiority, exploratory) _____ 8,9 _____

11

12

13

14 **Methods: Participants, interventions, and outcomes**

15

16 Study setting 9 Description of study settings (eg, community clinic, academic hospital) and list of countries where data will be collected. Reference to where list of study sites can be obtained _____ 9 _____

17

18

19 Eligibility criteria 10 Inclusion and exclusion criteria for participants. If applicable, eligibility criteria for study centres and individuals who will perform the interventions (eg, surgeons, psychotherapists) _____ 9,10 _____

20

21

22 Interventions 11a Interventions for each group with sufficient detail to allow replication, including how and when they will be administered _____ 8-12 _____

23

24

25 11b Criteria for discontinuing or modifying allocated interventions for a given trial participant (eg, drug dose change in response to harms, participant request, or improving/worsening disease) _____ 12 _____

26

27

28 11c Strategies to improve adherence to intervention protocols, and any procedures for monitoring adherence (eg, drug tablet return, laboratory tests) _____ 11,12 _____

29

30

31 11d Relevant concomitant care and interventions that are permitted or prohibited during the trial _____ N/A _____

32

33

34 Outcomes 12 Primary, secondary, and other outcomes, including the specific measurement variable (eg, systolic blood pressure), analysis metric (eg, change from baseline, final value, time to event), method of aggregation (eg, median, proportion), and time point for each outcome. Explanation of the clinical relevance of chosen efficacy and harm outcomes is strongly recommended _____ 10 _____

35

36

37

38

39

40 Participant timeline 13 Time schedule of enrolment, interventions (including any run-ins and washouts), assessments, and visits for participants. A schematic diagram is highly recommended (see Figure) _____ 11,12,table 1_ _____

41

42

43

44

45

46

1	Sample size	14	Estimated number of participants needed to achieve study objectives and how it was determined, including clinical and statistical assumptions supporting any sample size calculations	___10,11___
2				
3				
4	Recruitment	15	Strategies for achieving adequate participant enrolment to reach target sample size	___11___
5				
6				
7	Methods: Assignment of interventions (for controlled trials)			
8	Allocation:			
9				
10	Sequence	16a	Method of generating the allocation sequence (eg, computer-generated random numbers), and list of any factors for stratification. To reduce predictability of a random sequence, details of any planned restriction (eg, blocking) should be provided in a separate document that is unavailable to those who enrol participants or assign interventions	___8___
11	generation			
12				
13				
14				
15				
16	Allocation	16b	Mechanism of implementing the allocation sequence (eg, central telephone; sequentially numbered, opaque, sealed envelopes), describing any steps to conceal the sequence until interventions are assigned	___N/A___
17	concealment			
18	mechanism			
19				
20	Implementation	16c	Who will generate the allocation sequence, who will enrol participants, and who will assign participants to interventions	___8___
21				
22				
23				
24	Blinding (masking)	17a	Who will be blinded after assignment to interventions (eg, trial participants, care providers, outcome assessors, data analysts), and how	___8,9___
25				
26				
27		17b	If blinded, circumstances under which unblinding is permissible, and procedure for revealing a participant's allocated intervention during the trial	___N/A___
28				
29				
30				
31	Methods: Data collection, management, and analysis			
32				
33	Data collection	18a	Plans for assessment and collection of outcome, baseline, and other trial data, including any related processes to promote data quality (eg, duplicate measurements, training of assessors) and a description of study instruments (eg, questionnaires, laboratory tests) along with their reliability and validity, if known. Reference to where data collection forms can be found, if not in the protocol	___11,12___
34	methods			
35				
36				
37				
38				
39		18b	Plans to promote participant retention and complete follow-up, including list of any outcome data to be collected for participants who discontinue or deviate from intervention protocols	___11,12___
40				
41				
42				
43				
44				
45				
46				

1	Data management	19	Plans for data entry, coding, security, and storage, including any related processes to promote data quality (eg, double data entry; range checks for data values). Reference to where details of data management procedures can be found, if not in the protocol	_____12_____
2				
3				
4				
5	Statistical methods	20a	Statistical methods for analysing primary and secondary outcomes. Reference to where other details of the statistical analysis plan can be found, if not in the protocol	_____12_____
6				
7				
8		20b	Methods for any additional analyses (eg, subgroup and adjusted analyses)	_____N/A_____
9				
10		20c	Definition of analysis population relating to protocol non-adherence (eg, as randomised analysis), and any statistical methods to handle missing data (eg, multiple imputation)	_____12_____
11				
12				
13				
14	Methods: Monitoring			
15				
16	Data monitoring	21a	Composition of data monitoring committee (DMC); summary of its role and reporting structure; statement of whether it is independent from the sponsor and competing interests; and reference to where further details about its charter can be found, if not in the protocol. Alternatively, an explanation of why a DMC is not needed	_____11,12_____
17				
18				
19				
20				
21				
22		21b	Description of any interim analyses and stopping guidelines, including who will have access to these interim results and make the final decision to terminate the trial	_____N/A_____
23				
24				
25	Harms	22	Plans for collecting, assessing, reporting, and managing solicited and spontaneously reported adverse events and other unintended effects of trial interventions or trial conduct	_____11,12_____
26				
27				
28	Auditing	23	Frequency and procedures for auditing trial conduct, if any, and whether the process will be independent from investigators and the sponsor	_____12_____
29				
30				
31				
32	Ethics and dissemination			
33				
34	Research ethics approval	24	Plans for seeking research ethics committee/institutional review board (REC/IRB) approval	_____3,13_____
35				
36				
37	Protocol amendments	25	Plans for communicating important protocol modifications (eg, changes to eligibility criteria, outcomes, analyses) to relevant parties (eg, investigators, REC/IRBs, trial participants, trial registries, journals, regulators)	_____N/A_____
38				
39				
40				
41				
42				
43				
44				
45				
46				

1	Consent or assent	26a	Who will obtain informed consent or assent from potential trial participants or authorised surrogates, and how (see Item 32)	___9___
2				
3				
4		26b	Additional consent provisions for collection and use of participant data and biological specimens in ancillary studies, if applicable	___N/A___
5				
6				
7	Confidentiality	27	How personal information about potential and enrolled participants will be collected, shared, and maintained in order to protect confidentiality before, during, and after the trial	___9___
8				
9				
10	Declaration of interests	28	Financial and other competing interests for principal investigators for the overall trial and each study site	___Not applicable___
11				
12				
13	Access to data	29	Statement of who will have access to the final trial dataset, and disclosure of contractual agreements that limit such access for investigators	___15___
14				
15				
16	Ancillary and post-trial care	30	Provisions, if any, for ancillary and post-trial care, and for compensation to those who suffer harm from trial participation	___N/A___
17				
18				
19				
20	Dissemination policy	31a	Plans for investigators and sponsor to communicate trial results to participants, healthcare professionals, the public, and other relevant groups (eg, via publication, reporting in results databases, or other data sharing arrangements), including any publication restrictions	___13,14___
21				
22				
23				
24		31b	Authorship eligibility guidelines and any intended use of professional writers	___N/A___
25				
26		31c	Plans, if any, for granting public access to the full protocol, participant-level dataset, and statistical code	___N/A___
27				
28				
29	Appendices			
30				
31	Informed consent materials	32	Model consent form and other related documentation given to participants and authorised surrogates	___N/A___
32				
33				
34	Biological specimens	33	Plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies, if applicable	___N/A___
35				
36				

37 *It is strongly recommended that this checklist be read in conjunction with the SPIRIT 2013 Explanation & Elaboration for important clarification on the items.
 38 Amendments to the protocol should be tracked and dated. The SPIRIT checklist is copyrighted by the SPIRIT Group under the Creative Commons
 39 "[Attribution-NonCommercial-NoDerivs 3.0 Unported](https://creativecommons.org/licenses/by-nc-nd/3.0/)" license.
 40
 41
 42