

**Sodium tanshinone IIA sulfate adjunct therapy reduces high-sensitivity C-reactive protein level in coronary artery disease patients: a randomized controlled trial**

**Siming Li<sup>1,+</sup>, Yang Jiao<sup>1,2,+</sup>, Hanjay Wang<sup>3</sup>, Qinghua Shang<sup>1</sup>, Fang Lu<sup>4</sup>, Li Huang<sup>5</sup>, Jiangang Liu<sup>1</sup>, Hao Xu<sup>1\*</sup>, Keji Chen<sup>1</sup>**

<sup>1</sup> Cardiovascular Diseases Center, Xiyuan Hospital, China Academy of Chinese Medical Sciences, Beijing 100091, China.

<sup>2</sup> Graduate School, Beijing University of Chinese Medicine, Beijing 100029, China.

<sup>3</sup> Department of Cardiothoracic Surgery, Stanford University, Palo Alto, CA 94304, USA.

<sup>4</sup> Institute of Clinical Pharmacology, Xiyuan Hospital, China Academy of Chinese Medical Sciences, Beijing 100091, China.

<sup>5</sup> Integrative Cardiology Department, China-Japan Friendship Hospital, Beijing 100029, China.

\*Correspondence should be addressed to Hao Xu, xuhaotcm@hotmail.com.

+ These authors contributed equally to this work.

Supplement Table 1. Calculation of angina risk score

---

<b>Angina Frequency</b>	
<1 episode per week	0 points
1-6 episodes per week	2 points
1-3 episodes per day	4 points
>4 episodes per day	6 points
<b>Angina Duration, per episode</b>	
No angina	0 points
≤5 minutes	2 points
Between 5-10 minutes	4 points
≥10 minutes	6 points
<b>Angina Intensity</b>	
No angina	0 points
Relieved with rest. Does not affect daily life.	2 points
Relieved with medication. Can endure moderate exercise.	4 points
Angina frequently induced with activities of daily life (e.g. dressing, eating, walking)	6 points
<b>Nitroglycerin usage</b>	
<1 pill per week	0 points
1-4 pills per week	2 points
5-9 pills per week	4 points
≥10 pills per week	6 points

---

Total angina score equals the sum of the component scores. Minimum score = 0 points. Maximum score = 24 points.