

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email editorial.bmjopen@bmj.com

BMJ Open

The Association between the Use of Biomedical Services and the Holistic Use of Traditional East Asian Medicine among Outpatients in Korea

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-018414
Article Type:	Research
Date Submitted by the Author:	28-Jun-2017
Complete List of Authors:	Shim, Jae-Mahn; University of Seoul, Sociology Lee, Yun-Suk; University of Seoul, Sociology
Primary Subject Heading:	Complementary medicine
Secondary Subject Heading:	Sociology, Health services research, Public health
Keywords:	holism, co-utilization, East Asian medicine, Korea, utilization behavior, institutionalization

SCHOLARONE™
Manuscripts

only

1
2
3
4 **The Association between the Use of Biomedical Services and the Holistic Use of**
5 **Traditional East Asian Medicine among Outpatients in Korea**
6
7

8
9 **Authors**

10
11 1. Jae-Mahn Shim, Ph.D. (*Corresponding author)

12 Department of Sociology

13 University of Seoul

14 163 Seoulsiripdaero, Dongdaemungu

15 Seoul 130-742, Korea

16 Email: jaemahn.shim@gmail.com

17 Phone: +82-2-6490-2743

18 Fax: +82-2-6490-2734

19
20
21
22
23 2. Yun-Suk Lee, Ph.D.

24 Department of Sociology

25 University of Seoul

26 163 Seoulsiripdaero, Dongdaemungu

27 Seoul 130-742, Korea

28 Email: yslee@uos.ac.kr

29
30
31
32 **Keywords:** holism; co-utilization; East Asian medicine; Korea; utilization behavior

33
34
35 **Word count:** 3,560

Abstract

Objectives: The holistic use of a system of complementary and alternative medicine (CAM) is potentially linked to its treatment outcomes. This paper examines how the use of biomedicine is associated with the holistic use of CAM, focusing on traditional East Asian medicine (EM) that is uniquely integrated in the medical system in Korea.

Design/Settings: A representative national sample of EM outpatients in Korea.

Participants: 3,861 respondents to a national survey.

Methods: By using the 2011 Korean National Survey of EM Patients, ordered logistic regression models specify the relationship between EM outpatients' use of biomedicine and their holistic use of EM modalities.

Results: Among EM outpatients who used at least one EM modality in the past three months, people who used two (33.3%) or three (29.4%) modalities together are the two greatest in number, followed by users of four (18.1%), five (7.2%), six (2.1%), and seven (0.6%) modalities. Among these EM outpatients, 42.5% used biomedical services as well during the same period. The odds for EM users to use EM holistically are 17% greater among EM users who used biomedicine as well, compared to EM users who did not use biomedicine.

Conclusions: Health care community should recognize that CAM use likely becomes holistic as people use biomedicine concomitantly, when the practice rights over a CAM system are comprehensively and exclusively entitled to a group of CAM professionals who are independent from practitioners of biomedicine.

[Strengths and limitations of this study]

- It uses a national sample of EM users who visited EM facilities across Korea.
- It specifies several multivariate ordered logistic regression models to support the findings.
- All the measures are based upon self-reports of survey participants.

Introduction

The presence of complementary and alternative medicine (CAM) is substantial in contemporary health care systems around the world.^{1,2} Accordingly, the health care community is concerned with the ways in which CAM is integrated in the mainstream biomedicine^{3,4} and health care outcomes that CAM produces. Over 40 systematic reviews of CAM trials are currently registered in the Cochrane Library and debate the safety and efficacy of CAM. In addition, studies suggest that CAM services be provided holistically in medical trials as well as real-world practices so that CAM users can utilize all the related treatment modalities within a whole system of CAM.^{5,6} It is argued that the holistic utilization of various treatment modalities within a CAM system can maximize the treatment effects of CAM, compared to the selective and fragmented use of only some modalities of the whole system.⁷ In these studies, the holistic use of a CAM system refers to the utilization behavior of CAM users who use two or more treatment modalities that constitute the CAM system.⁸ In contrast, the selective and fragmented use refers to the utilization behavior of CAM users who use a certain modality of the CAM system and not others.

While the therapeutic effectiveness of the holistic use of CAM still needs to be adjudicated by more research, this paper attends to the possibility that CAM is institutionalized in some countries in a way that prevents the users from its holistic use. For example, the treatment modalities that together form a whole system of traditional East Asian medicine (EM), such as acupuncture, moxibustion, cupping, herbal remedies, and acupressure, are disconnected from one another in the Japanese medical system so that only herbal remedies are selectively incorporated into the practices of biomedical doctors;⁹⁻¹³ acupuncture and acupressure are

1
2
3
4 each relegated as pseudo-medicine to medical technicians, such as acupuncturists and
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

each relegated as pseudo-medicine to medical technicians, such as acupuncturists and
massage therapists who are permitted to practice only acupuncture and massage therapies.
These technicians are not allowed to practice herbal remedies or other EM modalities. Similar
observations are made that the otherwise rich practices of acupuncture are truncated and
simplified in the dominant biomedical health care systems in the U.S.¹⁴⁻¹⁶ and the U.K.¹⁷ In
the contemporary French biomedical system, a variety of CAM systems are reported to
become “balkanized” and their constituent treatment modalities are torn apart from one
another in practice.¹⁸ These studies together demonstrate that the intersection of a CAM
system with biomedicine can result in the fragmented and partial use of the CAM system.

In other countries, on the other hand, treatment modalities within a CAM system are held
together and institutionalized comprehensively into the national medical systems. For
example, several treatment modalities within EM are recognized as legitimate medicine in
China, Taiwan, and Korea, equivalent to biomedicine.^{3,19-25} These countries feature
distinctive systems of education and licensure for EM doctors who are legally permitted to
practice the whole range of modalities of EM, independent from medical doctors of
biomedicine. The EM doctors in Taiwan and Korea even hold the right to practice all the EM
modalities in such an exclusive way that doctors of biomedicine do not hold the right to
practice any of the EM modalities. Reciprocally, these EM doctors are not allowed to practice
biomedicine. In China, however, biomedical doctors are allowed to practice any EM
modalities as the doctors deem necessary for medical treatments. In return, EM doctors are
entitled to practice biomedicine as well. These countries show that a whole system of CAM is
institutionalized within the dominant biomedical system in several different ways.

1
2
3
4 Drawing on the literature of medical systems, this paper posits that these institution-level
5 variations are consequential for the extent to which CAM users utilize various treatment
6 modalities of a CAM system. The paper then examines users' utilization behavior by looking
7 at the extent to which CAM users utilize multiple treatment modalities within a CAM system
8 holistically. In accordance with previous studies,⁵⁻⁸ the holistic use is defined as CAM users'
9 utilization behavior in which two or more treatment modalities of a CAM system are used
10 together rather than a single modality being used in isolation from the other modalities. When
11 a user resorts to more modalities, its behavior is interpreted to be more holistic. The paper
12 pursues this investigation by examining the case of EM that is one of the popular CAM
13 practices in the world. In particular, it develops a specific hypothesis that elaborates the
14 unique institutional condition of EM in Korea and relates it to the utilization behavior of EM
15 users who may also use biomedical services concomitantly in the Korean health care system.
16 Therefore, the hypothesis focuses on how the use of biomedical services is associated with
17 the holistic use of EM in the Korean context.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

37 In Korea, the professional practice of EM is comprehensively and exclusively entitled to EM
38 doctors in separation from the professional practice of biomedicine by medical doctors. This
39 comprehensive sanction of all treatment modalities of EM within the medical system and, at
40 the same time, the exclusive entitlement of the practice rights of these modalities only to EM
41 doctors (and not biomedical doctors) likely project EM to Korean medical service users as a
42 system of medical practices that is very different and independent from biomedicine. EM can
43 be also viewed as a whole medical system that is composed of a variety of related treatment
44 modalities that are readily available for the needs of medical service users. In this
45 institutional condition, EM users who co-utilize biomedical services as well are likely those
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 who search for diverse medical resources of different kinds that the national medical system
5 provides for them. These EM users, when compared to EM users who do not use biomedical
6 services and thus do not seek for diverse medical resources, are likely to seek even more
7
8 diverse modalities that are available within EM itself. Therefore, this paper hypothesizes that
9
10 the EM users who also use biomedical services are likely to use EM more holistically.
11
12
13

14
15
16
17 *Hypothesis: Among EM users in Korea where certified EM professionals hold the*
18 *comprehensive and exclusive practice rights over EM, medical service users' use of*
19 *biomedical services is positively associated with their likelihood of using EM holistically.*
20
21
22
23
24

25 26 **Methods**

27
28
29
30 Data come from the 2011 Korean National Survey of EM Patients (NSEMP) that was
31 administered to a nationally representative sample of patients who visited (i.e. outpatients) or
32 were hospitalized (i.e. inpatients) in an EM facility as of September 2011. This survey used
33 the national sampling frame of 12,250 EM facilities that were registered in the national health
34 insurance system. This sampling frame was duly regarded as the national population of EM
35 facilities in Korea, since all medical service providers should be registered in the national
36 insurance system for reimbursement from the national government in the universal Korean
37 health care system. The survey then drew a stratified systematic sample of 471 EM facilities
38 (4% of the sampling frame). At each selected facility, the outpatient questionnaire of the
39 survey was administered to a random sample of 9 outpatients drawn from people in the
40 waiting area. When the facility was equipped with hospital beds, an additional random
41 sample of 8 inpatients was drawn from the list of inpatients and these inpatients participated
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 in the inpatient questionnaire. As a result, 3,926 outpatients and 1,581 inpatients participated
5
6 in the survey.
7
8
9

10 This paper analyzes only the responses from the outpatient EM users and not those from the
11
12 inpatient users, since only the outpatient questionnaire investigated the uses of EM in each of
13
14 the seven different EM modalities in detail; the inpatient questionnaire investigated whether a
15
16 respondent ever used any of the EM modalities without differentiating which modality was
17
18 used. These inpatient responses are ignorant of which and how many EM modalities were
19
20 used and, thus, how holistic the EM use was. As a result, the following analysis includes
21
22 responses from the final sample of 3,861 outpatient EM users. Its difference from the total
23
24 outpatient participants ($65 = 3,926 - 3,861$) is due to a further exclusion of 65 inpatient
25
26 respondents that are missing values in one or more of the variables that are included in the
27
28 following analysis.
29
30
31
32
33
34

35 The dependent variable, *the holistic use of EM*, is an ordinal variable that measures the
36
37 number of different EM modalities that were used together by a respondent in the past three
38
39 months. It is a composite measure that summarizes responses to seven distinct questions.
40
41 Each of the seven questions asked whether a respondent used one of the seven EM modalities
42
43 respectively (“have you used [a specific EM modality] for medical problems in the past three
44
45 months?”), such as 1) herbal extracts, 2) herbal pills/powders, 3) acupuncture, 4) moxibustion,
46
47 5) cupping, 6) chuna, and 7) manual treatments. The response to each question is coded 1 if
48
49 yes (0 if not). Thus, the values of the dependent variable range from 1 to 7. The focal
50
51 independent variable is *the use of biomedical services* which measures whether a respondent
52
53 visited a biomedical clinic or hospital where biomedical doctors provide medical services for
54
55
56
57
58
59
60

1
2
3
4 the medical conditions for which the respondent used EM modalities. It is coded 1 if a
5
6 respondent visited a biomedical clinic or hospital in the past (0 if not).
7
8
9

10
11 [Table 1]
12
13

14
15 A set of potential covariates, which may intervene in the relationship between holistic EM
16
17 use and the use of biomedical services, are incorporated in the analysis as control variables.
18
19 These control variables are the frequency of EM use, self-rated health status, gender, age,
20
21 marital status, the highest level of education, and monthly household income (Table 1 for
22
23 descriptive statistics). This paper uses ordered logistic regression models to test the
24
25 hypothesis about the relationship between holistic EM use and the use of biomedical services,
26
27 since it interprets different values in the dependent variable as ordered categories that refer to
28
29 the extent of holistic EM use. Results from negative binomial regression models, which treat
30
31 the dependent variable as a count measure, agree with the results reported here. For
32
33 comparison, results from negative binomial regression models are reported in Table A1 in the
34
35 Appendix.
36
37
38
39

40 41 42 **Results** 43 44 45

46 Among all current outpatient EM users who used at least one EM modality in the past three
47
48 months, EM users who used two or three modalities are the two greatest in number, followed
49
50 by those who used four, five, six, and seven modalities. The proportion of EM users who
51
52 used only one modality is only 9.4%. It turns out that more than 90% of current EM
53
54 outpatients in Korea used multiple EM modalities together when they ever resorted to EM.
55
56
57
58
59
60

1
2
3
4
5
6 Close to a half of these EM users (42.5%) also used biomedical services by visiting a
7
8 biomedical clinic or hospital. In addition, the bivariate tabulation on the relationship between
9
10 the extent of holistic EM use and the use of biomedical services shows that there is a positive
11
12 relationship between the two; EM users who utilized various EM modalities more holistically
13
14 were more likely to use biomedical services as well (Table 2). Put differently, EM users who
15
16 also used biomedical services were more likely to utilize EM modalities holistically than EM
17
18 users who did not use biomedical services.
19
20

21
22
23
24 [Table 2]
25
26

27
28 [Table 3]
29
30

31
32 Model 1 in Table 3 puts this bivariate association in odds ratio and finds it to be statistically
33
34 significant. The odds for EM users who also used biomedical services to utilize EM
35
36 holistically are greater than those for EM users who did not use biomedical services
37
38 (OR=1.27; 95% CI=1.13–1.42). The following two models show that this association remains
39
40 the same when respondents' frequency of EM use is controlled (Model 2) or when
41
42 respondents' health status is controlled (Model 3). The final Model 4 incorporates these two
43
44 confounders together and other possible confounders as control variables; the positive
45
46 association between holistic EM use and the use of biomedical services still persists. In this
47
48 final model, the odds for EM users who also used biomedicine to use EM holistically are
49
50 17% greater than those for EM users who did not use biomedicine (OR=1.17; 95% CI=1.04–
51
52 1.31). These results support the hypothesis unambiguously.
53
54
55
56
57

Discussion

To our knowledge, this is the first study in the global medical community that examines how the use of biomedical services is associated with the extent to which the users of CAM, involving EM, utilize various modalities of a CAM system holistically. It is surprising that this line of inquiry has been neglected in the literature, in spite of a potential link between the holistic-vs-selective utilization of a CAM system and its varying health care outcomes. While awaiting more empirical evidence on this link, this paper contributes to developing a much-needed analytical perspective that elaborates CAM utilization behavior beyond existent studies of the simple utilization rate and popularity of CAM among various populations.²⁵⁻³¹

As such, this perspective has grown out of a group of studies that investigate the complementary-versus-substitutive relationship between CAM use and biomedicine use.^{23,32-}

³⁵ Whereas these existing studies have investigated whether the utilization of fragmented CAM modalities increases or decreases the utilization of biomedical services by comparing the behavior of CAM users to the behavior of non-users, this paper originally focuses on CAM users only and investigates how their CAM utilization behavior is shaped by their use (vs. non-use) of biomedical services. In this sense, this paper provides CAM-centered evidence to the complementarity-versus-substitution debate; its finding suggests that CAM use can be complementary to biomedicine to the extent that users rely on a variety of treatment modalities within a CAM system even when they use biomedical services (i.e. the holistic co-utilization of CAM).

Conclusions

This finding is based upon the experiences of EM users in Korea where the professional practice of EM is comprehensively and exclusively entitled to EM doctors. It will take more research to generalize this finding in other institutional contexts and, at the same time, to discern any cross-national differences. The qualifications and characterizations that this paper has developed regarding the Korean medical system will certainly serve future research interest in this direction. A comparative study between the East Asian countries, such as China, Korea, and Japan where EM originated and is institutionalized differently, will shed more light.

The authors recommend that the CAM-centered perspective, asking how CAM use is reconstructed by biomedicine use, is especially relevant to studying CAM utilization behavior in societies where a system of CAM treatment modalities has existed for quite some time (e.g. traditional Indian medicine in the U.S., traditional African medicine in Africa, Ayurveda and Indian medical traditions in India, etc.). The international medical community will gain a lot from these future studies on their own accord. It will gain even more from these studies when the link between the holistic-versus-selective use of CAM and its varying health care outcomes is examined further.

Appendix

[Table A1]

Contributorship Statement

JMS conceived the study, analyzed the data, and wrote the manuscript. YSL conceived the study and helped modify the data analysis and revise the manuscript.

Acknowledgements

This work was supported by the 2014 Research Fund of the University of Seoul. The views in this paper are the authors' and should not be interpreted as those of the supporter. The authors thank Keith Woogerd for his copyediting the manuscript.

Competing Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The first and corresponding author received financial support from the University of Seoul for data analysis and manuscript writing.

Data Sharing Statement

The data will be available from the corresponding author upon request.

References

1. WHO. *WHO Traditional Medicine Strategy 2014-2023*. World Health Organization;2013.
2. Bodeker G, Ong C-K, Grundy C, Burford G, Shein K. *WHO Global Atlas of Traditional, Complementary and Alternative Medicine*. Kobe, Japan: WHO Centre for Health Development; 2005.
3. Xu J, Yang Y. Traditional Chinese Medicine in the Chinese Health Care System. *Health Policy*. 2009;90(2-3):133-139.
4. Kaboru BB, Falkenberg T, Ndulo J, Muchimba M, Solo K, Faxelid E. Communities' Views on Prerequisites for Collaboration between Modern and Traditional Health Sectors in Relation to STI/HIV/AIDS Care in Zambia. *Health Policy*. 2006;78(2-3):330-339.
5. MacPherson H, Nahin R, Paterson C, Cassidy CM, Lewith GT, Hammerschlag R. Developments in Acupuncture Research: Big-picture Perspectives from the Leading Edge. *J Altern Complement Med*. 2008;14(7):883-887.
6. Ritenbaugh C, Verhoef M, Fleishman S. Whole Systems Research: A Discipline for Studying Complementary and Alternative Medicine. *Altern Ther Health Med*. 2003;9:32-36.
7. Cao Y, Zhan H, Pang J, et al. Individually Integrated Traditional Chinese Medicine Approach in the Management of Knee Osteoarthritis: Study Protocol for a Randomized Controlled Trial. *Trials*. 2011;12:160.
8. Shim J-M, Kim J. Cross-national Differences in the Holistic Use of Traditional East Asian Medicine in East Asia. *Health Promot Int*. 2016. Doi: 10.1093/heapro/daw089.

- 1
2
3
4 9. Lock M. *East Asian Medicine in Urban Japan: Varieties of Medical Experience*. Vol
5 no. 4. Berkeley: University of California Press; 1980.
6
7
- 8
9 10. Lock M. The Organization and Practice of East Asian Medicine in Japan: Continuity
10 and Change. *Soc Sci Med*. 1980;14B(4):245-253.
11
- 12
13 11. Katayama K, Yoshino T, Munakata K, et al. Prescription of Kampo Drugs in the
14 Japanese Health Care Insurance Program. *Evid Based Complement Alternat Med*.
15 2013;2013:576973.
16
17
- 18
19 12. Kobayashi A, Uefuji M, Yasumo W. History and Progress of Japanese Acupuncture.
20 *Evid Based Complement Alternat Med*. 2010;7(3):359-365.
21
- 22
23 13. Tsutani K. The Evaluation of Herbal Medicines: an East Asian Perspective. In:
24 Lewith GT, Aldridge D, eds. *Clinical Research Methodology for Complementary*
25 *Therapies*. London: Hodder & Stoughton; 1993:365-393.
26
27
- 28
29 14. Barnes LL. The Acupuncture Wars: the Professionalizing of American Acupuncture.
30 *Med Anthropol*. 2003;22(3):261-301.
31
32
- 33
34 15. Wolpe PR. The Maintenance of Professional Authority: Acupuncture and the
35 American Physician. *Soc Probl*. 1985;32(5):409-424.
36
37
- 38
39 16. Braverman C, Baker C, Harris R. Acupuncture and Oriental Medicine (AOM) in the
40 United States. *Am Acupuncturist*. 2009;47:22-26.
41
42
- 43
44 17. Givati A, Hatton K. Traditional acupuncturists and higher education in Britain: The
45 dual, paradoxical impact of biomedical alignment on the holistic view. *Soc Sci Med*.
46 2015;131:173-180.
47
48
- 49
50 18. Ramsey M. Alternative Medicine in Modern France. *Medical Hist*. 1999;43(3):286.
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 19. Holliday I. Traditional Medicines in Modern Societies: An Exploration of
5
6 Integrationist Options through East Asian Experience. *J Med Philos.* 2003;28(3):373-
7
8 389.
9
- 10
11 20. Yoon K-J, Kim D-S. *Ui-Hanui Kongsangbalchunul Wihan Tonghapchok*
12
13 *Chopkeunbangsik Mosaek (Ways of Integrating Oriental Medicine and Western*
14
15 *Medicine)*. Korea Institute of Health and Social Affairs;2013.
16
- 17
18 21. Xu H, Chen K. Integrative Medicine: the Experience from China. *J Altern*
19
20 *Complement Med.* 2008;14(1):3-7.
21
- 22
23 22. Shin D. How Four Different Political Systems Have Shaped the Modernization of
24
25 Traditional Korean Medicine between 1900 and 1960. *Hist Sci.* 2008;17(3):225-241.
26
- 27
28 23. Shim J-M. The Relationship Between the Use of Complementary and Alternative
29
30 Medicine and the Use of Biomedical Services: Evidence From East Asian Medical
31
32 Systems. *Asia Pac J Public Health.* 2016;28(1):51-60.
33
- 34
35 24. Yu F, Takahashi T, Moriya J, et al. Traditional Chinese Medicine and Kampo: a
36
37 Review from the Distant Past for the Future. *J Int Med Res.* 2006;34(3):231-239.
38
- 39
40 25. Frass M, Strassl RP, Friehs H, Müllner M, Kundi M, Kaye AD. Use and Acceptance
41
42 of Complementary and Alternative Medicine Among the General Population and
43
44 Medical Personnel: A Systematic Review. *Ochsner J.* 2012;12(1):45-56.
45
- 46
47 26. Harris PE, Cooper KL, Relton C, Thomas KJ. Prevalence of Complementary and
48
49 Alternative Medicine (CAM) Use by the General Population: a Systematic Review
50
51 and Update. *Int J Clin Pract.* 2012;66(10):924-939.
52
- 53
54 27. Wai WT, Lan WS, Donnan SPB. Prevalence and Determinants of the Use of
55
56 Traditional Chinese Medicine in Hong Kong. *Asia Pac J Public Health.*
57
58 1995;8(3):167-170.
59
60

- 1
2
3
4 28. Chen FP, Chen TJ, Kung YY, et al. Use Frequency of Traditional Chinese Medicine
5 in Taiwan. *BMC Health Serv Res.* 2007;7:26.
6
7
8
9 29. Hori S, Mihaylov I, Vasconcelos J, McCoubrie M. Patterns of Complementary and
10 Alternative Medicine Use amongst Outpatients in Tokyo, Japan. *BMC Complement*
11 *Altern Med.* 2008;8(1):14.
12
13
14 30. Hong CD. Complementary and Alternative Medicine in Korea: Current Status and
15 Future Prospects. *J Altern Complement Med.* 2001;7(6):33-40.
16
17
18 31. Jin L. From Mainstream to Marginal? Trends in the Use of Chinese Medicine in
19 China from 1991 to 2004. *Soc Sci Med.* 2010;71(6):1063-1067.
20
21
22 32. Druss BG, Rosenheck RA. Association Between Use of Unconventional Therapies
23 and Conventional Medical Services. *JAMA.* 1999;282(7):651-656.
24
25
26 33. van Gameren E. Health Insurance and Use of Alternative Medicine in Mexico. *Health*
27 *Policy.* 2010;98(1):50-57.
28
29
30 34. Ayers SL, Kronenfeld JJ. Delays in Seeking Conventional Medical Care and
31 Complementary and Alternative Medicine Utilization. *Health Serv Res.*
32 2012;47(5):2081-2096.
33
34
35 35. Pagán JA, Pauly MV. Access to Conventional Medical Care and the Use of
36 Complementary and Alternative Medicine. *Health Aff.* 2005;24(1):255-262.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. Descriptive Statistics¹⁾ of Variables Used (N=3,861)

Variable	N	%	Variable	N	%
The Holistic Use of EM			Age		
1	363	9.4	less than 20	143	3.7
2	1,284	33.3	20s or 30s	894	23.2
3	1,135	29.4	40s	762	19.7
4	699	18.1	50s	833	21.6
5	276	7.2	60s	575	14.9
6	82	2.1	70s	522	13.5
7	22	0.6	80s or more	132	3.4
The Use of Biomedical Services			Marital Status		
Yes	1,641	42.5	Single	613	15.9
No	2,220	57.5	Married	2,783	72.1
EM Use Frequency ²⁾			Widowed/Divorced/Separated		
1 to 3 times	1,389	36.0	Others	440	11.4
4 to 10 times	1,289	33.4	Education		
11 to 30 times	885	22.9	No formal education	279	7.2
30 times or more	298	7.7	Elementary school	557	14.4
Self-Rated Health Status			Middle school	467	12.1
Very bad	103	2.7	High school	1,215	31.5
Bad	1,010	26.2	College or graduate school	1,343	34.8
Fair	1,651	42.8	Household Income		
Good	983	25.5	less than 1 million wons	733	19.0
Very good	114	3.0	1 to 1.9 million wons	789	20.4
Gender			2 to 2.9 million wons	831	21.5
Male	1,161	30.1	3 to 3.9 million wons	615	15.9
Female	2,700	69.9	4 to 4.9 million wons	446	11.6
			5 million wons or more	447	11.6

Note: ¹⁾ The socio-demographic characteristics of EM users show that EM is being used very widely across different social groups in age, education, and income. ²⁾ This refers to how many times a respondent has used EM services in the past three months. It is certain that, when a respondent uses EM less often, the respondent is less likely to use different treatment modalities of EM. However, a respondent who uses EM very often does not necessarily utilize different treatment modalities (e.g. a respondent who uses only acupuncture many times).

Table 2. The Bivariate Association between the Use of Biomedical Services and the Holistic Use of Traditional East Asian Medicine (EM)

	The Use of Biomedical Services			
	No	Yes	Total	
The Holistic Use of EM	1	227 (62.5)	136 (37.5)	363 (100.0)
(# of EM Modalities Used)	2	766 (59.7)	518 (40.3)	1284 (100.0)
	3	654 (57.6)	481 (42.4)	1135 (100.0)
	4	389 (55.6)	310 (44.4)	699 (100.0)
	5	138 (50.0)	138 (50.0)	276 (100.0)
	6	39 (47.6)	43 (52.4)	82 (100.0)
	7	7 (31.8)	15 (68.2)	22 (100.0)
Total		2,220 (57.5)	1641 (42.5)	3,861 (100.0)

Note: Percentages in parentheses. Pearson's χ^2 (6) = 22.8001; p-value = 0.001.

For peer review only

Table 3. Ordered Logistic Regression Models of Holistic EM Use upon the Use of Biomedical Services and Other Covariates

	Model 1		Model 2		Model 3		Model 4	
	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
Biomedical Service Use	1.27***	(1.13-1.42)	1.18**	(1.05-1.33)	1.23***	(1.09-1.38)	1.17**	(1.04-1.31)
EM Use Frequency (Ref= 1 to 3 times)								
4 to 10 times			2.29***	(2.00-2.64)			2.30***	(2.00-2.65)
11 to 30 times			3.38***	(2.90-3.96)			3.45***	(2.93-4.06)
31 or more times			3.40***	(2.71-4.27)			3.53***	(2.78-4.47)
Self-Rated Health Status (Ref= Very good)								
Very bad					0.88	(0.54-1.43)	0.58*	(0.35-0.95)
Bad					1.32	(0.93-1.88)	0.92	(0.64-1.32)
Fair					1.10	(0.78-1.55)	0.84	(0.59-1.19)
Good					0.91	(0.64-1.29)	0.78	(0.55-1.11)
Female							1.09	(0.96-1.24)
Age (Ref= less than 20)								
20s or 30s							2.96***	(1.98-4.44)
40s							3.28***	(2.13-5.05)
50s							3.06***	(2.00-4.70)
60s							2.45***	(1.58-3.79)
70s							2.29***	(1.46-3.58)
80s or more							2.33**	(1.37-3.97)
Marital Status (Ref= Single)								
Married							1.12	(0.91-1.39)
Widowed/Divorced/Separated							1.13	(0.83-1.52)
Others							1.08	(0.50-2.34)
Education (Ref= No formal education)								
Elementary school							1.05	(0.80-1.38)
Middle school							0.92	(0.68-1.25)
High school							0.97	(0.72-1.31)
College or graduate school							0.90	(0.66-1.24)
Household Income (Ref= less than 1 million wons)								
1 to 1.9 million wons							1.00	(0.82-1.22)
2 to 2.9 million wons							0.95	(0.77-1.17)
3 to 3.9 million wons							1.02	(0.81-1.28)
4 to 4.9 million wons							1.05	(0.82-1.35)
5 million wons or more							1.19	(0.92-1.54)
Observations			3,861		3,861		3,861	

Note: significant at 0.05 (*), 0.01 (**), 0.001 (***).

Table A1. The Replication of Table 3 in Negative Binomial Regression Models

	Model 1		Model 2		Model 3		Model 4	
	coeff.	(95% CI)	coeff.	(95% CI)	coeff.	(95% CI)	coeff.	(95% CI)
Biomedical Service Use	0.06**	(0.02-0.10)	0.04*	(0.01-0.08)	0.05**	(0.02-0.09)	0.04*	(0.01-0.08)
EM Use Frequency (Ref= 1 to 3 times)								
4 to 10 times			0.19***	(0.14-0.23)			0.18***	(0.14-0.23)
11 to 30 times			0.27***	(0.22-0.32)			0.28***	(0.22-0.33)
31 or more times			0.27***	(0.20-0.35)			0.28***	(0.21-0.36)
Self-Rated Health Status (Ref= Very good)								
Very bad					-0.02	(-0.18-0.14)	-0.11	(-0.27-0.06)
Bad					0.06	(-0.06-0.17)	-0.02	(-0.14-0.10)
Fair					0.02	(-0.09-0.13)	-0.04	(-0.15-0.08)
Good					-0.02	(-0.13-0.10)	-0.05	(-0.17-0.07)
Female							0.02	(-0.02-0.06)
Age (Ref= less than 20)								
20s or 30s							0.24***	(0.10-0.38)
40s							0.26***	(0.11-0.41)
50s							0.25***	(0.10-0.39)
60s							0.21**	(0.05-0.36)
70s							0.18*	(0.03-0.33)
80s or more							0.17	(-0.01-0.35)
Marital Status (Ref= Single)								
Married							0.03	(-0.04-0.10)
Widowed/Divorced/Separated							0.03	(-0.06-0.13)
Others							-0.02	(-0.32-0.28)
Education (Ref= No formal education)								
Elementary school							0.00	(-0.09-0.09)
Middle school							-0.02	(-0.12-0.08)
High school							-0.01	(-0.11-0.09)
College or graduate school							-0.02	(-0.12-0.09)
Household Income (Ref= less than 1 million wons)								
1 to 1.9 million wons							0.01	(-0.06-0.07)
2 to 2.9 million wons							-0.00	(-0.07-0.06)
3 to 3.9 million wons							0.00	(-0.07-0.08)
4 to 4.9 million wons							0.02	(-0.06-0.10)
5 million wons or more							0.04	(-0.05-0.12)
Constant	1.04***	(1.01-1.06)	0.89***	(0.85-0.93)	1.02***	(0.91-1.13)	0.67***	(0.49-0.85)
Observations		3,861		3,861		3,861		3,861

Note: significant at 0.05 (*), 0.01 (**), 0.001 (***)

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	1
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	2 to 4
Objectives	3	State specific objectives, including any prespecified hypotheses	4 to 5
Methods			
Study design	4	Present key elements of study design early in the paper	5 to 6
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	5 to 6
Participants	6	(a) <i>Cohort study</i> —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	n.a.
		<i>Case-control study</i> —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls	n.a.
		<i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	5 to 6
		(b) <i>Cohort study</i> —For matched studies, give matching criteria and number of exposed and unexposed	n.a.
		<i>Case-control study</i> —For matched studies, give matching criteria and the number of controls per case	n.a.
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	6 to 7
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	6 to 7
Bias	9	Describe any efforts to address potential sources of bias	6
Study size	10	Explain how the study size was arrived at	6
Quantitative variables	11	(a) Describe all statistical methods, including those used to control for confounding	7
		(b) Describe any methods used to examine subgroups and interactions	n.a.
		(c) Explain how missing data were addressed	6
		(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed	n.a.
		<i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed	n.a.
Statistical methods	12	<i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	n.a.
		(e) Describe any sensitivity analyses	7

Continued on next page

Results

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	6
		(b) Give reasons for non-participation at each stage	6
		(c) Consider use of a flow diagram	n.a.
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	7
		(b) Indicate number of participants with missing data for each variable of interest	6
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	n.a.
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	n.a.
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	n.a.
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	7
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	8
		(b) Report category boundaries when continuous variables were categorized	7
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	n.a.
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	10
Discussion			
Key results	18	Summarise key results with reference to study objectives	9
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	10
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	9 to 10
Generalisability	21	Discuss the generalisability (external validity) of the study results	10
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	11

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.

BMJ Open

The Association between the Use of Biomedical Services and the Holistic Use of Traditional East Asian Medicine: a National Survey of Outpatients in South Korea

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-018414.R1
Article Type:	Research
Date Submitted by the Author:	30-Oct-2017
Complete List of Authors:	Shim, Jae-Mahn; Korea University, Sociology Lee, Yun-Suk; University of Seoul, Sociology
Primary Subject Heading:	Complementary medicine
Secondary Subject Heading:	Sociology, Health services research, Public health
Keywords:	holism, co-utilization, East Asian medicine, Korea, utilization behavior, institutionalization

SCHOLARONE™
Manuscripts

Peer Review Only

1
2
3
4 **The Association between the Use of Biomedical Services and the Holistic Use of**
5 **Traditional East Asian Medicine: a National Survey of Outpatients in South Korea**
6
7

8
9 **Authors**

10
11 1. Jae-Mahn Shim, Ph.D. (*Corresponding author)

12 Department of Sociology

13 Korea University

14 145 Anam-Ro, Seoungbuk-Gu

15 Seoul 02841, Korea

16 Email: jaemahn.shim@gmail.com

17 Phone: +82-2-3290-2088

18 Fax: +82-2-953-2142

19
20
21
22
23 2. Yun-Suk Lee, Ph.D.

24 Department of Sociology

25 University of Seoul

26 163 Seoulsiripdaero, Dongdaemungu

27 Seoul 130-742, Korea

28 Email: yslee@uos.ac.kr

29
30
31
32 **Keywords:** holism; co-utilization; East Asian medicine; Korea; utilization behavior

33
34
35 **Word count:** 3,769

Abstract

Objectives: The holistic use of a system of complementary and alternative medicine (CAM) is potentially linked to its treatment outcomes. This paper examines how the use of biomedicine is associated with the holistic use of CAM, focusing on traditional East Asian medicine (EM) that is uniquely integrated in the medical system in South Korea.

Design/Settings: A representative national sample of EM outpatients in South Korea.

Participants: 3,861 survey respondents.

Methods: By using the 2011 Korean National Survey of EM Patients, ordered logistic regression models specify the relationship between EM outpatients' use of biomedicine and their holistic use of EM modalities.

Results: Among EM outpatients who used at least one EM modality in the past three months, people who used two (33.3%) or three (29.4%) modalities together are the two greatest in number, followed by users of four (18.1%), five (7.2%), six (2.1%), and seven (0.6%) modalities. The odds for EM users to use EM holistically are 17% greater among EM users who used biomedicine as well, compared to EM users who did not use biomedicine.

Conclusions: Health care community should recognize that CAM use likely becomes holistic as people use biomedicine concomitantly, when the practice rights over a CAM system are comprehensively and exclusively entitled to a group of CAM professionals who are independent from practitioners of biomedicine.

[Strengths and limitations of this study]

- It uses a national sample of EM users who visited EM facilities across South Korea.
- It specifies several multivariate ordered logistic regression models to support the findings.
- All the measures are based upon self-reports of survey participants.

Introduction

The presence of complementary and alternative medicine (CAM) is substantial in contemporary health care systems around the world.^{1,2} The health care community is concerned with the ways in which CAM is integrated in the mainstream biomedicine^{3,4} and subsequent health care outcomes that CAM produces. Studies suggest that CAM services be provided holistically so that CAM users can utilize all the related treatment modalities within a whole system of CAM.^{5,6} In the literature, the holistic use of a CAM system refers to the utilization behavior of CAM users who use two or more treatment modalities together that constitute the CAM system.^{7,8} In the selective and fragmented use, on the contrary, people use only a certain modality of the CAM system and not the other modalities. It is argued that the holistic use can maximize the treatment effects of CAM, compared to the selective and fragmented use of only one modality out of multiple interrelated modalities in the whole system.⁸

This paper aims to investigate what generates this difference in the behavior of CAM users. Drawing on the literature of medical systems, the paper posits that the ways in which CAM is institutionalized in medical systems are consequential for the extent to which CAM users utilize various treatment modalities of a CAM system holistically. In accordance with previous studies,⁵⁻⁸ the paper defines the holistic use of CAM as CAM users' utilization behavior in which two or more treatment modalities of a CAM system are used together rather than a single modality being used in isolation from the other modalities of the CAM system. When a user resorts to more modalities, its behavior is interpreted to be more holistic.

1
2
3
4 The paper pursues this investigation by examining the case of a whole system of traditional
5
6 East Asian medicine (EM) that is composed of multiple treatment modalities, such as
7
8 acupuncture, moxibustion, cupping, herbal remedies, and acupressure. In particular, the paper
9
10 develops a specific hypothesis that elaborates the unique institutional condition of EM in
11
12 South Korea and relates it to the utilization behavior of EM users who may also use
13
14 biomedical services concomitantly that are readily available in the national health care system
15
16 of the country. The hypothesis focuses on how the use of biomedical services is associated
17
18 with the holistic use of EM in the Korean context.
19
20
21
22
23

24 In Korea, the professional practice of EM is comprehensively and exclusively entitled to EM
25
26 doctors in separation from the professional practice of biomedicine by medical doctors.⁹⁻¹²
27

28 Korea features distinctive systems of education and licensure for EM doctors who are legally
29
30 permitted to practice the whole range of modalities of EM, independent from medical doctors
31
32 of biomedicine. The EM doctors even hold the right to practice all the EM modalities in such
33
34 an exclusive way that doctors of biomedicine do not hold the right to practice any of the EM
35
36 modalities. Reciprocally, these EM doctors are not allowed to practice biomedicine. This
37
38 comprehensive sanction of all treatment modalities of EM within the medical system and, at
39
40 the same time, the exclusive entitlement of the practice rights of these modalities only to EM
41
42 doctors (and not biomedical doctors) likely project EM to medical service users as a system
43
44 of medical practices that is distinct and independent from biomedicine. EM is also viewed as
45
46 a whole medical system that is composed of a variety of interrelated treatment modalities that
47
48 are readily available for the needs of medical service users.
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 In this institutional condition in Korea, EM users who co-utilize biomedical services as well
5
6 are likely those who search for diverse medical resources of different kinds that the national
7
8 medical system provides for them. These EM users, when compared to EM users who do not
9
10 use biomedical services and thus do not seek for diverse medical resources, are likely to seek
11
12 even more diverse modalities that are available within EM itself. Therefore, this paper
13
14 hypothesizes that the EM users who also use biomedical services are likely to use EM more
15
16 holistically.
17
18
19
20

21
22 *Hypothesis: Among EM users in Korea where certified EM professionals hold the*
23
24 *comprehensive and exclusive practice rights over EM, medical service users' use of*
25
26 *biomedical services is positively associated with their likelihood of using EM holistically.*
27
28
29

30 **Methods**

31
32
33
34
35 Data come from the 2011 Korean National Survey of EM Patients (NSEMP) that was
36
37 administered to a nationally representative sample of patients who visited (i.e. outpatients) or
38
39 were hospitalized (i.e. inpatients) in an EM facility as of September 2011. This survey used
40
41 the national sampling frame of 12,250 EM facilities that were registered in the national health
42
43 insurance system. This sampling frame was duly regarded as the national population of EM
44
45 facilities in Korea, since all medical service providers should be registered in the national
46
47 insurance system for reimbursement from the national government in the universal Korean
48
49 health care system. The survey then drew a stratified systematic sample of 471 EM facilities
50
51 (4% of the sampling frame). At each selected facility, the outpatient questionnaire of the
52
53 survey was administered to a random sample of 9 outpatients drawn from people in the
54
55
56
57
58
59
60

1
2
3
4 waiting area. When the facility was equipped with hospital beds, an additional random
5
6 sample of 8 inpatients was drawn from the list of inpatients and these inpatients participated
7
8 in the inpatient questionnaire. As a result, 3,926 outpatients and 1,581 inpatients participated
9
10 in the survey.
11

12
13
14
15 This paper analyzes only the responses from the outpatient EM users, since only the
16
17 outpatient questionnaire investigated the uses of EM in each of the seven different EM
18
19 modalities in detail; the inpatient questionnaire investigated whether a respondent ever used
20
21 any of the EM modalities without differentiating which modality was used. These inpatient
22
23 responses give no information of which and how many EM modalities were used and, thus,
24
25 how holistic the EM use was. As a result, the following analysis includes responses from the
26
27 final sample of 3,861 outpatient EM users. Its difference from the total outpatient participants
28
29 (65 = 3,926 – 3,861) is due to a further exclusion of 65 inpatient respondents that are missing
30
31 values in one or more of the variables that are included in the following analysis. This study,
32
33 which uses publically available survey data, is granted an exemption from requiring ethics
34
35 approval by the Institutional Review Board of Korea University.
36
37
38

39
40
41
42 The dependent variable, *the holistic use of EM*, is an ordinal variable that measures the
43
44 number of different EM modalities that were used together by a respondent in the past three
45
46 months. It is a composite measure that summarizes responses to seven distinct questions.
47
48 Each of the seven questions asked whether a respondent used one of the seven EM modalities
49
50 respectively (“have you used [a specific EM modality] for medical problems in the past three
51
52 months?”), such as 1) herbal extracts, 2) herbal pills/powders, 3) acupuncture, 4) moxibustion,
53
54 5) cupping, 6) chuna, and 7) manual treatments. The response to each question is coded 1 if
55
56
57
58
59
60

yes (0 if not). Thus, the values of the dependent variable range from 1 to 7. The focal independent variable is *the use of biomedical services* which measures whether a respondent visited a biomedical clinic or hospital where biomedical doctors provide medical services for the medical conditions for which the respondent used EM modalities. It is coded 1 if a respondent visited a biomedical clinic or hospital (0 if not).

Table 1. Descriptive Statistics¹⁾ of Variables Used (N=3,861)

Variable	N	%	Variable	N	%
The Holistic Use of EM			Age		
1	363	9.4	less than 20	143	3.7
2	1,284	33.3	20s or 30s	894	23.2
3	1,135	29.4	40s	762	19.7
4	699	18.1	50s	833	21.6
5	276	7.2	60s	575	14.9
6	82	2.1	70s	522	13.5
7	22	0.6	80s or more	132	3.4
The Use of Biomedical Services			Marital Status		
Yes	1,641	42.5	Single	613	15.9
No	2,220	57.5	Married	2,783	72.1
EM Use Frequency ²⁾			Widowed/Divorced/Separated		
1 to 3 times	1,389	36.0	Others	440	11.4
4 to 10 times	1,289	33.4	Education		
11 to 30 times	885	22.9	No formal education	279	7.2
30 times or more	298	7.7	Elementary school	557	14.4
Self-Rated Health Status			Middle school	467	12.1
Very bad	103	2.7	High school	1,215	31.5
Bad	1,010	26.2	College or graduate school	1,343	34.8
Fair	1,651	42.8	Household Income		
Good	983	25.5	less than 1 million wons	733	19.0
Very good	114	3.0	1 to 1.9 million wons	789	20.4
Gender			2 to 2.9 million wons	831	21.5
Male	1,161	30.1	3 to 3.9 million wons	615	15.9
Female	2,700	69.9	4 to 4.9 million wons	446	11.6
			5 million wons or more	447	11.6

Note: ¹⁾ The socio-demographic characteristics of EM users show that EM is being used very widely across different social groups in age, education, and income. ²⁾ This refers to how many times a respondent has used EM services in the past three months. It is certain that, when a respondent uses EM less often, the respondent is less likely to use different treatment modalities of EM. However, a respondent who uses EM very often does not necessarily utilize different treatment modalities (e.g. a respondent who uses only acupuncture many times).

A set of potential covariates, which may intervene in the relationship between holistic EM use and the use of biomedical services, are incorporated in the analysis as control variables. These control variables are the frequency of EM use, self-rated health status, gender, age, marital status, the highest level of education, and monthly household income (Table 1 for

1
2
3
4 descriptive statistics). This paper uses ordered logistic regression models to test the
5
6 hypothesis about the relationship between holistic EM use and the use of biomedical services,
7
8 since it interprets different values in the dependent variable as ordered categories that refer to
9
10 the extent of holistic EM use. Results from negative binomial regression models, which treat
11
12 the dependent variable as a count measure, agree with the results reported here. For
13
14 comparison, results from negative binomial regression models are reported in Table A1 in the
15
16 Appendix.
17
18
19
20

21 **Results**

22
23
24
25
26 Among all current outpatient EM users who used at least one EM modality in the past three
27
28 months, EM users who used two or three modalities are the two greatest in number, followed
29
30 by those who used four, five, six, and seven modalities. The proportion of EM users who
31
32 used only one modality is only 9.4%. It turns out that more than 90% of current EM
33
34 outpatients in Korea used multiple EM modalities together when they ever resorted to EM.
35
36
37
38

39
40 Close to a half of these EM users (42.5%) also used biomedical services by visiting a
41
42 biomedical clinic or hospital. In addition, the bivariate tabulation on the relationship between
43
44 the extent of holistic EM use and the use of biomedical services shows that there is a positive
45
46 relationship between the two; EM users who utilized various EM modalities more holistically
47
48 were more likely to use biomedical services as well (Table 2). Put differently, EM users who
49
50 also used biomedical services were more likely to utilize EM modalities holistically than EM
51
52 users who did not use biomedical services.
53
54
55
56
57
58
59
60

Table 2. The Bivariate Association between the Use of Biomedical Services and the Holistic Use of Traditional East Asian Medicine (EM)

	The Use of Biomedical Services		
	No	Yes	Total
The Holistic Use of EM	1 227 (62.5)	136 (37.5)	363 (100.0)
(# of EM Modalities Used)	2 766 (59.7)	518 (40.3)	1284 (100.0)
	3 654 (57.6)	481 (42.4)	1135 (100.0)
	4 389 (55.6)	310 (44.4)	699 (100.0)
	5 138 (50.0)	138 (50.0)	276 (100.0)
	6 39 (47.6)	43 (52.4)	82 (100.0)
	7 7 (31.8)	15 (68.2)	22 (100.0)
Total	2,220 (57.5)	1641 (42.5)	3,861 (100.0)

Note: Percentages in parentheses. Pearson's χ^2 (6) = 22.8001; p-value = 0.001.

Table 3. Ordered Logistic Regression Models of Holistic EM Use upon the Use of Biomedical Services and Other Covariates

	Model 1		Model 2		Model 3		Model 4	
	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
Biomedical Service Use	1.27***	(1.13-1.42)	1.18**	(1.05-1.33)	1.23***	(1.09-1.38)	1.17**	(1.04-1.31)
EM Use Frequency (Ref= 1 to 3 times)								
4 to 10 times			2.29***	(2.00-2.64)			2.30***	(2.00-2.65)
11 to 30 times			3.38***	(2.90-3.96)			3.45***	(2.93-4.06)
31 or more times			3.40***	(2.71-4.27)			3.53***	(2.78-4.47)
Self-Rated Health Status (Ref= Very good)								
Very bad					0.88	(0.54-1.43)	0.58*	(0.35-0.95)
Bad					1.32	(0.93-1.88)	0.92	(0.64-1.32)
Fair					1.10	(0.78-1.55)	0.84	(0.59-1.19)
Good					0.91	(0.64-1.29)	0.78	(0.55-1.11)
Female							1.09	(0.96-1.24)
Age (Ref= less than 20)								
20s or 30s							2.96***	(1.98-4.44)
40s							3.28***	(2.13-5.05)
50s							3.06***	(2.00-4.70)
60s							2.45***	(1.58-3.79)
70s							2.29***	(1.46-3.58)
80s or more							2.33**	(1.37-3.97)
Marital Status (Ref= Single)								
Married							1.12	(0.91-1.39)
Widowed/Divorced/Separated							1.13	(0.83-1.52)
Others							1.08	(0.50-2.34)
Education (Ref= No formal education)								
Elementary school							1.05	(0.80-1.38)
Middle school							0.92	(0.68-1.25)
High school							0.97	(0.72-1.31)
College or graduate school							0.90	(0.66-1.24)
Household Income (Ref= less than 1 million wons)								
1 to 1.9 million wons							1.00	(0.82-1.22)
2 to 2.9 million wons							0.95	(0.77-1.17)
3 to 3.9 million wons							1.02	(0.81-1.28)
4 to 4.9 million wons							1.05	(0.82-1.35)
5 million wons or more							1.19	(0.92-1.54)
Observations			3,861		3,861		3,861	
								3,861

Note: significant at 0.05 (*), 0.01 (**), 0.001 (***).

1
2
3
4 Model 1 in Table 3 puts this bivariate association in odds ratio and finds it to be statistically
5 significant. The odds for EM users who also used biomedical services to utilize EM
6
7
8 holistically are greater than those for EM users who did not use biomedical services
9
10 (OR=1.27; 95% CI=1.13–1.42). The two subsequent models show that this association
11
12 remains the same when respondents' frequency of EM use is controlled (Model 2) or when
13
14 respondents' health status is controlled (Model 3). The final Model 4 incorporates these two
15
16 confounders together and other possible confounders as control variables; the positive
17
18 association between holistic EM use and the use of biomedical services still persists. In this
19
20 final model, the odds for EM users who also used biomedicine to use EM holistically are
21
22 17% greater than those for EM users who did not use biomedicine (OR=1.17; 95% CI=1.04–
23
24 1.31). These results support the hypothesis unambiguously.
25
26
27
28
29

30 **Discussion**

31
32
33
34
35 This paper found that people who used biomedicine were more likely to use EM holistically
36
37 in South Korea. Among EM outpatients who used at least one EM modality in the past three
38
39 months, people who used two or three modalities together are the two greatest in number,
40
41 followed by users of four, five, six, and seven modalities. The odds for EM users to use EM
42
43 holistically are greater among EM users who used biomedicine as well, compared to EM
44
45 users who did not use biomedicine. It is a limitation of this study that it used measures based
46
47 on self-reports of survey participants. However, it is worth noting that the paper used a
48
49 national sample of EM users who visited EM facilities across South Korea and specified
50
51 multivariate regression models to show the robustness of these findings.
52
53
54
55
56
57
58
59
60

1
2
3
4 To our knowledge, this is the first study in the global medical community that examines how
5
6 the use of biomedical services is associated with the extent to which the users of CAM utilize
7
8 various modalities of a CAM system holistically. It is surprising that this line of inquiry has
9
10 been neglected in the literature, in spite of a potential link between the holistic-vs-selective
11
12 utilization of a CAM system and its health care outcomes. Furthermore, it is very probable
13
14 that CAM users shape their specific ways of using CAM in relation with biomedical services
15
16 that are available to them. In this sense, this paper contributes to developing a much-needed
17
18 analytical perspective that elaborates CAM utilization behavior beyond existent studies of the
19
20 simple utilization rate and the popularity of CAM.¹³⁻¹⁹
21
22
23
24
25

26 As such, this perspective has grown out of a group of studies that investigate the
27
28 complementary-versus-substitutive relationship between CAM use and biomedicine use.^{9,20-23}
29
30 Whereas these existing studies have investigated whether the utilization of fragmented CAM
31
32 modalities increases or decreases the utilization of biomedical services by comparing the
33
34 behavior of CAM users to the behavior of non-users, this paper originally focuses on CAM
35
36 users only and investigates how their CAM utilization behavior is shaped by their use (vs.
37
38 non-use) of biomedical services. In this sense, this paper provides CAM-centered evidence to
39
40 the complementarity-versus-substitution debate; its finding suggests that CAM use can be
41
42 complementary to biomedicine to the extent that users rely on a variety of treatment
43
44 modalities within a CAM system even when they use biomedical services (i.e. the holistic co-
45
46 utilization of CAM).
47
48
49
50
51
52

53 The findings in this paper suggest that the knowledge of how CAM is institutionalized in a
54
55 medical system can generate reasonable predictions about how CAM users behave. It is
56
57
58
59
60

1
2
3
4 known that the various modalities of EM are disconnected from one another in the Japanese
5
6 medical system so that only herbal remedies are selectively incorporated into the practices of
7
8 biomedical doctors;²⁴⁻²⁸ acupuncture and acupressure are each relegated as pseudo-medicine
9
10 to medical technicians, such as acupuncturists and massage therapists who are permitted to
11
12 practice only acupuncture and massage therapies. These technicians are not allowed to
13
14 practice herbal remedies or other EM modalities. Similar observations are made that the
15
16 otherwise rich practices of acupuncture are truncated and simplified in the dominant
17
18 biomedical health care systems in the U.S.²⁹⁻³¹ and the U.K.³² In the contemporary French
19
20 biomedical system, a variety of CAM systems are reported to become “balkanized” and their
21
22 constituent treatment modalities are torn apart from one another in practice.³³ In these
23
24 institutional contexts, this paper suggest, the intersection of a CAM system with biomedicine
25
26 can result in the fragmented and partial use of the CAM system as people use biomedical
27
28 services concomitantly.
29
30
31
32
33
34

35 **Conclusions**

36
37
38
39 Health care community should recognize that CAM use likely becomes holistic as people use
40
41 biomedicine concomitantly, when the practice rights over a CAM system are
42
43 comprehensively and exclusively entitled to a group of CAM professionals who are
44
45 independent from practitioners of biomedicine. This conclusion is based upon the experiences
46
47 of EM users in Korea. It will take more research to generalize this finding against other
48
49 institutional contexts and, at the same time, to discern any cross-national differences. A
50
51 comparative study between the East Asian countries, such as China, Korea, and Japan where
52
53 EM originated and is institutionalized differently, will shed more light.
54
55
56
57
58
59
60

1
2
3
4
5
6 The authors recommend that the CAM-centered perspective, asking how CAM use is
7
8 reconstructed by biomedicine use, is especially relevant to studying CAM utilization behavior
9
10 in societies where a relatively coherent system of CAM has existed for quite some time (e.g.
11
12 traditional Indian medicine in the U.S., traditional African medicine in Africa, Ayurveda and
13
14 Indian medical traditions in India, etc.). The international medical community will gain a lot
15
16 from these future studies. It will gain even more from the studies when the link between the
17
18 holistic-versus-selective use of CAM and its varying health care outcomes is examined
19
20 further.
21
22
23
24
25

26 **Appendix**

27
28
29
30 [Table A1]
31
32
33
34

35 **Contributorship Statement**

36
37 JMS conceived the study, analyzed the data, and wrote the manuscript. YSL helped conceive
38
39 the study, modify the data analysis, and revise the manuscript.
40
41
42
43

44 **Acknowledgements**

45
46 This work was supported by the Ministry of Education of the Republic of Korea and the
47
48 National Research Foundation of Korea (NRF-2015S1A5A8017517). The views in this paper
49
50 are the authors' and should not be interpreted as those of the supporter. The authors thank
51
52 Keith Woogerd for his copyediting the manuscript.
53
54
55
56
57
58
59
60

Competing Interests

The authors declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The first and corresponding author received financial support from the Ministry of Education of the Republic of Korea and the National Research Foundation of Korea (NRF-2015S1A5A8017517) for data analysis and manuscript writing.

Data Sharing Statement

The data are available from the corresponding author upon request.

References

1. WHO. *WHO Traditional Medicine Strategy 2014-2023*. World Health Organization;2013.
2. Bodeker G, Ong C-K, Grundy C, Burford G, Shein K. *WHO Global Atlas of Traditional, Complementary and Alternative Medicine*. Kobe, Japan: WHO Centre for Health Development; 2005.
3. Xu J, Yang Y. Traditional Chinese Medicine in the Chinese Health Care System. *Health Policy*. 2009;90(2-3):133-139.
4. Kaboru BB, Falkenberg T, Ndulo J, Muchimba M, Solo K, Faxelid E. Communities' Views on Prerequisites for Collaboration between Modern and Traditional Health

- 1
2
3
4 Sectors in Relation to STI/HIV/AIDS Care in Zambia. *Health Policy*. 2006;78(2–
5 3):330-339.
6
7
8
9 5. MacPherson H, Nahin R, Paterson C, Cassidy CM, Lewith GT, Hammerschlag R.
10 Developments in Acupuncture Research: Big-picture Perspectives from the Leading
11 Edge. *J Altern Complement Med*. 2008;14(7):883-887.
12
13 6. Ritenbaugh C, Verhoef M, Fleishman S. Whole Systems Research: A Discipline for
14 Studying Complementary and Alternative Medicine. *Altern Ther Health Med*.
15 2003;9:32-36.
16
17 7. Shim J-M, Kim J. Cross-national Differences in the Holistic Use of Traditional East
18 Asian Medicine in East Asia. *Health Promot Int*. 2016. Doi: 10.1093/heapro/daw089.
19
20 8. Cao Y, Zhan H, Pang J, et al. Individually Integrated Traditional Chinese Medicine
21 Approach in the Management of Knee Osteoarthritis: Study Protocol for a
22 Randomized Controlled Trial. *Trials*. 2011;12:160.
23
24 9. Shim J-M. The Relationship Between the Use of Complementary and Alternative
25 Medicine and the Use of Biomedical Services: Evidence From East Asian Medical
26 Systems. *Asia Pac J Public Health*. 2016;28(1):51-60.
27
28 10. Holliday I. Traditional Medicines in Modern Societies: An Exploration of
29 Integrationist Options through East Asian Experience. *J Med Philos*. 2003;28(3):373-
30 389.
31
32 11. Yoon K-J, Kim D-S. *Ui-Hanui Kongsangbalchunul Wihan Tonghapchok*
33 *Chopkeunbangsik Mosaek (Ways of Integrating Oriental Medicine and Western*
34 *Medicine)*. Korea Institute of Health and Social Affairs;2013.
35
36 12. Shin D. How Four Different Political Systems Have Shaped the Modernization of
37 Traditional Korean Medicine between 1900 and 1960. *Hist Sci*. 2008;17(3):225-241.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4 13. Frass M, Strassl RP, Friehs H, Müllner M, Kundi M, Kaye AD. Use and Acceptance
5
6 of Complementary and Alternative Medicine Among the General Population and
7
8 Medical Personnel: A Systematic Review. *Ochsner J.* 2012;12(1):45-56.
9
- 10
11 14. Harris PE, Cooper KL, Relton C, Thomas KJ. Prevalence of Complementary and
12
13 Alternative Medicine (CAM) Use by the General Population: a Systematic Review
14
15 and Update. *Int J Clin Pract.* 2012;66(10):924-939.
16
- 17
18 15. Wai WT, Lan WS, Donnan SPB. Prevalence and Determinants of the Use of
19
20 Traditional Chinese Medicine in Hong Kong. *Asia Pac J Public Health.*
21
22 1995;8(3):167-170.
23
- 24
25 16. Chen FP, Chen TJ, Kung YY, et al. Use Frequency of Traditional Chinese Medicine
26
27 in Taiwan. *BMC Health Serv Res.* 2007;7:26.
28
- 29
30 17. Hori S, Mihaylov I, Vasconcelos J, McCoubrie M. Patterns of Complementary and
31
32 Alternative Medicine Use amongst Outpatients in Tokyo, Japan. *BMC Complement*
33
34 *Altern Med.* 2008;8(1):14.
35
- 36
37 18. Hong CD. Complementary and Alternative Medicine in Korea: Current Status and
38
39 Future Prospects. *J Altern Complement Med.* 2001;7(6):33-40.
40
- 41
42 19. Jin L. From Mainstream to Marginal? Trends in the Use of Chinese Medicine in
43
44 China from 1991 to 2004. *Soc Sci Med.* 2010;71(6):1063-1067.
45
- 46
47 20. Druss BG, Rosenheck RA. Association Between Use of Unconventional Therapies
48
49 and Conventional Medical Services. *JAMA.* 1999;282(7):651-656.
50
- 51
52 21. van Gameren E. Health Insurance and Use of Alternative Medicine in Mexico. *Health*
53
54 *Policy.* 2010;98(1):50-57.
55
56
57
58
59
60

- 1
2
3
4 22. Ayers SL, Kronenfeld JJ. Delays in Seeking Conventional Medical Care and
5 Complementary and Alternative Medicine Utilization. *Health Serv Res.*
6 2012;47(5):2081-2096.
7
8
9
10
11 23. Pagán JA, Pauly MV. Access to Conventional Medical Care and the Use of
12 Complementary and Alternative Medicine. *Health Aff.* 2005;24(1):255-262.
13
14
15 24. Lock M. *East Asian Medicine in Urban Japan: Varieties of Medical Experience.* Vol
16 no. 4. Berkeley: University of California Press; 1980.
17
18
19 25. Lock M. The Organization and Practice of East Asian Medicine in Japan: Continuity
20 and Change. *Soc Sci Med.* 1980;14B(4):245-253.
21
22
23
24 26. Katayama K, Yoshino T, Munakata K, et al. Prescription of Kampo Drugs in the
25 Japanese Health Care Insurance Program. *Evid Based Complement Alternat Med.*
26 2013;2013:576973.
27
28
29
30
31 27. Kobayashi A, Uefuji M, Yasumo W. History and Progress of Japanese Acupuncture.
32 *Evid Based Complement Alternat Med.* 2010;7(3):359-365.
33
34
35 28. Tsutani K. The Evaluation of Herbal Medicines: an East Asian Perspective. In:
36 Lewith GT, Aldridge D, eds. *Clinical Research Methodology for Complementary*
37 *Therapies.* London: Hodder & Stoughton; 1993:365-393.
38
39
40
41 29. Barnes LL. The Acupuncture Wars: the Professionalizing of American Acupuncture.
42 *Med Anthropol.* 2003;22(3):261-301.
43
44
45
46 30. Wolpe PR. The Maintenance of Professional Authority: Acupuncture and the
47 American Physician. *Soc Probl.* 1985;32(5):409-424.
48
49
50
51 31. Braverman C, Baker C, Harris R. Acupuncture and Oriental Medicine (AOM) in the
52 United States. *Am Acupuncturist.* 2009;47:22-26.
53
54
55
56
57
58
59
60

- 1
2
3
4 32. Givati A, Hatton K. Traditional acupuncturists and higher education in Britain: The
5
6 dual, paradoxical impact of biomedical alignment on the holistic view. *Soc Sci Med*.
7
8 2015;131:173-180.
9
10 33. Ramsey M. Alternative Medicine in Modern France. *Medical Hist*. 1999;43(3):286.
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Table A1. The Replication of Table 3 in Negative Binomial Regression Models

	Model 1		Model 2		Model 3		Model 4	
	coeff.	(95% CI)	coeff.	(95% CI)	coeff.	(95% CI)	coeff.	(95% CI)
Biomedical Service Use	0.06**	(0.02-0.10)	0.04*	(0.01-0.08)	0.05**	(0.02-0.09)	0.04*	(0.01-0.08)
EM Use Frequency (Ref = 1 to 3 times)								
4 to 10 times			0.19***	(0.14-0.23)			0.18***	(0.14-0.23)
11 to 30 times			0.27***	(0.22-0.32)			0.28***	(0.22-0.33)
31 or more times			0.27***	(0.20-0.35)			0.28***	(0.21-0.36)
Self-Rated Health Status (Ref = Very good)								
Very bad					-0.02	(-0.18-0.14)	-0.11	(-0.27-0.06)
Bad					0.06	(-0.06-0.17)	-0.02	(-0.14-0.10)
Fair					0.02	(-0.09-0.13)	-0.04	(-0.15-0.08)
Good					-0.02	(-0.13-0.10)	-0.05	(-0.17-0.07)
Female							0.02	(-0.02-0.06)
Age (Ref = less than 20)								
20s or 30s							0.24***	(0.10-0.38)
40s							0.26***	(0.11-0.41)
50s							0.25***	(0.10-0.39)
60s							0.21**	(0.05-0.36)
70s							0.18*	(0.03-0.33)
80s or more							0.17	(-0.01-0.35)
Marital Status (Ref = Single)								
Married							0.03	(-0.04-0.10)
Widowed/Divorced/Separated							0.03	(-0.06-0.13)
Others							-0.02	(-0.32-0.28)
Education (Ref = No formal education)								
Elementary school							0.00	(-0.09-0.09)
Middle school							-0.02	(-0.12-0.08)
High school							-0.01	(-0.11-0.09)
College or graduate school							-0.02	(-0.12-0.09)
Household Income (Ref = less than 1 million wons)								
1 to 1.9 million wons							0.01	(-0.06-0.07)
2 to 2.9 million wons							-0.00	(-0.07-0.06)
3 to 3.9 million wons							0.00	(-0.07-0.08)
4 to 4.9 million wons							0.02	(-0.06-0.10)
5 million wons or more							0.04	(-0.05-0.12)
Constant	1.04***	(1.01-1.06)	0.89***	(0.85-0.93)	1.02***	(0.91-1.13)	0.67***	(0.49-0.85)
Observations		3,861		3,861		3,861		3,861

Note: significant at 0.05 (*), 0.01 (**), 0.001 (***).

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation	Page No
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract	1
		(b) Provide in the abstract an informative and balanced summary of what was done and what was found	1
Introduction			
Background/rationale	2	Explain the scientific background and rationale for the investigation being reported	2 to 3
Objectives	3	State specific objectives, including any prespecified hypotheses	3 to 4
Methods			
Study design	4	Present key elements of study design early in the paper	4 to 6
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection	4 to 6
Participants	6	(a) <i>Cohort study</i> —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up	n.a.
		<i>Case-control study</i> —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls	n.a.
		<i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants	4 to 6
		(b) <i>Cohort study</i> —For matched studies, give matching criteria and number of exposed and unexposed	n.a.
		<i>Case-control study</i> —For matched studies, give matching criteria and the number of controls per case	n.a.
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable	5 to 6
Data sources/measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group	5 to 6
Bias	9	Describe any efforts to address potential sources of bias	5
Study size	10	Explain how the study size was arrived at	5
Quantitative variables	11	(a) Describe all statistical methods, including those used to control for confounding	6
		(b) Describe any methods used to examine subgroups and interactions	n.a.
		(c) Explain how missing data were addressed	5
		(d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed	n.a.
		<i>Case-control study</i> —If applicable, explain how matching of cases and controls was addressed	n.a.
Statistical methods	12	<i>Cross-sectional study</i> —If applicable, describe analytical methods taking account of sampling strategy	n.a.
		(e) Describe any sensitivity analyses	6

Continued on next page

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60**Results**

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed	5
		(b) Give reasons for non-participation at each stage	5
		(c) Consider use of a flow diagram	n.a.
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders	6
		(b) Indicate number of participants with missing data for each variable of interest	5
		(c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)	n.a.
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time	n.a.
		<i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure	n.a.
		<i>Cross-sectional study</i> —Report numbers of outcome events or summary measures	6
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included	7
		(b) Report category boundaries when continuous variables were categorized	6
		(c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period	n.a.
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses	8
Discussion			
Key results	18	Summarise key results with reference to study objectives	8
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias	8
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence	8 to 9
Generalisability	21	Discuss the generalisability (external validity) of the study results	10
Other information			
Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based	12

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is available at www.strobe-statement.org.