## Supplementary material

Table 1. In-scope ICD9/10 alcohol related diagnoses

Alcohol diagnoses <sup>a</sup>	ICD10 codes	ICD9 codes	
Mental and behavioural disorders			
Acute alcohol intoxication	F10.0; F10.1	303.0–303.03; 305.0–305.03	
Alcohol dependence syndrome	F10.2	303.9–303.93	
Alcohol withdrawal	F10.3; F10.4	291.0; 291.81	
Alcohol psychotic disorder	F10.5	291.3; 291.5	
Alcohol amnesic syndrome	F10.6	291.1	
Residual and late-onset alcohol psychiatric disorder	F10.7	291.2; 291.4	
Other	F10.8; F10.9	291.82; 291.89; 291.9	
Alcohol-related diseases			
Alcoholic pseudo-Cushing syndrome	E24.4	255.0	
Alcoholic nervous system degeneration	G31.2	331.7	
Alcoholic polyneuropathy	G62.1	357.5	
Alcoholic myopathy	G72.1	359.4	
Alcoholic cardiomyopathy	142.6	425.5	
Alcoholic gastritis	K29.2	535.30; 535.31	
Alcoholic liver disease	K70-K70.9	571.0-571.3	
Alcoholic pancreatitis	K86.0	577.1	
'Other' alcohol disorders			
Infant damage due to alcohol <sup>b</sup>	O35.4; P04.3; Q86.0	655.43; 760.71	
Other	R78.0; T51; T51.0; T51.8; T51.9; Y90–Y90.9	790.3; 980.0; 980.8; 980.9; E860.0	
Poisoning	X45; X65; Y15	E86.09; E950.09; E98.05	

Rehabilitation/history of alcohol use disorder Z50.2; Z71.4; Z86.41 V57.89, V65.42, V11.3 Encountering health services due to alcohol problems Z72.1 –

Reprinted with permission from *Heavy maternal alcohol consumption and cerebral palsy in the* offspring by O'Leary, C. M., Watson, I., D'Antoine, H., Stanley, F. and Bower, C. (2012), Developmental Medicine & Child Neurology, 54: 224–230

<sup>&</sup>lt;sup>a</sup> Some women may have more than one diagnosis.

<sup>&</sup>lt;sup>b</sup> Recorded on maternal record.

Table 2. Coding of timing of maternal alcohol-related diagnosis in relation to pregnancy

	Timing of maternal alcohol related diagnosis in relation to pregnancy					
	During	≤1 yr pre-pregnancy	≤1 yr after pregnancy	> yr before pregnancy	>1 yr after pregnancy	
During pregnancy	Yes	Possible	Possible	Possible	Possible	
≤ 1 yr pre pregnancy	No	Yes	Possible	Possible	Possible	
≤ 1 yr after pregnancy	No	No	Yes	Possible	Possible	
>1 yr before pregnancy	No	No	No	Yes	Possible	
>1 yr after pregnancy	No	No	No	No	Yes	

Reprinted with permission from *Heavy maternal alcohol consumption and cerebral palsy in the* offspring by O'Leary, C. M., Watson, I., D'Antoine, H., Stanley, F. and Bower, C. (2012), Developmental Medicine & Child Neurology, 54: 224–230