

Table S4. Percentage of Women Who Continued Index Medication Therapy, Discontinued Index Medication Therapy, and Switched to Other Therapies at the Various Time Points

| Time Point | Medication Cohort, % | | | | |
|--------------|----------------------|-----------------|--------------------|-----------------|---------------------|
| | LARC (n=3,928) | LA (n=3,525) | SARC (n=33,000) | TA (n=1,108) | Total (N=41,561) |
| 3 mo | | | | | |
| Continued | 73.9 | 42.9 | 54.3 | 34.5 | 54.7 |
| Discontinued | 8.7 | 17.0 | 23.7 | 37.4 | 22.1 |
| Switched | 17.4 | 40.1 | 22.0 | 28.2 | 23.3 |
| 6 mo | | | | | |
| Continued | 61.8 | 13.0 | 42.6 | 22.3 | 41.4 |
| Discontinued | 12.0 | 26.0 | 29.7 | 43.1 | 28.1 |
| Switched | 26.3 | 61.0 | 27.7 | 34.6 | 30.6 |
| 1 y | | | | | |
| Continued | (n=3,177) | (n=2,914) | (n=27,021) | (n=811) | (n=33,923) |
| Continued | 52.0 | 3.5 | 33.1 | 14.2 | 31.9 |
| Discontinued | 14.9 | 29.5 | 34.9 | 47.8 | 32.9 |
| Switched | 33.1 | 67.0 | 32.0 | 38.0 | 35.2 |
| 2 y | | | | | |
| Continued | (n=2,103) | (n=2,056) | (n=18,496) | (n=366) | (n=23,021) |
| Continued | 42.4 | 1.6 | 23.6 | 6.0 | 23.1 |
| Discontinued | 17.5 | 29.6 | 40.3 | 55.2 | 37.5 |
| Switched | 40.1 | 68.8 | 36.1 | 38.8 | 39.5 |
| 5 y | | | | | |
| Continued | (n=602) | (n=765) | (n=6,215) | (n=0) | (n=7,582) |
| Continued | 15.9 | 0.7 | 10.9 | N/A | 10.3 |
| Discontinued | 34.2 | 27.7 | 48.6 | N/A | 45.4 |
| Switched | 49.8 | 71.6 | 40.5 | N/A | 44.4 |

Abbreviations: LA, leuprolide acetate; LARC, long-acting reversible contraceptive steroid; N/A, not available; SARC, short-acting reversible contraceptive steroid; TA, tranexamic acid.