

Supplemental Table 1. Cytokine and chemokine levels in culture supernatants of wild-type bone marrow-derived macrophages infected with *Mycoplasma pneumoniae* (*Mp*).

Hours post-infection	4		8		16	
	Mock	<i>Mp</i> -infected	Mock	<i>Mp</i> -infected	Mock	<i>Mp</i> -infected
Cytokine						
IL-1a	1.26*	33.17 ± 0.76	2.19 ± 1.07	249.26 ± 16.22	2.20 ± 1.16	870.86 ± 181.61
IL-1b	1.28*	13.81 ± 3.09	1.28*	376.01 ± 72.74	1.28*	1262.46 ± 223.27
IL-6	.059*	6695.25 ± 355.57	1.44 ± 1.0	14568 ± 1008.05	1.48 ± 0.65	22933**
TNF-a	1.53*	1389.62 ± 424.94	2.04 ± 1.01	1518.43 ± 529.53	4.34 ± 0.56	1300.50 ± 130.23
G-CSF	1.25*	974.62 ± 45.36	1.25*	9475.75 ± 1293.42	1.25*	15882 ± 566.09
IP-10	2.97 ± 1.05	87.30 ± 7.13	10.18 ± 0.73	4047.5 ± 391.66	27.88 ± 6.66	4899.25 ± 301.13
KC	3.20 ± 0.32	6383.25 ± 334.27	6.17 ± 1.92	9631.25 ± 1063.31	5.01 ± 0.98	10982.25 ± 942.22
MCP-1	272.87 ± 11.15	871.92 ± 80.07	407.80 ± 69.82	1782 ± 131.03	654.98 ± 35.03	2380.75 ± 97.02
MIP-1a	124.48 ± 4.67	6889.03 ± 327.58	180.33 ± 8.86	8500.53 ± 792.78	186.4 ± 6.56	8027.78 ± 538.17
MIP-1b	179.67 ± 10.42	3302.00 ± 704.72	272.93 ± 23.23	12553.25 ± 2135.92	324.54 ± 17.48	10332.50 ± 2875.84
MIP-2	63.77 ± 5.30	12972.50 ± 696.30	132.19 ± 6.56	13495.75 ± 826.78	166.89 ± 25.82	13610.25**

*Denotes values that were below limit of detection

**Denotes values that reached multiplex assay detection limit

Values are represented in linear scale and were normalized by subtracting uninfected control values from infected values