

Supplementary Table VI. Predictors of T₂ smoking cessation using complete case analysis

Quit smoking at T ₂ (n=865)					
		Yes (n=194) n (%) or M (SD)	No (n=671) n (%) or M (SD)	Univariable OR (95% CI)	Multivariable OR [^] (95% CI)
Trial allocation	Intervention	115 (59%)	373 (55%)	1.16 (0.84 to 1.61)	1.16 (0.65 to 1.33)
	Control	79 (41%)	298 (45%)		
Site	Liverpool	100 (51%)	350 (52%)	1.03 (0.74 to 1.43)	0.90 (0.59 to 1.37)
	Cambridge	94 (49%)	329 (48%)		
Age (years)	Up to 65 years	71 (37%)	270 (40%)	- Reference -	- Reference -
	66 to 70 years	84 (43%)	295 (44%)	1.08 (0.75 to 1.55)	1.08 (0.75 to 1.57)
	Over 70 years	39 (20%)	106 (16%)	1.49 (0.94 to 2.37)	1.54 (0.97 to 2.46)
Gender	Male	144 (74%)	471 (69%)	0.76 (0.53 to 1.10)	0.73 (0.50 to 1.07)
	Female	50 (26%)	208 (31%)		
Marital group	Married/cohabiting	140 (72%)	458 (68%)	0.86 (0.60 to 1.24)	0.94 (0.64 to 1.38)
	Not married/cohabiting	54 (28%)	220 (32%)		
IMD	Quintile 1 (most deprived)	55 (28%)	218 (32%)	- Reference -	- Reference -
	Quintile 2	25 (13%)	83 (12%)	1.21 (0.70 to 2.10)	1.21 (0.69 to 2.13)
	Quintile 3	34 (18%)	102 (15%)	1.37 (0.83 to 2.25)	1.25 (0.74 to 2.11)
	Quintile 4	37 (19%)	112 (17%)	1.38 (0.85 to 2.24)	1.43 (0.82 to 2.49)
	Quintile 5 (least deprived)	43 (22%)	164 (24%)	1.08 (0.68 to 1.71)	1.12 (0.62 to 2.05)
Lung cancer experience	No	117 (61%)	408 (60%)	0.91 (0.65 to 1.28)	0.93 (0.65 to 1.33)
	Yes	74 (39%)	271 (40%)		
Cancer distress (T₀)⁺		2.27 (0.28) <i>10.01 (2.74)</i>	2.23 (0.28) <i>9.63 (2.78)</i>	1.69 (0.95 to 3.02)	1.99 (1.08 to 3.67)

[^] Adjusted for T₀ cancer distress, recruitment site, gender, age, marital group, deprivation and experience of lung cancer.

⁺ Log_n scores in normal text, original scores in italics (analyses performed using log_n scores).