

Supplemental Data (for publication only)

Supplemental Table 1. Cancer case counts, population estimates, and incidence rates by histology and year of diagnosis, Chinese males

Year of Diagnosis	Population Estimates	All lung Cancer		Small cell carcinoma		NSCLC		Squamous cell carcinoma		Adenocarcinoma		Large cell carcinoma		NSCLC NOS	
		Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)
1990	346,928	132	56.4 (46.4-67.9)	9	3.6 (1.6-6.9)	121	51.3 (41.7-62.2)	31	13.9 (9.0-20.3)	56	22.6 (16.7-29.9)	12	5.4 (2.5-9.9)	22	9.4 (5.6-14.5)
1991	362,532	155	61.0 (51.3-72.0)	14	5.7 (3.0-9.7)	140	54.9 (45.7-65.3)	42	16.5 (11.6-22.7)	53	18.9 (14.1-24.9)	15	6.2 (3.4-10.3)	30	13.2 (8.6-19.2)
1992	378,145	187	67.7 (57.9-78.6)	16	5.1 (2.8-8.3)	166	59.9 (50.7-70.2)	43	15.5 (11.1-21.2)	67	23.6 (18.1-30.2)	18	6.4 (3.7-10.3)	38	14.3 (9.8-19.9)
1993	393,756	177	59.9 (51.0-69.8)	21	7.2 (4.4-11.1)	151	50.3 (42.3-59.3)	38	13.2 (9.1-18.3)	56	17.1 (12.9-22.4)	12	4.2 (2.1-7.4)	45	15.8 (11.3-21.3)
1994	409,370	157	52.3 (44.1-61.6)	19	6.2 (3.6-9.7)	129	41.6 (34.4-49.7)	33	10.6 (7.2-15.0)	53	16.3 (12.0-21.4)	17	5.2 (3.0-8.4)	26	9.5 (6.0-14.1)
1995	424,978	174	55.1 (46.9-64.3)	20	6.9 (4.1-10.8)	147	45.6 (38.2-53.9)	35	10.8 (7.4-15.1)	71	20.1 (15.6-25.5)	10	2.9 (1.4-5.4)	31	11.8 (7.8-16.8)
1996	440,591	159	45.7 (38.7-53.6)	15	4.4 (2.4-7.3)	133	37.7 (31.4-44.9)	42	12.1 (8.6-16.4)	48	13.5 (9.9-18.0)	11	2.9 (1.5-5.3)	32	9.2 (6.2-13.1)
1997	456,202	176	48.1 (41.0-55.9)	17	4.3 (2.5-6.9)	155	42.0 (35.5-49.4)	32	9.2 (6.3-13.1)	65	16.2 (12.5-20.7)	6	1.5 (0.6-3.4)	52	15.1 (11.1-19.8)
1998	471,817	193	51.4 (44.2-59.5)	21	5.4 (3.3-8.3)	166	44.2 (37.5-51.7)	36	10.1 (7.0-14.1)	69	17.7 (13.7-22.5)	11	2.9 (1.4-5.3)	50	13.4 (9.8-17.8)
1999	487,423	210	55.1 (47.7-63.3)	19	5.0 (3.0-7.7)	188	49.2 (42.2-57.0)	29	7.6 (5.0-11.0)	81	21.7 (17.0-27.1)	15	3.8 (2.1-6.3)	63	16.2 (12.3-20.7)
2000	503,040	235	56.9 (49.7-64.8)	19	4.7 (2.8-7.5)	214	51.5 (44.7-59.1)	36	8.2 (5.7-11.3)	86	19.8 (15.8-24.5)	14	3.2 (1.7-5.4)	78	20.4 (16.0-25.5)
2001	516,489	208	48.8 (42.3-56.0)	17	3.7 (2.2-6.0)	178	41.7 (35.7-48.5)	32	7.5 (5.1-10.6)	78	17.7 (13.9-22.1)	6	1.3 (0.5-2.9)	62	15.2 (11.6-19.6)
2002	529,938	229	50.3 (43.9-57.3)	26	6.0 (3.9-8.8)	194	41.7 (36.0-48.1)	36	7.7 (5.4-10.7)	83	17.7 (14.0-22.0)	9	2.0 (0.9-3.7)	66	14.3 (11.0-18.3)
2003	543,387	261	53.0 (46.6-59.9)	17	3.5 (2.1-5.7)	233	46.6 (40.8-53.1)	31	6.2 (4.2-8.8)	95	18.6 (15.0-22.8)	11	2.2 (1.1-4.0)	96	19.6 (15.8-23.9)
2004	556,836	245	47.5 (41.7-53.9)	23	4.4 (2.8-6.7)	213	41.0 (35.6-46.9)	31	6.1 (4.2-8.7)	106	19.8 (16.2-24.0)	10	2.0 (1.0-3.7)	66	13.0 (10.1-16.6)
2005	570,285	265	50.3 (44.4-56.8)	22	4.0 (2.5-6.0)	228	43.0 (37.6-49.0)	44	8.5 (6.2-11.4)	91	16.9 (13.6-20.8)	12	2.4 (1.2-4.2)	81	15.2 (12.1-19.0)
2006	583,734	278	50.1 (44.4-56.4)	27	4.8 (3.2-7.0)	234	41.9 (36.7-47.7)	27	5.1 (3.3-7.4)	118	20.6 (17.0-24.6)	9	1.7 (0.8-3.2)	80	14.5 (11.5-18.1)
2007	597,183	294	50.7 (45.0-56.9)	25	4.3 (2.8-6.4)	248	42.5 (37.4-48.2)	41	7.1 (5.1-9.7)	126	21.1 (17.6-25.2)	6	0.9 (0.3-2.1)	75	13.3 (10.5-16.7)
2008	610,632	250	41.2 (36.2-46.7)	17	2.8 (1.6-4.5)	219	35.8 (31.2-40.9)	34	5.6 (3.9-7.8)	114	18.4 (15.2-22.2)	8	1.3 (0.6-2.6)	63	10.5 (8.0-13.5)
2009	624,081	297	47.0 (41.8-52.7)	24	3.7 (2.4-5.5)	248	39.0 (34.3-44.2)	42	6.6 (4.7-9.0)	127	19.6 (16.3-23.4)	6	1.0 (0.4-2.1)	73	11.8 (9.3-14.9)
2010	637,529	276	41.7 (36.9-47.0)	27	3.9 (2.6-5.7)	238	36.0 (31.5-40.9)	43	6.6 (4.8-8.9)	143	21.6 (18.2-25.5)	6	0.7 (0.3-1.7)	46	7.0 (5.1-9.4)

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (18 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

--Statistic not displayed due to fewer than 5 cases.

Supplemental Table 2. Cancer case counts, population estimates, and incidence rates by histology and year of diagnosis, Chinese females

Year of Diagnosis	Population Estimates	All lung Cancer		Small cell carcinoma		NSCLC		Squamous cell carcinoma		Adenocarcinoma		Large cell carcinoma		NSCLC NOS	
		Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)
1990	357,922	89	32.7 (26.1-40.5)	8	3.1 (1.3-6.0)	79	28.7 (22.5-35.9)	10	3.4 (1.6-6.2)	40	14.1 (9.9-19.3)	15	5.6 (3.1-9.3)	14	5.6 (3.0-9.4)
1991	376,552	89	29.9 (23.9-36.9)	5	1.9 (0.6-4.3)	80	26.5 (20.9-33.1)	10	3.6 (1.7-6.6)	55	17.9 (13.4-23.4)	--	--	11	3.8 (1.9-6.7)
1992	395,188	117	36.5 (30.0-43.8)	8	2.5 (1.1-4.9)	104	31.8 (25.8-38.6)	11	3.3 (1.6-5.9)	63	19.4 (14.8-24.9)	9	2.7 (1.2-5.1)	21	6.4 (4.0-9.9)
1993	413,827	91	26.7 (21.4-32.8)	--	--	86	25.1 (20.0-31.0)	8	2.4 (1.0-4.8)	51	14.5 (10.7-19.1)	12	3.5 (1.8-6.1)	15	4.7 (2.6-7.7)
1994	432,466	110	30.3 (24.8-36.5)	5	1.5 (0.5-3.4)	99	26.8 (21.7-32.7)	12	3.3 (1.7-5.8)	66	17.5 (13.5-22.3)	7	1.9 (0.7-3.9)	14	4.2 (2.3-7.0)
1995	451,104	124	32.5 (27.0-38.8)	8	2.2 (0.9-4.2)	111	29.0 (23.8-35.0)	11	3.1 (1.5-5.4)	78	20.0 (15.8-25.0)	--	--	18	4.9 (2.9-7.8)
1996	469,739	120	29.5 (24.4-35.4)	7	1.7 (0.7-3.4)	107	26.1 (21.4-31.6)	7	1.9 (0.8-3.8)	67	15.9 (12.3-20.2)	9	2.2 (1.0-4.2)	24	6.2 (3.9-9.2)
1997	488,378	135	31.5 (26.4-37.4)	--	--	125	28.8 (23.9-34.4)	9	2.1 (1.0-4.0)	75	17.3 (13.6-21.7)	7	1.4 (0.6-3.0)	34	8.0 (5.5-11.2)
1998	507,017	140	31.1 (26.1-36.7)	7	1.5 (0.6-3.1)	123	27.0 (22.4-32.3)	12	2.8 (1.5-4.9)	71	15.4 (12.0-19.4)	8	1.5 (0.7-3.1)	32	7.3 (5.0-10.3)
1999	525,654	114	23.9 (19.7-28.8)	8	1.9 (0.8-3.6)	101	21.0 (17.1-25.5)	9	2.0 (0.9-3.7)	60	12.2 (9.3-15.8)	8	1.6 (0.7-3.2)	24	5.2 (3.3-7.7)
2000	544,294	139	27.9 (23.4-32.9)	--	--	131	26.1 (21.8-31.0)	7	1.4 (0.5-2.8)	86	17.2 (13.7-21.2)	--	--	34	6.8 (4.7-9.5)
2001	561,345	164	30.8 (26.2-35.9)	6	1.1 (0.4-2.5)	152	28.4 (24.0-33.3)	7	1.3 (0.5-2.7)	94	17.4 (14.0-21.3)	6	1.1 (0.4-2.4)	45	8.5 (6.2-11.4)
2002	578,396	172	30.4 (26.0-35.3)	6	1.0 (0.4-2.3)	158	27.8 (23.6-32.5)	6	1.1 (0.4-2.4)	88	15.3 (12.3-18.9)	8	1.4 (0.6-2.7)	56	10.0 (7.6-13.0)
2003	595,447	183	30.6 (26.3-35.4)	--	--	170	28.4 (24.3-33.0)	9	1.5 (0.7-2.9)	115	19.1 (15.7-22.9)	8	1.4 (0.6-2.7)	38	6.5 (4.6-8.9)
2004	612,498	177	27.8 (23.9-32.3)	--	--	165	25.9 (22.1-30.2)	11	1.7 (0.9-3.1)	104	16.2 (13.2-19.7)	7	1.1 (0.5-2.4)	43	6.8 (4.9-9.2)
2005	629,549	191	28.9 (25.0-33.4)	8	1.2 (0.5-2.4)	173	26.2 (22.4-30.4)	13	1.9 (1.0-3.3)	110	16.5 (13.6-19.9)	9	1.5 (0.7-2.8)	41	6.2 (4.5-8.5)
2006	646,600	214	31.2 (27.1-35.7)	--	--	200	29.1 (25.2-33.4)	7	1.0 (0.4-2.0)	130	18.7 (15.6-22.3)	7	1.1 (0.4-2.3)	56	8.3 (6.3-10.8)
2007	663,651	185	25.4 (21.9-29.4)	8	1.1 (0.5-2.2)	163	22.4 (19.1-26.2)	9	1.2 (0.6-2.3)	96	13.3 (10.7-16.3)	6	0.8 (0.3-1.8)	52	7.1 (5.3-9.4)
2008	680,702	179	23.8 (20.4-27.6)	9	1.3 (0.6-2.4)	162	21.4 (18.2-25.0)	8	1.1 (0.5-2.2)	119	15.7 (12.9-18.8)	--	--	31	4.1 (2.8-5.9)
2009	697,753	220	27.5 (24.0-31.5)	--	--	197	24.6 (21.3-28.4)	12	1.6 (0.8-2.8)	139	17.3 (14.5-20.4)	5	0.7 (0.2-1.6)	41	5.1 (3.7-7.0)
2010	714,800	219	26.7 (23.3-30.6)	7	0.9 (0.4-2.0)	201	24.5 (21.2-28.2)	14	1.7 (0.9-2.9)	148	18.0 (15.2-21.2)	--	--	36	4.4 (3.0-6.1)

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (18 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

--Statistic not displayed due to fewer than 5 cases.

Supplemental Table 3. Cancer case counts, population estimates, and incidence rates by histology and year of diagnosis, foreign-born Chinese males

Year of Diagnosis	Population Estimates	All lung Cancer		Small cell carcinoma		NSCLC		Squamous cell carcinoma		Adenocarcinoma		Large cell carcinoma		NSCLC NOS	
		Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)
1990	237,869	120	59.6 (48.6-73.2)	8	3.7 (1.6-9.4)	110	54.2 (43.7-67.2)	28	15.0 (9.5-23.4)	50	23.0 (16.7-31.9)	11	5.9 (2.6-12.7)	21	10.4 (6.2-17.6)
1991	249,430	145	66.2 (55.2-79.3)	14	6.6 (3.4-12.8)	130	59.1 (48.8-71.6)	42	19.0 (13.4-27.2)	48	19.9 (14.5-27.7)	16	7.6 (4.2-13.9)	24	12.6 (7.7-20.3)
1992	261,508	174	73.7 (62.6-87.0)	14	5.3 (2.8-10.7)	155	65.2 (54.8-77.7)	41	17.1 (12.0-24.6)	63	25.8 (19.6-34.2)	17	7.2 (4.0-13.2)	34	15.1 (10.1-22.7)
1993	271,779	159	63.2 (53.1-75.2)	20	8.0 (4.7-13.9)	134	52.1 (43.2-63.1)	35	14.3 (9.7-21.5)	48	16.9 (12.4-23.7)	11	4.5 (2.1-9.9)	40	16.4 (11.4-23.8)
1994	279,887	138	54.6 (45.3-66.0)	14	5.3 (2.8-10.7)	115	43.6 (35.5-53.7)	31	11.4 (7.5-17.7)	42	15.7 (11.2-22.7)	17	5.7 (3.3-10.9)	25	10.8 (6.7-17.5)
1995	288,079	157	57.5 (48.5-68.5)	19	7.0 (4.1-12.5)	130	46.8 (38.8-56.8)	32	11.5 (7.7-17.6)	62	20.2 (15.4-27.1)	7	2.6 (1.0-7.4)	29	12.5 (8.1-19.3)
1996	296,143	141	46.3 (38.8-55.8)	13	4.4 (2.3-9.4)	116	37.6 (30.8-46.3)	39	12.9 (9.0-19.1)	40	12.6 (9.0-18.6)	9	2.6 (1.2-7.2)	28	9.4 (6.1-15.1)
1997	306,316	163	50.1 (42.5-59.5)	15	4.3 (2.4-8.9)	144	43.8 (36.7-52.7)	33	10.4 (7.1-16.0)	60	16.9 (12.8-23.0)	5	1.4 (0.5-5.9)	46	15.1 (10.9-21.5)
1998	316,488	178	53.4 (45.6-63.1)	20	5.9 (3.6-10.7)	153	45.8 (38.6-54.8)	33	10.7 (7.2-16.4)	64	18.3 (14.0-24.7)	9	2.8 (1.2-7.4)	47	14.0 (10.2-20.0)
1999	327,459	176	51.8 (44.2-61.1)	16	4.7 (2.7-9.5)	157	46.0 (38.9-54.9)	25	7.4 (4.7-12.6)	66	20.0 (15.3-26.7)	16	4.4 (2.5-9.1)	50	14.2 (10.4-20.0)
2000	339,388	198	52.5 (45.3-61.3)	14	3.4 (1.9-7.8)	182	48.4 (41.5-57.0)	35	9.0 (6.2-14.0)	67	17.3 (13.3-23.2)	13	3.4 (1.8-7.8)	67	18.8 (14.5-25.1)
2001	345,876	195	51.1 (44.1-59.9)	19	4.7 (2.8-9.2)	166	43.6 (37.1-51.9)	33	8.6 (5.9-13.6)	66	16.9 (13.0-22.8)	6	1.5 (0.5-5.7)	61	16.6 (12.6-22.6)
2002	350,687	211	51.9 (45.0-60.5)	23	5.7 (3.6-10.4)	180	43.7 (37.5-51.7)	31	7.4 (5.0-12.3)	82	19.8 (15.7-25.9)	8	2.1 (0.9-6.5)	59	14.4 (10.9-20.1)
2003	354,392	237	54.8 (48.0-63.4)	18	4.1 (2.4-8.6)	210	48.1 (41.8-56.2)	29	6.6 (4.4-11.3)	85	19.2 (15.3-25.1)	10	2.3 (1.1-6.6)	86	20.1 (16.0-26.2)
2004	356,851	230	50.5 (44.1-58.6)	23	5.0 (3.1-9.6)	197	42.9 (37.1-50.5)	33	7.5 (5.1-12.4)	96	20.2 (16.3-26.1)	9	2.1 (0.9-6.5)	59	13.2 (10.0-18.6)

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (18 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

--Statistic not displayed due to fewer than 5 cases.

Supplemental Table 4. Cancer case counts, population estimates, and incidence rates by histology and year of diagnosis, US-born Chinese males

Year of Diagnosis	Population Estimates	All lung Cancer		Small cell carcinoma		NSCLC		Squamous cell carcinoma		Adenocarcinoma		Large cell carcinoma		NSCLC NOS	
		Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)
1990-1992	347,433	41	41.8 (28.6-58.4)	--	--	37	38.0 (25.3-54.2)	6	5.6 (1.8-12.7)	18	18.4 (9.6-31.1)	--	--	11	12.6 (6.0-22.6)
1993-1995	405,173	65	51.7 (38.4-67.6)	9	10.0 (3.8-19.9)	56	43.4 (31.5-57.8)	8	5.6 (2.2-11.3)	30	21.4 (14.1-30.9)	--	--	14	13.8 (6.5-24.5)
1996-1998	450,333	60	46.9 (35.0-61.1)	5	3.3 (1.0-7.6)	53	42.7 (31.2-56.5)	12	9.1 (4.4-16.3)	22	18.2 (10.9-27.8)	--	--	15	12.2 (6.5-20.3)
1999-2001	498,475	107	75.7 (61.4-92.1)	8	7.1 (2.9-13.7)	99	68.6 (55.2-84.0)	11	7.1 (3.5-12.4)	54	35.0 (26.0-45.8)	--	--	32	25.5 (17.0-36.2)
2002-2004	577,303	87	50.0 (39.8-61.7)	--	--	80	45.1 (35.5-56.2)	9	4.8 (2.1-8.9)	38	21.0 (14.7-28.8)	--	--	29	16.9 (11.2-24.1)

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (18 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

--Statistic not displayed due to fewer than 5 cases.

Supplemental Table 5. Cancer case counts, population estimates, and incidence rates by histology and year of diagnosis, foreign-born Chinese females

Year of Diagnosis	Population Estimates	All lung Cancer		Small cell carcinoma		NSCLC		Squamous cell carcinoma		Adenocarcinoma		Large cell carcinoma		NSCLC NOS	
		Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)
1990	255,888	80	33.6 (26.4-43.3)	5	2.3 (0.7-8.4)	73	30.1 (23.4-39.4)	9	3.4 (1.5-9.4)	37	14.9 (10.4-22.5)	14	6.0 (3.2-12.4)	13	5.8 (3.0-12.2)
1991	269,107	72	28.2 (21.9-36.9)	--	--	64	24.6 (18.8-32.8)	8	3.4 (1.4-9.0)	42	15.8 (11.3-23.0)	--	--	11	4.4 (2.1-10.0)
1992	283,791	108	39.5 (32.2-48.9)	7	2.5 (1.0-7.6)	97	34.9 (28.1-43.8)	10	3.4 (1.6-8.6)	59	21.6 (16.3-29.1)	11	3.8 (1.9-9.0)	17	6.1 (3.5-11.7)
1993	297,704	77	26.9 (21.1-34.8)	--	--	72	25.0 (19.5-32.7)	8	2.8 (1.2-7.6)	43	14.5 (10.4-20.9)	8	2.8 (1.2-7.7)	13	4.9 (2.6-10.1)
1994	309,588	91	29.7 (23.8-37.5)	--	--	82	26.2 (20.8-33.7)	11	3.7 (1.8-8.4)	56	17.4 (13.1-23.8)	5	1.6 (0.5-6.0)	10	3.5 (1.7-8.3)
1995	321,492	101	31.6 (25.7-39.3)	6	1.9 (0.7-6.1)	90	28.2 (22.6-35.5)	10	3.3 (1.6-7.7)	64	19.5 (15.0-25.9)	--	--	13	4.4 (2.3-9.0)
1996	333,166	113	32.6 (26.8-40.1)	6	1.7 (0.6-5.6)	101	28.9 (23.5-36.0)	7	2.2 (0.9-6.2)	61	16.7 (12.7-22.5)	8	2.4 (1.0-6.4)	25	7.6 (4.9-12.4)
1997	347,525	127	34.8 (28.9-42.2)	--	--	115	31.0 (25.5-38.0)	8	2.2 (0.9-5.9)	66	17.9 (13.8-23.7)	7	1.6 (0.6-5.2)	34	9.3 (6.4-14.1)
1998	361,232	127	32.1 (26.7-39.0)	6	1.5 (0.5-4.9)	112	28.0 (23.0-34.5)	11	3.0 (1.5-6.7)	66	16.2 (12.5-21.5)	8	1.8 (0.8-5.2)	27	7.0 (4.6-11.3)
1999	374,159	102	24.6 (20.0-30.6)	8	2.1 (0.9-5.6)	90	21.4 (17.2-27.1)	5	1.3 (0.4-4.5)	52	12.1 (9.0-16.8)	7	1.6 (0.6-4.8)	26	6.4 (4.2-10.5)
2000	388,009	131	29.4 (24.5-35.5)	--	--	125	27.9 (23.2-33.9)	7	1.5 (0.6-4.5)	81	18.1 (14.3-23.2)	--	--	34	7.7 (5.3-11.6)
2001	396,185	146	31.0 (26.2-37.1)	5	1.1 (0.4-4.1)	136	28.7 (24.0-34.6)	8	1.7 (0.7-4.7)	86	18.0 (14.4-23.0)	5	1.0 (0.3-3.9)	37	7.9 (5.6-11.8)
2002	403,279	158	31.7 (26.9-37.7)	5	1.0 (0.3-4.0)	143	28.6 (24.1-34.4)	6	1.2 (0.5-4.3)	76	15.0 (11.8-19.6)	9	1.8 (0.8-4.9)	52	10.6 (7.9-14.8)
2003	408,853	173	32.9 (28.2-39.0)	--	--	163	31.0 (26.4-36.8)	6	1.1 (0.4-4.2)	110	20.8 (17.1-25.8)	8	1.6 (0.7-4.6)	39	7.5 (5.3-11.3)
2004	413,064	168	30.2 (25.8-35.9)	--	--	154	27.6 (23.4-33.1)	7	1.3 (0.5-4.4)	102	18.1 (14.7-22.9)	6	1.2 (0.4-4.3)	39	7.0 (5.0-10.8)

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (18 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

--Statistic not displayed due to fewer than 5 cases.

Supplemental Table 6. Cancer case counts, population estimates, and incidence rates by histology and year of diagnosis, US-born Chinese females

Year of Diagnosis	Population Estimates	All lung Cancer		Small cell carcinoma		NSCLC		Squamous cell carcinoma		Adenocarcinoma		Large cell carcinoma		NSCLC NOS	
		Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)	Case Count	Incidence rate (95% CI)
1990-1992	325,702	42	37.8 (26.3-52.2)	6	5.1 (1.8-11.1)	35	31.1 (20.7-44.4)	--	--	23	19.4 (11.6-30.0)	--	--	5	5.8 (1.5-13.8)
1993-1995	384,147	60	41.6 (31.3-54.0)	--	--	55	37.5 (27.8-49.3)	--	--	35	23.0 (15.7-32.3)	7	4.8 (1.8-9.9)	11	8.4 (4.0-15.0)
1996-1998			24.4 (17.2-33.2)	--	--		37	22.5 (15.6-31.0)	--		--				23
1999-2001	474,487	62	31.5 (24.1-40.3)	--	--	57	28.8 (21.7-37.3)	7	3.4 (1.4-6.9)	31	15.5 (10.5-21.9)	--	--	15	7.7 (4.3-12.6)
2002-2004	549,017	62	27.6 (21.0-35.5)	--	--	59	26.0 (19.6-33.5)	8	3.7 (1.6-7.1)	32	14.0 (9.5-19.8)	--	--	18	7.8 (4.5-12.3)

Rates are per 100,000 and age-adjusted to the 2000 US Std Population (18 age groups - Census P25-1130) standard; Confidence intervals (Tiwari mod) are 95% for rates.

--Statistic not displayed due to fewer than 5 cases.