

Supplementary Materials: Reference Intervals for Non-Fasting CVD Lipids and Inflammation Markers in Pregnant Indigenous Australian Women

Tracy L. Schumacher, Christopher Oldmeadow, Don Clausen, Loretta Weatherall, Lyniece Keogh, Kirsty G. Pringle and Kym M. Rae

Table S 1: Distributions, fractional polynomial terms and goodness of fit values used in creating of reference ranges.

| CVD factor | Distribution | Fractional polynomial terms | | | Goodness of fit (p) | |
|-------------------|--------------|-----------------------------|---|----|---------------------|----------------------------|
| | | M | S | G | | |
| Cholesterol | men | -0.5 | 3 | -1 | 0 | 0.362 |
| Triglycerides | men | -2 | 2 | 2 | 0 | 0.366 |
| Cystatin-C | men | 2 | 2 | 3 | 0 | 0.358 |
| ALP (16-22 years) | en | 3 | 3 | 3 | 0 | 0.583 |
| ALP (22+ years) | en | 2 | 3 | 3 | 0 | Not available ¹ |

¹ Insufficient degrees of freedom for test of means. Results for the ranges provided for ALP 16-22 years should be treated with caution.

Table S 2. Non-fasting total cholesterol ¹percentiles for Indigenous Australian women during pregnancy

| Gestation week | 2.5 th | 5 th | 10 th | 25 th | 50 th | 75 th | 90 th | 95 th | 97.5 th |
|----------------|-------------------|-----------------|------------------|------------------|------------------|------------------|------------------|------------------|--------------------|
| 12 | 3.60 | 3.72 | 3.87 | 4.19 | 4.68 | 5.26 | 5.78 | 6.09 | 6.38 |
| 13 | 3.63 | 3.76 | 3.93 | 4.27 | 4.81 | 5.45 | 6.00 | 6.34 | 6.64 |
| 14 | 3.65 | 3.80 | 3.98 | 4.35 | 4.93 | 5.61 | 6.19 | 6.55 | 6.86 |
| 15 | 3.68 | 3.84 | 4.04 | 4.43 | 5.04 | 5.76 | 6.37 | 6.73 | 7.06 |
| 16 | 3.71 | 3.88 | 4.09 | 4.50 | 5.14 | 5.89 | 6.52 | 6.90 | 7.23 |
| 17 | 3.74 | 3.92 | 4.13 | 4.57 | 5.24 | 6.01 | 6.66 | 7.05 | 7.39 |
| 18 | 3.77 | 3.95 | 4.18 | 4.63 | 5.33 | 6.13 | 6.79 | 7.19 | 7.53 |
| 19 | 3.81 | 3.99 | 4.23 | 4.70 | 5.41 | 6.23 | 6.91 | 7.31 | 7.66 |
| 20 | 3.84 | 4.03 | 4.27 | 4.76 | 5.49 | 6.33 | 7.02 | 7.43 | 7.78 |
| 21 | 3.87 | 4.07 | 4.32 | 4.82 | 5.57 | 6.43 | 7.13 | 7.54 | 7.90 |
| 22 | 3.90 | 4.11 | 4.37 | 4.87 | 5.65 | 6.52 | 7.23 | 7.64 | 8.00 |
| 23 | 3.94 | 4.15 | 4.41 | 4.93 | 5.72 | 6.60 | 7.32 | 7.74 | 8.10 |
| 24 | 3.97 | 4.19 | 4.46 | 4.99 | 5.79 | 6.69 | 7.41 | 7.83 | 8.20 |
| 25 | 4.01 | 4.23 | 4.50 | 5.05 | 5.86 | 6.77 | 7.50 | 7.92 | 8.29 |
| 26 | 4.05 | 4.27 | 4.55 | 5.10 | 5.93 | 6.85 | 7.59 | 8.01 | 8.38 |
| 27 | 4.09 | 4.31 | 4.60 | 5.16 | 6.00 | 6.93 | 7.67 | 8.10 | 8.46 |
| 28 | 4.13 | 4.36 | 4.65 | 5.22 | 6.07 | 7.01 | 7.75 | 8.18 | 8.55 |
| 29 | 4.17 | 4.41 | 4.70 | 5.28 | 6.14 | 7.08 | 7.83 | 8.26 | 8.63 |
| 30 | 4.22 | 4.45 | 4.76 | 5.34 | 6.21 | 7.16 | 7.91 | 8.34 | 8.71 |
| 31 | 4.26 | 4.50 | 4.81 | 5.40 | 6.28 | 7.24 | 8.00 | 8.43 | 8.79 |
| 32 | 4.31 | 4.56 | 4.87 | 5.47 | 6.35 | 7.32 | 8.08 | 8.51 | 8.87 |
| 33 | 4.36 | 4.61 | 4.92 | 5.53 | 6.42 | 7.39 | 8.16 | 8.59 | 8.95 |
| 34 | 4.42 | 4.67 | 4.98 | 5.60 | 6.50 | 7.47 | 8.24 | 8.67 | 9.04 |
| 35 | 4.47 | 4.73 | 5.05 | 5.66 | 6.57 | 7.55 | 8.32 | 8.75 | 9.12 |
| 36 | 4.53 | 4.79 | 5.11 | 5.73 | 6.65 | 7.64 | 8.4 | 8.84 | 9.20 |
| 37 | 4.59 | 4.85 | 5.18 | 5.81 | 6.73 | 7.72 | 8.49 | 8.92 | 9.29 |
| 38 | 4.65 | 4.92 | 5.25 | 5.88 | 6.81 | 7.80 | 8.58 | 9.01 | 9.37 |
| 39 | 4.72 | 4.98 | 5.32 | 5.96 | 6.89 | 7.89 | 8.66 | 9.10 | 9.46 |
| 40 | 4.79 | 5.06 | 5.39 | 6.04 | 6.98 | 7.98 | 8.75 | 9.19 | 9.55 |

¹reference range for non-pregnant women: <5.5mmol/L [21]

Table S 3. Non-fasting triglycerides¹ percentiles for Indigenous Australian women during pregnancy

| Gestation week | 2.5 th | 5 th | 10 th | 25 th | 50 th | 75 th | 90 th | 95 th | 97.5 th |
|----------------|-------------------|-----------------|------------------|------------------|------------------|------------------|------------------|------------------|--------------------|
| 12 | 0.75 | 0.84 | 0.95 | 1.15 | 1.40 | 1.70 | 2.05 | 2.32 | 2.61 |
| 13 | 0.77 | 0.86 | 0.97 | 1.18 | 1.43 | 1.74 | 2.10 | 2.38 | 2.68 |
| 14 | 0.79 | 0.89 | 1.00 | 1.21 | 1.47 | 1.78 | 2.16 | 2.45 | 2.76 |
| 15 | 0.82 | 0.91 | 1.03 | 1.24 | 1.51 | 1.83 | 2.22 | 2.52 | 2.85 |
| 16 | 0.84 | 0.94 | 1.06 | 1.28 | 1.55 | 1.89 | 2.29 | 2.60 | 2.94 |
| 17 | 0.87 | 0.97 | 1.09 | 1.31 | 1.59 | 1.94 | 2.36 | 2.68 | 3.03 |
| 18 | 0.89 | 1.00 | 1.12 | 1.35 | 1.64 | 2.00 | 2.43 | 2.76 | 3.14 |
| 19 | 0.92 | 1.03 | 1.16 | 1.39 | 1.69 | 2.06 | 2.50 | 2.85 | 3.24 |
| 20 | 0.95 | 1.06 | 1.19 | 1.44 | 1.74 | 2.12 | 2.58 | 2.95 | 3.35 |
| 21 | 0.98 | 1.10 | 1.23 | 1.48 | 1.80 | 2.19 | 2.67 | 3.05 | 3.47 |
| 22 | 1.02 | 1.13 | 1.27 | 1.53 | 1.86 | 2.26 | 2.76 | 3.15 | 3.59 |
| 23 | 1.05 | 1.17 | 1.31 | 1.58 | 1.92 | 2.34 | 2.85 | 3.26 | 3.72 |
| 24 | 1.09 | 1.21 | 1.36 | 1.63 | 1.98 | 2.41 | 2.95 | 3.37 | 3.85 |
| 25 | 1.12 | 1.25 | 1.40 | 1.69 | 2.05 | 2.49 | 3.05 | 3.49 | 3.99 |
| 26 | 1.16 | 1.29 | 1.45 | 1.74 | 2.11 | 2.58 | 3.15 | 3.61 | 4.14 |
| 27 | 1.20 | 1.34 | 1.50 | 1.80 | 2.19 | 2.67 | 3.26 | 3.74 | 4.29 |
| 28 | 1.24 | 1.38 | 1.55 | 1.86 | 2.26 | 2.76 | 3.37 | 3.87 | 4.44 |
| 29 | 1.29 | 1.43 | 1.60 | 1.92 | 2.34 | 2.85 | 3.49 | 4.01 | 4.61 |
| 30 | 1.33 | 1.48 | 1.66 | 1.99 | 2.41 | 2.95 | 3.61 | 4.15 | 4.77 |
| 31 | 1.38 | 1.53 | 1.71 | 2.06 | 2.5 | 3.05 | 3.73 | 4.29 | 4.95 |
| 32 | 1.42 | 1.58 | 1.77 | 2.13 | 2.58 | 3.15 | 3.86 | 4.45 | 5.13 |
| 33 | 1.47 | 1.63 | 1.83 | 2.20 | 2.67 | 3.26 | 4.00 | 4.60 | 5.31 |
| 34 | 1.52 | 1.69 | 1.89 | 2.27 | 2.76 | 3.37 | 4.13 | 4.76 | 5.50 |
| 35 | 1.58 | 1.75 | 1.96 | 2.35 | 2.85 | 3.48 | 4.27 | 4.93 | 5.70 |
| 36 | 1.63 | 1.81 | 2.02 | 2.43 | 2.94 | 3.60 | 4.42 | 5.10 | 5.90 |
| 37 | 1.68 | 1.87 | 2.09 | 2.51 | 3.04 | 3.72 | 4.57 | 5.27 | 6.11 |
| 38 | 1.74 | 1.93 | 2.16 | 2.59 | 3.14 | 3.84 | 4.72 | 5.45 | 6.33 |
| 39 | 1.80 | 1.99 | 2.23 | 2.67 | 3.24 | 3.97 | 4.88 | 5.64 | 6.55 |
| 40 | 1.86 | 2.06 | 2.30 | 2.76 | 3.35 | 4.10 | 5.04 | 5.83 | 6.78 |

¹reference range for non-pregnant women: <2.00mmol/L [21]

Table S 4. Non-fasting Cystatin C¹ percentiles for Indigenous Australian women during pregnancy

| Gestation week | 2.5th | 5th | 10th | 25th | 50th | 75th | 90th | 95th | 97.5th |
|-----------------------|-------------------------|-----------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|--------------------------|
| 12 | 0.53 | 0.56 | 0.60 | 0.65 | 0.72 | 0.78 | 0.86 | 0.91 | 0.97 |
| 13 | 0.53 | 0.56 | 0.59 | 0.65 | 0.71 | 0.78 | 0.86 | 0.91 | 0.97 |
| 14 | 0.53 | 0.56 | 0.59 | 0.65 | 0.71 | 0.78 | 0.86 | 0.91 | 0.97 |
| 15 | 0.52 | 0.55 | 0.59 | 0.65 | 0.71 | 0.78 | 0.86 | 0.91 | 0.97 |
| 16 | 0.52 | 0.55 | 0.59 | 0.64 | 0.71 | 0.78 | 0.86 | 0.92 | 0.98 |
| 17 | 0.52 | 0.55 | 0.59 | 0.64 | 0.71 | 0.78 | 0.86 | 0.92 | 0.98 |
| 18 | 0.52 | 0.55 | 0.59 | 0.65 | 0.71 | 0.78 | 0.87 | 0.93 | 0.99 |
| 19 | 0.52 | 0.55 | 0.59 | 0.65 | 0.71 | 0.79 | 0.87 | 0.94 | 1.00 |
| 20 | 0.52 | 0.56 | 0.59 | 0.65 | 0.72 | 0.79 | 0.88 | 0.95 | 1.02 |
| 21 | 0.53 | 0.56 | 0.60 | 0.66 | 0.73 | 0.80 | 0.89 | 0.96 | 1.03 |
| 22 | 0.53 | 0.56 | 0.60 | 0.66 | 0.73 | 0.81 | 0.91 | 0.98 | 1.05 |
| 23 | 0.54 | 0.57 | 0.61 | 0.67 | 0.74 | 0.82 | 0.92 | 0.99 | 1.07 |
| 24 | 0.55 | 0.58 | 0.62 | 0.68 | 0.75 | 0.84 | 0.94 | 1.01 | 1.09 |
| 25 | 0.55 | 0.59 | 0.63 | 0.69 | 0.77 | 0.85 | 0.95 | 1.03 | 1.11 |
| 26 | 0.56 | 0.60 | 0.64 | 0.70 | 0.78 | 0.87 | 0.97 | 1.05 | 1.14 |
| 27 | 0.57 | 0.61 | 0.65 | 0.72 | 0.80 | 0.89 | 1.00 | 1.08 | 1.17 |
| 28 | 0.59 | 0.62 | 0.66 | 0.73 | 0.81 | 0.91 | 1.02 | 1.11 | 1.20 |
| 29 | 0.60 | 0.64 | 0.68 | 0.75 | 0.83 | 0.93 | 1.05 | 1.14 | 1.24 |
| 30 | 0.62 | 0.65 | 0.70 | 0.77 | 0.86 | 0.95 | 1.08 | 1.17 | 1.28 |
| 31 | 0.63 | 0.67 | 0.71 | 0.79 | 0.88 | 0.98 | 1.11 | 1.21 | 1.32 |
| 32 | 0.65 | 0.69 | 0.73 | 0.81 | 0.90 | 1.01 | 1.14 | 1.25 | 1.37 |
| 33 | 0.67 | 0.71 | 0.76 | 0.84 | 0.93 | 1.04 | 1.18 | 1.29 | 1.42 |
| 34 | 0.69 | 0.73 | 0.78 | 0.86 | 0.96 | 1.08 | 1.22 | 1.34 | 1.47 |
| 35 | 0.71 | 0.76 | 0.81 | 0.89 | 0.99 | 1.11 | 1.26 | 1.39 | 1.53 |
| 36 | 0.74 | 0.78 | 0.83 | 0.92 | 1.03 | 1.15 | 1.31 | 1.44 | 1.59 |
| 37 | 0.76 | 0.81 | 0.86 | 0.96 | 1.06 | 1.19 | 1.36 | 1.49 | 1.65 |
| 38 | 0.79 | 0.84 | 0.89 | 0.99 | 1.10 | 1.23 | 1.41 | 1.55 | 1.72 |
| 39 | 0.82 | 0.87 | 0.93 | 1.02 | 1.14 | 1.28 | 1.46 | 1.62 | 1.79 |
| 40 | 0.85 | 0.90 | 0.96 | 1.06 | 1.18 | 1.33 | 1.52 | 1.68 | 1.87 |

¹reference range for non-pregnant women: <2.00mmol/L [21]

Table S 5. Non-fasting ALP¹ percentiles for Indigenous Australian women during pregnancy aged 16 to less than 22 years

| Gestation week | 2.5 th | 5 th | 10 th | 25 th | 50 th | 75 th | 90 th | 95 th | 97.5 th |
|----------------|-------------------|-----------------|------------------|------------------|------------------|------------------|------------------|------------------|--------------------|
| 12 | 42.40 | 44.61 | 47.37 | 52.56 | 59.49 | 68.28 | 78.88 | 87.36 | 97.06 |
| 13 | 42.15 | 44.44 | 47.27 | 52.62 | 59.75 | 68.79 | 79.68 | 88.39 | 98.32 |
| 14 | 42.00 | 44.36 | 47.29 | 52.82 | 60.18 | 69.51 | 80.74 | 89.72 | 99.94 |
| 15 | 41.96 | 44.40 | 47.44 | 53.17 | 60.80 | 70.47 | 82.10 | 91.39 | 101.95 |
| 16 | 42.03 | 44.58 | 47.74 | 53.70 | 61.64 | 71.69 | 83.78 | 93.42 | 104.38 |
| 17 | 42.25 | 44.91 | 48.21 | 54.43 | 62.72 | 73.21 | 85.81 | 95.86 | 107.26 |
| 18 | 42.61 | 45.40 | 48.86 | 55.38 | 64.06 | 75.04 | 88.23 | 98.73 | 110.64 |
| 19 | 43.15 | 46.08 | 49.72 | 56.57 | 65.69 | 77.22 | 91.06 | 102.07 | 114.55 |
| 20 | 43.87 | 46.96 | 50.80 | 58.01 | 67.62 | 79.76 | 94.32 | 105.91 | 119.02 |
| 21 | 44.80 | 48.06 | 52.11 | 59.73 | 69.88 | 82.69 | 98.06 | 110.28 | 124.09 |
| 22 | 45.94 | 49.40 | 53.68 | 61.75 | 72.48 | 86.04 | 102.3 | 115.21 | 129.79 |
| 23 | 47.32 | 50.98 | 55.53 | 64.08 | 75.47 | 89.84 | 107.06 | 120.73 | 136.17 |
| 24 | 48.95 | 52.84 | 57.67 | 66.76 | 78.85 | 94.11 | 112.38 | 126.88 | 143.24 |
| 25 | 50.84 | 54.98 | 60.12 | 69.78 | 82.64 | 98.87 | 118.29 | 133.69 | 151.06 |
| 26 | 53.02 | 57.42 | 62.90 | 73.19 | 86.88 | 104.15 | 124.82 | 141.20 | 159.65 |
| 27 | 55.49 | 60.19 | 66.02 | 76.99 | 91.58 | 109.98 | 131.99 | 149.42 | 169.05 |
| 28 | 58.28 | 63.29 | 69.51 | 81.21 | 96.77 | 116.38 | 139.84 | 158.4 | 179.30 |
| 29 | 61.39 | 66.74 | 73.38 | 85.86 | 102.46 | 123.38 | 148.39 | 168.17 | 190.43 |
| 30 | 64.86 | 70.56 | 77.65 | 90.97 | 108.68 | 131.00 | 157.67 | 178.76 | 202.48 |
| 31 | 68.68 | 74.77 | 82.34 | 96.56 | 115.46 | 139.27 | 167.72 | 190.21 | 215.48 |
| 32 | 72.88 | 79.39 | 87.46 | 102.64 | 122.81 | 148.22 | 178.56 | 202.53 | 229.46 |
| 33 | 77.48 | 84.42 | 93.04 | 109.23 | 130.76 | 157.87 | 190.22 | 215.78 | 244.47 |
| 34 | 82.49 | 89.89 | 99.09 | 116.37 | 139.32 | 168.24 | 202.73 | 229.97 | 260.54 |
| 35 | 87.92 | 95.82 | 105.62 | 124.05 | 148.53 | 179.36 | 216.13 | 245.15 | 277.70 |
| 36 | 93.79 | 102.22 | 112.67 | 132.31 | 158.41 | 191.26 | 230.43 | 261.34 | 295.98 |
| 37 | 100.13 | 109.10 | 120.24 | 141.17 | 168.97 | 203.96 | 245.67 | 278.57 | 315.43 |
| 38 | 106.93 | 116.49 | 128.35 | 150.64 | 180.24 | 217.49 | 261.88 | 296.88 | 336.09 |
| 39 | 114.23 | 124.40 | 137.02 | 160.74 | 192.24 | 231.87 | 279.09 | 316.31 | 357.97 |
| 40 | 122.03 | 132.85 | 146.28 | 171.50 | 205.00 | 247.13 | 297.32 | 336.87 | 381.13 |

¹reference range for non-pregnant women: <35-140 U/L (16<22 yrs) [21]

Table S 6. Non-fasting ALP¹ percentiles for Indigenous Australian women during pregnancy aged 22 years and older

| Gestation week | 2.5 th | 5 th | 10 th | 25 th | 50 th | 75 th | 90 th | 95 th | 97.5 th |
|----------------|-------------------|-----------------|------------------|------------------|------------------|------------------|------------------|------------------|--------------------|
| 12 | 30.64 | 34.92 | 39.89 | 48.29 | 57.76 | 67.38 | 76.16 | 81.48 | 86.14 |
| 13 | 30.86 | 35.10 | 40.05 | 48.47 | 58.07 | 67.92 | 77.03 | 82.59 | 87.48 |
| 14 | 31.07 | 35.29 | 40.24 | 48.73 | 58.48 | 68.61 | 78.06 | 83.88 | 89.04 |
| 15 | 31.30 | 35.53 | 40.50 | 49.08 | 59.03 | 69.46 | 79.29 | 85.39 | 90.83 |
| 16 | 31.57 | 35.82 | 40.84 | 49.55 | 59.73 | 70.49 | 80.74 | 87.15 | 92.89 |
| 17 | 31.89 | 36.18 | 41.28 | 50.15 | 60.59 | 71.73 | 82.43 | 89.18 | 95.26 |
| 18 | 32.28 | 36.64 | 41.83 | 50.90 | 61.64 | 73.20 | 84.40 | 91.51 | 97.96 |
| 19 | 32.77 | 37.21 | 42.51 | 51.81 | 62.90 | 74.92 | 86.66 | 94.17 | 101.03 |
| 20 | 33.37 | 37.91 | 43.33 | 52.90 | 64.38 | 76.91 | 89.25 | 97.20 | 104.49 |
| 21 | 34.10 | 38.76 | 44.33 | 54.21 | 66.11 | 79.20 | 92.19 | 100.63 | 108.40 |
| 22 | 34.98 | 39.77 | 45.51 | 55.73 | 68.12 | 81.81 | 95.52 | 104.47 | 112.78 |
| 23 | 36.02 | 40.96 | 46.90 | 57.50 | 70.41 | 84.78 | 99.26 | 108.78 | 117.66 |
| 24 | 37.25 | 42.36 | 48.52 | 59.53 | 73.01 | 88.11 | 103.43 | 113.58 | 123.09 |
| 25 | 38.68 | 43.98 | 50.37 | 61.85 | 75.96 | 91.84 | 108.08 | 118.9 | 129.11 |
| 26 | 40.33 | 45.84 | 52.5 | 64.47 | 79.26 | 96.00 | 113.23 | 124.78 | 135.74 |
| 27 | 42.23 | 47.96 | 54.90 | 67.42 | 82.94 | 100.60 | 118.90 | 131.26 | 143.03 |
| 28 | 44.39 | 50.36 | 57.62 | 70.72 | 87.03 | 105.68 | 125.14 | 138.36 | 151.02 |
| 29 | 46.83 | 53.07 | 60.65 | 74.39 | 91.54 | 111.27 | 131.97 | 146.12 | 159.75 |
| 30 | 49.58 | 56.10 | 64.04 | 78.45 | 96.51 | 117.39 | 139.42 | 154.58 | 169.27 |
| 31 | 52.65 | 59.48 | 67.80 | 82.93 | 101.96 | 124.06 | 147.53 | 163.77 | 179.60 |
| 32 | 56.06 | 63.21 | 71.94 | 87.85 | 107.92 | 131.32 | 156.33 | 173.73 | 190.79 |
| 33 | 59.84 | 67.34 | 76.50 | 93.23 | 114.40 | 139.20 | 165.85 | 184.51 | 202.90 |
| 34 | 64.01 | 71.87 | 81.49 | 99.09 | 121.43 | 147.72 | 176.12 | 196.12 | 215.95 |
| 35 | 68.58 | 76.84 | 86.94 | 105.46 | 129.05 | 156.90 | 187.18 | 208.62 | 230.00 |
| 36 | 73.59 | 82.26 | 92.88 | 112.37 | 137.27 | 166.79 | 199.05 | 222.04 | 245.08 |
| 37 | 79.05 | 88.15 | 99.31 | 119.84 | 146.12 | 177.41 | 211.79 | 236.42 | 261.26 |
| 38 | 84.98 | 94.54 | 106.27 | 127.88 | 155.63 | 188.79 | 225.41 | 251.80 | 278.56 |
| 39 | 91.41 | 101.45 | 113.78 | 136.54 | 165.82 | 200.95 | 239.95 | 268.22 | 297.05 |
| 40 | 98.36 | 108.91 | 121.87 | 145.82 | 176.73 | 213.94 | 255.45 | 285.72 | 316.76 |

¹reference range for non-pregnant women: <30-110 U/L(22<120 yrs) [21]

Table S 7. Percentile bands for hs-CRP and GGT during pregnancy for Indigenous Australian women

| Inflamm-atory Marker | Unit | 2.5 th | 5 th | 10 th | 25 th | 50 th | 75 th | 90 th | 95 th | 97.5 th |
|----------------------|--------|-------------------|-----------------|------------------|------------------|------------------|------------------|------------------|------------------|--------------------|
| hsCRP ¹ | (mg/L) | 0.50 | 0.90 | 1.50 | 3.00 | 6.00 | 10.00 | 18.00 | 25.00 | 34.60 |
| GGT ² | (U/L) | 5.0 | 5.0 | 6.0 | 8.0 | 11.0 | 16.0 | 26.0 | 37.0 | 52.0 |

1 Reference range for non-pregnant women: 1.00-3.00 mg/L.

2 Reference range for non-pregnant women: 9-36 U/L



