

Web-only table 2. The prevalence of demographic, psychosocial and structural factors among FSW, IBBS 2012 Nepal.

Factors	FSW N (%)
Total	610 (100)
HIV prevalence	6 (1)
Demographic	
Age in years	
16-19	85 (14)
20-24	130 (21)
≥25	395(65)
Illiterate	196(32)
Married	360(59)
Behavioural	
Use of condom with most recent client	461(75)
Consistent condom use	236(59)
Structural	
Having dependents in family	341(56)
Own home	311(51)
Forced sex in last one year	125(21)
Knowledge regarding confidential HTC	397(65)
Participated in discussion in HIV	84(29)
Participated in HIV awareness program	169(28)
Participated in AIDS day	238(39)
Participated in Condom Day	207(34)
Participated in Workshop for HIV	110(18)
Received demonstration on Condon Use	61(10)
Comprehensive Knowledge on condom	122(20)
Psychosocial	
Stigma	295(48)
Suicidal thought	210(34)
Distressed / Depressed	268 (44)
SSQS^	
Satisfied	530(87)
Dissatisfied	80(3)

*MSM/TG – Men who have sex with men/transgender, FSW – female sex workers, HIV- Human Immunodeficiency Virus., AIDS- Acquired immune deficiency syndrome, HTC-HIV Testing and Counselling, ^SSQS- The Social Support Questionnaire Score.