

BMJ Open

Experience with the systematic evaluation and reconfirmation of a national set of health system performance indicators

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2016-014772
Article Type:	Research
Date Submitted by the Author:	17-Oct-2016
Complete List of Authors:	Fekri, Omid; Canadian Institute for Health Information, Health System Performance Leeb, Kira; CIHI, Health System Performance Gurevich, Yana; CIHI, Health System Performance
Primary Subject Heading:	Health services research
Secondary Subject Heading:	Health policy
Keywords:	Consensus methods, Health system performance, Health indicators, Evaluation

SCHOLARONE™
Manuscripts

Title page

Date: October 17, 2016

Manuscript title: Experience with the systematic evaluation and reconfirmation of a national set of health system performance indicators

Authors:

Omid Fekri
Kira Leeb
Yana Gurevich

Affiliation: Canadian Institute for Health Information

Address for all authors:

Canadian Institute for Health Information
4110 Yonge Street, Suite 300
Toronto, Ontario, Canada, M2P 2B7

Keywords: Consensus methods; Health system performance; Health indicators; Evaluation

Corresponding author:

Omid Fekri
Canadian Institute for Health Information
4110 Yonge Street, Suite 300
Toronto, Ontario, Canada, M2P 2B7
Phone: (416) 549-5389
Fax: (416) 481-2950
Email: omid.fekri@utoronto.ca

Word count: 3,598

*EVALUATION OF HSP INDICATORS****Experience with the systematic evaluation and reconfirmation of a national set of health system performance indicators***

Objectives: Evaluating an existing suite of health system performance (HSP) indicators for continued reporting using a systematic criteria-based assessment and national consensus conference.

Design: Modified-Delphi approach with technical and leadership groups, an online survey of stakeholders, and convening a national consensus conference.

Setting: A national health information steward, the Canadian Institute for Health Information (CIHI).

Participants: A total of 73 participants, comprised of 61 conference attendants/stakeholders from across Canada, and 12 national health information steward staff.

Primary and secondary outcome measures: Indicator dispositions of retention, additional stakeholder consultation, further redevelopment, or retirement.

Results: Four dimensions (usability, importance, scientific soundness, and feasibility) typically used to select measures for reporting were expanded to 18 criteria grouped under the four dimensions through a process of research and testing. Definitions for each criterion were developed and piloted. Once the definitions were established, 56 of CIHI's publicly reported HSP indicators were evaluated against the criteria using modified Delphi approaches. Of the 56 HSP indicators evaluated, 9 measures were ratified for retirement, 7 were identified for additional consultation, and 3 for further research and development. A pre-Consensus Conference survey soliciting feedback from stakeholders on indicator recommendations received 48 responses (response rate of 79%).

Conclusions: A systematic evaluation of HSP indicators informed the development of objective recommendations for continued reporting. The evaluation was a fruitful exercise to identify technical considerations for calculating indicators, furthering our understanding of how measures are used by stakeholders, as well as harmonizing actions that could be taken to ensure relevancy, reduce indicator chaos, and build consensus with stakeholders.

Article summary/Strengths and limitations of this study:

- This exercise utilized an extensive suite of criteria to evaluate health system performance indicators.
- Multiple evaluation modalities were held to solicit feedback from evaluators.
- A large number of stakeholders participated in an in-person consensus conference.
- Assessment criteria and processes may not apply in other evaluation contexts.

1 Introduction

2 Health indicators offer valuable insight into the performance of health systems and the health of populations. As the
3 discipline of health system performance (HSP) measurement has grown over the decades, so too have the number of
4 available health measures. In Europe alone, journal publications related to performance indicators increased at a rate of
5 ~20% annually between 2000 and 2009 [1]. However, the adage of *more is better* is not entirely true for this discipline, and
6 runs contrary to the notion of providing concise findings [2]. Health measure producers and users are constrained with
7 finite resources, and must make important decisions on which indicators they deem important, have high utility, are valid,
8 and are feasible. Periodic reviews of indicators and conceptual frameworks can ensure their continued relevance and
9 efficacy [3].
10

11
12
13 Two national agencies, the Canadian Institute for Health Information (CIHI) and Statistics Canada, have collaborated for
14 more than 15 years on developing and publicly reporting health measures for health regions, provinces and territories as
15 part of the Health Information Roadmap [4]. Over the years, the number of indicators has increased from 13 in 2000 to
16 more than 80 in 2014. This in part reflects the growing information needs of health care systems in general. For example,
17 new indicators measuring outcomes, wait times, and patient safety were the areas of focus for development in recent
18 years. CIHI also expanded its indicator reporting over the years by refining the granularity of public reporting, and in 2007
19 began public reporting of health indicators for acute care hospitals in Canada. The indicators were developed and reported
20 on according to the CIHI–Statistics Canada Health Indicator Framework [5]. In 2012, the suite of publicly available indicators
21 at the hospital level was expanded substantially and in 2015, was expanded again to include indicators for long-term care
22 homes.
23
24

25
26 After a period of rapid growth in public reporting of indicators and capacity-building activities across the country, health
27 system managers identified that having too many indicators to monitor and respond to was not achieving the goal of
28 helping understand how well the health care system was performing. This phenomenon was termed “indicator chaos” [6],
29 and initiated a new focus on streamlining indicator reporting and development activities.
30

31 Partly in response to this notion of indicator chaos, but also in efforts to ensure relevancy and efficiency, CIHI initiated a
32 program of work aimed at streamlining health system reporting in Canada. As part of this work, CIHI developed a new
33 Health System Performance Framework to better reflect the relationship between indicator measurement and health
34 system goals [5]. CIHI also recognized the need to ensure that the indicators being produced and reported reflected these
35 goals and contributed to a broader understanding of health system performance rather than continuing to add to the
36 reporting and monitoring burden across the country. This required a systematic indicator evaluation process that could be
37 repeated periodically to inform indicator reporting initiatives across the organization and possibly beyond.
38
39

40 Coincidentally, every five years (1999, 2004, 2009, 2014), CIHI and Statistics Canada invite stakeholders from across the
41 country to a national Consensus Conference on Health Indicators to discuss priority-setting of indicator development and
42 reporting for the next half decade [7,8,9,10]. The latest such conference (held in 2014) provided an opportunity to present
43 the results of the internal evaluation of publicly reported indicators and to validate the results with stakeholders.
44
45

46 This paper describes CIHI’s experience piloting the evaluation of a set of HSP indicators using a systematic criteria-based
47 assessment tool and process. The results of the pilot — including achieving reconfirmation through a national consensus
48 process — and possible next steps for broader implementation of the strategy are also presented in the paper.
49
50
51
52
53
54
55
56
57
58
59
60

Methods

This project had four distinct components:

1. Process and criteria development for systematic evaluation of HSP indicators
2. Internal CIHI modified-Delphi sessions
3. Pre-conference survey of stakeholders on indicator recommendations
4. Presentation and ratification of results at the national Consensus Conference

Systematic Evaluation of HSP Indicators

The Institute of Medicine's (IOM's) *Recommendations for Measure Selection Criteria* [11] — Usability, Importance, Scientific Soundness and Feasibility — are consistently used in the evaluation and selection of health measures [12]. While many examples in the literature employ these four domains of criteria, we saw the need to expand the dimensions to include other criteria within three of the four domains. Through a process of research and testing, we arrived at a total of 18 criteria points organized around the four IOM domains (see Table 1) that were feasible to apply and that held meaning to our project objective regarding continued reporting of indicators. There is congruence between these criteria and CIHI's Data Quality Framework [13] dimensions of Accuracy, Timeliness, Comparability, Usability and Relevance. Over a period of two months, 56 of CIHI's suite of HSP indicators were assessed against these 18 criteria. To aid evaluators in their subsequent reviews, we created a one-page summary for each indicator denoting results for each evaluation criterion [14].

Internal CIHI modified-Delphi sessions

Two groups within CIHI participated in the evaluation. First, a technical group of experts (n=6) (comprised of epidemiologists, methodologists and statisticians) independently reviewed each indicator and criterion point, and provided a Likert Scale score between 1 and 9 recommending continued reporting or otherwise (with 1 representing strong disagreement for continued reporting and 9 indicating strong agreement for continued reporting). Evaluators were instructed to produce a Likert score and disposition recommendation based on a holistic assessment of all 18 criteria. We therefore forewent weighting evaluation criteria. This allowed for flexibility and context in instances where some criteria proved more informative than others.

Likert scores were averaged and presented at an internal in-person Delphi session as a basis for discussions, but were not automatically tied to a final result of continued indicator reporting. Beginning with the lowest average scores, each indicator was discussed, pertinent commentary synthesized, and a final consensus reached on a disposition recommendation. Disposition options for indicators were Retain, Recommend Further Research and Development (R&D) or Consultation, or Retire.

Recommendations of the technical group's Delphi sessions were then presented to the CIHI HSP leadership group (n=6) (comprised of senior managers and researchers) who performed a similar exercise. First, they were asked to independently review all results to date (including indicator assessments and Likert scores, commentary and disposition recommendations). Results of their individual assessments were collated and presented at an in-person session. Disposition recommendations for each of the 56 indicators were solidified. The RAND/UCLA Appropriateness Method [15] guided our internal iterative modified-Delphi sessions.

Pre-conference survey of stakeholders on indicator recommendations

A pre-Consensus Conference survey solicited initial feedback on recommendations. The online survey was available for a period of six weeks prior to the conference. Consensus Conference participants were chosen from an existing list of CIHI partners, stakeholders, and clients; common roles of participants were hospital/health region CEOs, academics and researchers, representatives from ministries of health, clinicians, and national collaboration partners. An electronic survey was emailed to conference participants along with background documentation on the evaluation process, methodology, and recommendations. The survey asked respondents whether they Agreed, Disagreed or had No Opinion on the recommendation to retire select HSP indicators as per recommendations from CIHI's internal review.

National Consensus Conference presentation

EVALUATION OF HSP INDICATORS

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

There were 61 participants at the invitational in-person Consensus Conference held in Toronto on October 16 and 17, 2014. Results of the pre-conference survey were presented. A threshold of 70% agreement by respondents was used to automatically pass recommendations or to otherwise hold further group discussion at the conference. An external moderator facilitated discussion and voting on final indicator dispositions.

For peer review only

Results

Systematic evaluation of HSP indicators

The systematic evaluation of HSP indicators was a holistic process considering 18 criteria points. Some criteria differentiated indicators more than others. For example, a small number of criteria resulted in mostly uniform findings for the suite of HSP indicators. However, when assessed alongside remaining criteria, important contextual considerations can be gleaned. Notable findings are summarized below by criterion.

Usability

The *Granularity of reporting* criterion identified nuances inherent within public reporting purposes. There are approximately 100 administrative health regions in Canada, and approximately 600 acute care hospitals. Twenty-nine indicators are reported at the regional level, and 27 are reported at the hospital/facility level. All indicators are reported at an aggregate provincial/territorial and national level.

With respect to *Pan-Canadian coverage*, 44 of 56 indicators provided complete pan-Canadian coverage (all provinces and territories). The province of Quebec does not have available or comparable data for a dozen indicators. Similar to the criterion of pan-Canadian coverage, the *Comprehensiveness* criterion assessed the inclusiveness of health services providers that submit data toward the indicator. For example, the mental illness hospitalization indicator includes data on mental health patients treated in general hospitals only, while hospitalizations at free-standing psychiatric institutions are not included due to the differences in data collection.

For the *Usage* criterion, we polled CIHI HSP staff responsible for interacting with clients on indicators and data requests. This provided a proxy for the level and extent of the indicator's usage by clients. The 56 indicators under evaluation were rated as High (n=33), Medium (n=15) or Low usage (n=8).

With regard to *Dimensionality*, breakdowns of indicator results by dimensions of sex and socio-economic status (SES) are available where applicable. Thus, 15 indicators are reportable by SES and 14 are reportable by sex.

In terms of *Timeliness, Reporting frequency and Accessibility*, all 56 indicators were publicly reported annually within 10 months of the relevant data being available for analysis. At the time of the evaluation, all HSP indicators were accessible publicly through online publications such as the Health Indicators e-Publication. Additionally, a majority of facility-level indicators are available to providers through private online tools to allow for more granular breakdowns of results and peer comparative reports.

For the *Trendability* criterion, it was found that time trends vary by indicator. For example, the set of facility-level indicators was largely first reported beginning with 2007 data. Results for select regional indicators dated back to 1997. Overall, regional indicators possessed almost twice as many available years of results compared with facility indicators, a nature of the timing of reporting programs.

Importance

As a proxy measure for *Relevance*, an environmental scan was conducted to understand stakeholder utilization of indicators. A total of 232 instances online were recorded. The top five indicators were Hospital Standardized Mortality Ratio (HSMR) (n=23), 30-Day Overall Readmission (n=18), Wait Times for Hip Fracture Repair (n=17), Ambulatory Care Sensitive Conditions (n=14), and Caesarean Section Rate (n=13).

Detailed statements on the *Actionability* of each indicator were provided to evaluation participants. Specifically, summations on the purpose of indicator, strengths, caveats, and scientific evidence in support were considered.

To measure the degree of *Stakeholder follow-up*, we reviewed all instances of patient-level data requests from providers. In 2013-14, there were 298 requests, with 11 facility-level readmission indicators accounting for 58% of all requests (n=173).

1 The criterion *Sufficient volumes* quantifies the proportion of indicator results that are suppressed per CIHI's data privacy
2 protocols. In general, indicator results with cell counts less than five are suppressed, and results based on less than 50
3 denominator cases per hospital are flagged as low volume and unstable rates. Facility-level indicators are particularly
4 affected by low volumes and suppressed results: 23 of 27 facility-level indicators had at least one-fifth of all results flagged
5 as low volume. A further 7 of these indicators had at least one-fifth of all results suppressed due to small cell counts. At the
6 extreme, we note the 28-Day Readmission After Stroke and Acute Myocardial Infarction (AMI) indicators with ~75% low
7 volume rates and one-third of all results suppressed.
8
9

10 We performed *Significance of variation* analysis to determine the variability within indicator results. For example, the Hip
11 Fracture Surgical Procedures Performed Within 48 Hours indicators (both Within One and Across facilities) had the lowest
12 relative standard deviation values of 16% and 17%, respectively, indicating minimal differences across indicator results.
13
14

15 Scientific Soundness

16 The criterion *Data quality* garnered the greatest discussion during Delphi reviews. Limitations of using administrative data
17 were considered. Examples of concern include the inability to assess indications for angiography for AMI patients for the
18 indicator Use of Coronary Angiography Following AMI, and the ability to properly identify denominator cases for the
19 Hysterectomy indicator.
20
21

22 The evaluation revealed that *Validity reviews* were performed for each indicator on an annual basis. These included
23 significance testing of risk factors, monitoring of diagnosis and procedure coding updates, and outlier and significant change
24 analyses. Indicators recommended for further consultation and R&D were identified as such mainly for the purpose of
25 seeking feedback from stakeholders on the validity and clinical relevance of current calculation methodologies.
26
27

28 The criterion *Participation bias* assessed whether data submission and participation in the calculation of indicator results
29 were a nature of voluntary participation. All but two indicators — Physician Specialists and General/Family Physicians per
30 100,000 Population — required mandatory participation. In other instances, such as indicators produced for long-term care
31 facilities, participation is not yet mandatory across the country, and therefore results published may contain a participation
32 bias.
33
34

35 Feasibility

36 *Production cost* was considered based on the extent of staff resources required to produce each indicator. Indicators with
37 complex linkages across multiple databases and those requiring building of episodes of care necessitate a larger degree of
38 resources.
39
40

41 Modified-Delphi sessions of CIHI technical and leadership groups

42 Mean Likert scores, recommendations and rationale are noted in tables 2, 3 and 4. Nine indicators were recommended as
43 candidates for retirement (Table 3), seven were identified as requiring additional consultation and three were
44 recommended to undergo further redevelopment (Table 2). Thirty-five indicators were recommended for retention (Table
45 4). The rationale to retain these HSP indicators was based on holistic decisions across 18 criteria points. Although retained
46 indicators correlate strongly with high mean Likert scores, this was only one contributor to the recommendation.
47
48

49 CIHI leadership and technical groups identified indicators for additional consultation and redevelopment. These indicator
50 recommendations were not forwarded to Consensus Conference participants, but were instead identified for internal R&D
51 efforts in the interim.
52

53 Pre-Consensus Conference survey

54 Forty-eight Consensus Conference participants completed the online survey (response rate of 79%). Eighty-five percent of
55 conference participants had more than 10 years of health care experience. Geographic distribution of respondents
56 correlated well with Canada's population across provinces/territories. The mean survey agreement score (as a percentage
57 of responses) for all nine indicators proposed for retirement was 70%, and was used as a benchmark for automatic
58 ratification.
59
60

EVALUATION OF HSP INDICATORSNational Consensus Conference

Of the nine indicators recommended for retirement, six received more than 70% agreement as a proportion of responses in the pre-conference survey, and therefore were automatically accepted for retirement (Table 3). The remaining three indicators were discussed as a group, and subsequently also ratified for retirement by conference participants. The majority of indicators recommended for retirement were condition-specific readmission indicators. Ultimately, the decision to retire these indicators was based on appropriateness for continued public reporting. While these indicators were ratified for retirement over concerns of rate stability and small numbers, facilities can continue to calculate and monitor these indicators through CIHI private reporting tools. Consensus on retiring these indicators was achieved with greater ease given that a provider's capacity to continue to privately monitor performance would be maintained.

Two contextual health human resources indicators at the regional level — Physician Specialists and General/Family Physicians per 100,000 Population — were also included in the modified Delphi review process, and rated low in Likert Scale scoring (both received a mean score of 3.2). While these indicators provide some context on HSP characteristics, they are already reported elsewhere within CIHI. It was agreed to continue reporting on these indicators but outside of the HSP framework.

Table 4 lists 35 HSP indicators retained for continued public reporting. Although retained indicators correlate strongly with high mean Likert scores, this was only one contributor to the recommendation. For example, the regional level Caesarean Section Rate indicator received a mean Likert score of 4.8 from the technical group, but was retained for public reporting after discussion by the leadership group due to continued concerns over high rates in Canada and therefore a need for continued monitoring.

Discussion

This exercise proved to be an informative, objective, systematic, transparent, inclusive, and likely repeatable process for evaluating and reconfirming a national set of HSP indicators. Overall, the approach of using 18 sub-criteria was manageable and informative, with feedback from participants that the added information and context made it easier to make a final disposition recommendation for each indicator. The overall timeline of the evaluation process from inception to completion was 18 months. Three distinct phases stand out, each requiring approximately six months to complete: initial R&D of the evaluation plan, executing the evaluation internally at CIHI, and achieving consensus across stakeholders.

An initial Likert score of indicators provides a baseline to proceed with group Delphi reviews. We found it beneficial to begin with the lowest scores and work our way to the highest rated indicators. We also found it operational to have our technical group first review indicators and to pass on recommendations to a leadership group that would consider these in addition to their knowledge and understanding of the use of HSP information in the field. The iterative process of having participants first review indicators independent of other Delphi members and to then convene as a group to discuss findings allowed for a balanced and participatory discussion among participants. These iterative methods ensured a summative process whereby findings were transparent and confirmed at each stage.

The national Consensus Conference provided an opportunity to pilot-test the results of a rigorous, mostly internal methodology for evaluating indicators produced by CIHI. While stakeholders appreciated the opportunity to review and ratify our findings, stakeholders indicated that, going forward, they were comfortable with CIHI implementing a systematic evaluation of the indicators and making decisions about reporting. There was congruence in opinion on the suitability of HSP indicators for public reporting throughout the evaluation process, beginning with Likert scores and assessments from CIHI technical staff, to CIHI leadership, and lastly with stakeholders.

Strengths and weaknesses of the study

While the overall evaluation required considerable time and resources, there are important benefits to such a comprehensive process. We ensured a transparent and sequential evaluation, whereby discourse and findings were accumulated and presented in a summarized manner at each phase. We solicited feedback from evaluators both independently and in a group manner (also polling online and in-person at the consensus conference). An external moderator facilitating discussion can ensure independence during the consensus process. The aforementioned processes have been described as favourable conceptual approaches to aid exercises of indicator development, maintenance, and evaluation [14].

Shekelle [16] notes there is little agreement on methodologies for developing performance indicators, and this can be said of the same regarding their evaluation. Nonetheless, Stelfox and Straus [14] emphasize the importance of clearly establishing the chosen evaluation criteria in advance of launching a consensus process. In the majority of the studies we reviewed and cite, a smaller number of evaluation criteria were applied: most often, usability, importance, scientific soundness, and feasibility (or a variation thereof that drew upon similar domains). Conversely, we found it helpful to apply multiple sub-criteria to comprehensively reflect the evaluation of indicators for their suitability of ongoing public reporting. Furthermore, providing a more granular evaluation schema for participants ensured more consistent definitions of domains and structured evidence/results for evaluators' consideration. Nonetheless, while these evaluation criteria were informative and applicable to this precise context, not all would apply for other evaluation purposes. Further efforts are necessary to determine the level of customization required to ensure that the process and criteria are applicable to other sectors of care and types of indicators.

In addition to convening an in-person consensus conference (or expert panel) to evaluate indicators, Santana et al [17] forwarded their evaluation survey to 101 trauma centres across four countries involved in the use and assessment of injury-care indicators. Moreover, a novel subsequent processes has been described by Bobrovitz et al [18] whereby the discussion occurring throughout the consensus conference is transcribed and undergoes qualitative content analysis to identify key themes raised throughout the deliberations. These additional activities can provide complementary evidence to the evaluation process, such as qualitative findings to an otherwise objective and quantitative exercise, and reaching a broader group of stakeholders and users of health measures.

*EVALUATION OF HSP INDICATORS***Conclusion**

The proliferation of health measures required to fulfill reporting gaps occurred with minimal consideration to alignment and utility with pre-existing indicators. Not surprisingly, then, stakeholders were overwhelmingly in favour of implementing a process that would result in a leaner, more applicable suite of HSP indicators.

CIHI will gradually expand this evaluation methodology to applicable sectors of care. We will also continue to work with external partners to reduce indicator chaos and increase alignment with reporting requirements across the country [6].

This exercise generated identified analytical alignment actions that can be taken at CIHI throughout indicator production and maintenance with a view to reduce indicator chaos. Furthermore, we gained new knowledge about how the HSP indicators we produce are used by stakeholders both through an internet-based environmental scan and via discussions held at the Consensus Conference [10].

In line with established practices of convening a Consensus Conference every five years, we feel that it is highly beneficial to inform those discussions with a wholesale and systematic criteria-based review of indicators just prior. A broad consultation process encompassing diverse public health stakeholders from across the country helps ensure the development and use of indicators most appropriately reflecting the health of populations and the performance of health systems [19]. Similarly, a retrospective exercise on national HSP practices can identify important lessons, of which the selection of indicators suitable for public reporting is an integral component [20].

Contributors

OF participated in the design of the evaluation, performed data analysis, facilitated internal Delphi sessions, and drafted the manuscript. KL participated in the design of the evaluation, facilitated internal Delphi sessions, organized the national consensus conference, and drafted the manuscript. YG participated in the design of the evaluation, facilitated internal Delphi sessions, and drafted the manuscript. All authors reviewed and approve the final version.

Funding statement

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests

We have read and understood the BMJ policy on declaration of interests and declare that we have no competing interests.

Data sharing statement

No additional data are available.

For peer review only

References

1. Klazinga N, Fischer C, ten Asbroek A. Health services research related to performance indicators and benchmarking in Europe. *J Health Serv Res Policy*, 2011;16:38-47. http://hsr.sagepub.com/content/16/suppl_2/38.long (accessed 16 October 2016).
2. Kelley E, Arispe I, Homes J. Beyond the initial indicators: lessons from the OECD Health Care Quality Indicators Project and the US National Healthcare Quality Report. *Int J Qual Health Care*, September 2006:45-51. http://intqhc.oxfordjournals.org/content/18/suppl_1/45.long (accessed 16 October 2016).
3. Carinci F, Van Gool K, Mainz J et al. Towards actionable international comparisons of health system performance: expert revision of the OECD framework and quality indicators. *Int J Qual Health Care*, 2015:1-10. <http://intqhc.oxfordjournals.org/content/27/2/137> (accessed 16 October 2016).
4. Canadian Institute for Health Information. Roadmap Initiative ... Launching the Process. Ottawa:2000. https://www.cihi.ca/en/profile_roadmap_launch_pdf_en.pdf (accessed 16 October 2016).
5. Canadian Institute for Health Information. A Performance Measurement Framework for the Canadian Health System. Ottawa:2013. https://secure.cihi.ca/free_products/HSP-Framework-ENweb.pdf (accessed 16 October 2016).
6. Saskatchewan Health Quality Council. Think big, start small, act now: tackling indicator chaos: a report on a national summit: Saskatoon, May 30-31, 2011. <http://hqc.sk.ca/Portals/0/documents/tracking-indicator-choas.pdf> (accessed 16 October 2016).
7. Canadian Institute for Health Information. National Consensus conference on Population Health Indicators. Ottawa: 1999. https://secure.cihi.ca/free_products/phi.pdf (accessed 16 October 2016).
8. Canadian Institute for Health Information. The Health Indicators Project: The Next 5 Years. Report from the Second Consensus Conference on Population Health Indicators. Ottawa:2005. https://secure.cihi.ca/free_products/Consensus_Conference_e.pdf (accessed 16 October 2016).
9. Canadian Institute for Health Information. The Health Indicators Project: Report From the Third Consensus Conference on Health Indicators. Ottawa:2009. https://secure.cihi.ca/free_products/82-230-XWE_e.PDF (accessed 16 October 2016).
10. Canadian Institute for Health Information. Rethink, Renew, Retire: Report from the Fourth Consensus Conference on Evaluating Priorities for Canada's Health Indicators. Ottawa: 2015. https://secure.cihi.ca/free_products/Rethink_Renew_Retire.pdf (accessed 16 October 2016).
11. Hurtado M, Swift E, Corrigan J, editors, Committee on the National Quality Report on Health Care Delivery, Board on Health Care Services. Envisioning the National Health Care Quality Report, Washington, DC, 2001, National Academy press. http://www.ncbi.nlm.nih.gov/books/NBK223318/pdf/Bookshelf_NBK223318.pdf (accessed 16 October 2016).
12. Mattke S. When should measures be updated? Development of a conceptual framework for maintenance of quality-of-care measures. *Qual Saf Health Care* 2008;17:182-186. <http://qualitysafety.bmj.com/content/17/3/182.abstract> (accessed 16 October 2016).
13. Canadian Institute for Health Information. The CIHI Data Quality Framework, 2009. Ottawa:2009. https://www.cihi.ca/en/data_quality_framework_2009_en.pdf (accessed 16 October 2016).
14. Stelfox, H & Straus, S. Measuring quality of care: considering conceptual approaches to quality indicator development and evaluation. *J Clinical Epidemiology*. 2013; 66: 1328-1337.
15. Fitch K, Bernstein S, Aguilar M, Burnand B, LaCalle JR, Lazaro P, et al. The RAND/UCLA Appropriateness method user's manual. Santa Monica, CA: RAND Corporation, 2001. http://www.rand.org/pubs/monograph_reports/MR1269 (accessed 16 October 2016).
16. Shekelle, P. Quality indicators and performance measures: methods for development need more standardization. *J Clinical Epidemiology*. 2013; 66: 1338-1339.
17. Santana MJ, Stelfox HT, Asbridge M, Ball CG, Cameron P, Dyer D et al. Development and evaluation of evidence-informed quality indicators for adult injury care. *Annals of Surgery*. 2014 Jan;259(1):186-192.
18. Bobrovitz et al.: A qualitative analysis of a consensus process to develop quality indicators of injury care. *Implementation Science*. 2013 8:45.

- 1 19. Klazinga N, Stronks K, Delnoij D, Verhoeff A. Indicators without a cause. Reflections on the development and use of
2 indicators in health care from a public health perspective. *Int J Qual Health Care*, 2001;13:433-438.
3 <http://intqhc.oxfordjournals.org/content/intqhc/13/6/433.full.pdf> (accessed 16 October 2016).
- 4 20. van den Berg, M et al.: The Dutch health care performance report: seven years of health care performance
5 assessment in the Netherlands. *Health Research Policy and Systems* 2014 12:1.
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

EVALUATION OF HSP INDICATORS

Table 1 – Evaluation criteria

Domain/Criterion	Definition
<u>Usability</u>	
· Granularity of reporting	Reporting at national, provincial/territorial, regional and facility levels
· Pan-Canadian coverage	Extent of participation from all provinces and territories
· Comprehensiveness	Proportion of providers submitting data for the indicator
· Usage	Level and extent of usage
· Dimensionality	Ability to break down results by age, sex, socio-economic status, and other dimensions
· Timeliness	Latest year of available results
· Reporting frequency	Whether indicator is reported quarterly, annually or other
· Accessibility	Whether the indicator is publicly and/or privately reported
· Trendability	Number of years of available results for trending
<u>Importance</u>	
· Relevance	Environmental scan identified uses of indicator by stakeholders
· Actionability	Extent to which providers can meaningfully influence the indicator
· Stakeholder follow-up	Number of data and methodological requests within last fiscal year
· Sufficient volumes	Percentage of results suppressed (due to low counts)
· Significance of variation	Degree of variation across reported values
<u>Scientific Soundness</u>	
· Data quality	Strength of data quality, ability to validate results, based on standards
· Validity review	Extent and frequency of reviewing indicator's validity/methodology
· Participation bias	Mandatory or voluntary participation by providers
<u>Feasibility</u>	
· Production cost	Extent of staff/resources to produce indicator

Table 2 – Indicators identified for additional consultation and further redevelopment

Type	Indicator	Mean Likert Score	Rationale
ADDITIONAL CONSULTATION			
Region	Hip Replacement	5.0	There are concerns of utility and actionability for these indicators as they represent procedure counts per population.
	Knee Replacement	4.8	
	Coronary Artery Bypass Graft (CABG)	6.6	
	Percutaneous Coronary Intervention (PCI)	6.6	
	Cardiac Revascularization	6.6	
Facility	Vaginal Birth After Caesarean Section	4.4	There are concerns of validity and utility for these indicators.
	Birth Trauma	5.4	
FURTHER REDEVELOPMENT			
Region	Hysterectomy	4.4	R&D is required to improve identification of appropriate denominator cases.
Facility	Nursing Sensitive Adverse Events for Medical Patients	6.8	There is an opportunity for incorporation within newly developed Hospital Harm indicator.
	Nursing Sensitive Adverse Events for Surgical Patients	6.8	

EVALUATION OF HSP INDICATORS

Table 3 – Indicators recommended for retirement

Type	Indicator	Mean Likert Score	Rationale	Pre-Consensus Conference Survey Agreement (as a % of responses)
Facility	28-Day Readmission After Prostatectomy	5.2	These indicators have low volumes of cases leading to unstable rates as well as to the suppression of a large number of results for public reporting. Furthermore, these cases are included in the Surgical/Medical Readmission indicators, and can still be derived through private reporting tools.	82%*
	28-Day Readmission After Hysterectomy	5.6		80%*
	90-Day Readmission After Knee Replacement	6.4		73%*
	90-Day Readmission After Hip Replacement	6.4		72%*
	28-Day Readmission After Stroke	6.2		58%
	Use of Coronary Angiography Following AMI	6.4	Angiography may not be indicated for every AMI patient, depending on his or her clinical history, and the clinical appropriateness of angiography is difficult to ascertain from the administrative hospitalization data. Therefore, it is challenging to interpret and compare the results for this indicator.	78%*
	Hip Fracture Surgical Procedures Performed Within One Facility (48 Hours)	6.4	This indicator does not measure the true proportion of surgeries performed within 48 hours of admission to an acute care hospital, since it does not account for transfers across hospitals. Many patients are transferred from their initial admitting acute care facility to another facility for surgery. The indicator Hip Fracture Surgical Procedures Performed Within 48 Hours, which measures total time across all acute care facilities, will continue to be produced and reported on.	72%*
	28-Day Readmission After AMI	6.4	Concerns have been raised regarding hospitals' ability to take action on this indicator. It is felt that with the regionalization of cardiac care, it is more appropriate to measure readmission after AMI at the regional level (by patient residence) than at the hospital level. In addition, having a low volume of cases leads to unstable rates and to the suppression of a large number of results for public reporting. Therefore, it was proposed to keep the Readmission After AMI indicator at the regional level and to retire the facility-level indicator. Furthermore, readmissions after AMI are included in the 30-Day Overall Readmission indicator at the facility level.	59%
	Primary Caesarean Section Rate	4.6	A new indicator (Low-Risk Caesarean Section) measures the rate of deliveries via Caesarean section among singleton term cephalic pregnancies for women without placenta previa or previous C-section. Since this new indicator is limited to women who have not had a previous C-section, it can take the place of Primary Caesarean Section Rate and be a better indicator of appropriateness.	57%

* Passing the threshold (of 70% agreement among responses) for automatic ratification.

Table 4 – Indicators retained

Type	Indicator	Mean Likert Score
Region	30-Day AMI In-Hospital Mortality	8.8
	30-Day Stroke In-Hospital Mortality	8.8
	Hospital Standardized Mortality Ratio (HSMR)	8.8
	Ambulatory Care Sensitive Conditions	8.6
	Wait Times for Hip Fracture Repair	8.4
	30-Day Readmission for Mental Illness	7.8
	Repeat Hospital Stays for Mental Illness	7.8
	Self-Injury Hospitalization	7.6
	30-Day AMI Readmission	7.4
	Hospitalized Hip Fracture Event	7.2
	Hospitalized Strokes	7.2
	Hospitalized AMI Event	7.0
	Inflow/Outflow Ratio	7.0
	30-Day Readmission: Patients Age 19 and Younger	6.8
	30-Day Obstetric Readmission	6.8
	30-Day Medical Readmission	6.8
	30-Day Surgical Readmission	6.4
	Mental Illness Patient Days	6.2
	Mental Illness Hospitalization	6.0
	Injury Hospitalization	5.4
Caesarean Section Rate	4.8	
Facility	30-Day AMI In-Hospital Mortality	8.8
	30-Day Stroke In-Hospital Mortality	8.6
	Hip Fracture Surgery Within 48 Hours	8.4
	30-Day Overall Readmission	8.0
	30-Day In-Hospital Mortality Following Major Surgery	8.0
	30-Day Readmission: Patients Age 19 and Younger	7.8
	30-Day Obstetric Readmission	7.8
	30-Day Medical Readmission	7.6
	30-Day Surgical Readmission	7.4
	In-Hospital Hip Fracture In Elderly (Age 65+) Patients	7.4
	Obstetric Trauma — Vaginal Delivery With Instrument	7.4
	Obstetric Trauma — Vaginal Delivery Without Instrument	7.4
	Caesarean Section Rate	6.8
	Low-Risk Caesarean Section	6.8

BMJ Open

A systematic approach to evaluating and confirming the utility of a suite of national health system performance (HSP) indicators in Canada: a modified Delphi study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2016-014772.R1
Article Type:	Research
Date Submitted by the Author:	24-Dec-2016
Complete List of Authors:	Fekri, Omid; Canadian Institute for Health Information, Health System Performance Leeb, Kira; CIHI, Health System Performance Gurevich, Yana; CIHI, Health System Performance
Primary Subject Heading:	Health services research
Secondary Subject Heading:	Health policy
Keywords:	Consensus methods, Health system performance, Health indicators, Evaluation

SCHOLARONE™
Manuscripts

Title page

1
2
3
4 **Date:** December 16, 2016
5
6

Manuscript title:

8 A systematic approach to evaluating and confirming the utility of a suite of national health system performance (HSP)
9 indicators in Canada: a modified Delphi study
10

Authors:

12 Omid Fekri
13 Kira Leeb
14 Yana Gurevich
15
16

17
18
19 **Affiliation:** Canadian Institute for Health Information
20

Address for all authors:

22 Canadian Institute for Health Information
23 4110 Yonge Street, Suite 300
24 Toronto, Ontario, Canada, M2P 2B7
25
26
27

28
29 **Keywords:** Consensus methods; Health system performance; Health indicators; Evaluation
30
31

Corresponding author:

32 Omid Fekri
33 Canadian Institute for Health Information
34 4110 Yonge Street, Suite 300
35 Toronto, Ontario, Canada, M2P 2B7
36 Phone: (416) 549-5389
37 Fax: (416) 481-2950
38 Email: omid.fekri@utoronto.ca
39
40
41
42

43
44 **Word count:** 4,328
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS

1 A systematic approach to evaluating and confirming the utility of a suite of national health system performance (HSP)
2 indicators in Canada: a modified Delphi study
3

4
5 **Objectives:** Evaluating an existing suite of health system performance (HSP) indicators for continued reporting using a
6 systematic criteria-based assessment and national consensus conference.
7

8 **Design:** Modified-Delphi approach with technical and leadership groups, an online survey of stakeholders, and convening a
9 national consensus conference.
10

11 **Setting:** A national health information steward, the Canadian Institute for Health Information (CIHI).
12
13

14 **Participants:** A total of 73 participants, comprised of 61 conference attendants/stakeholders from across Canada, and 12
15 national health information steward staff.
16

17 **Primary and secondary outcome measures:** Indicator dispositions of retention, additional stakeholder consultation, further
18 redevelopment, or retirement.
19

20
21 **Results:** Four dimensions (usability, importance, scientific soundness, and feasibility) typically used to select measures for
22 reporting were expanded to 18 criteria grouped under the four dimensions through a process of research and testing.
23 Definitions for each criterion were developed and piloted. Once the definitions were established, 56 of CIHI's publicly
24 reported HSP indicators were evaluated against the criteria using modified Delphi approaches. Of the 56 HSP indicators
25 evaluated, 9 measures were ratified for retirement, 7 were identified for additional consultation, and 3 for further research
26 and development. A pre-Consensus Conference survey soliciting feedback from stakeholders on indicator
27 recommendations received 48 responses (response rate of 79%).
28
29

30 **Conclusions:** A systematic evaluation of HSP indicators informed the development of objective recommendations for
31 continued reporting. The evaluation was a fruitful exercise to identify technical considerations for calculating indicators,
32 furthering our understanding of how measures are used by stakeholders, as well as harmonizing actions that could be taken
33 to ensure relevancy, reduce indicator chaos, and build consensus with stakeholders.
34
35

Article summary/Strengths and limitations of this study:

- 37 • This exercise utilized an extensive suite of criteria to evaluate health system performance indicators.
- 38 • Multiple evaluation modalities were used to solicit feedback from evaluators.
- 39 • A large number of stakeholders participated in an in-person consensus conference.
- 40 • Assessment criteria and processes may not apply in other evaluation contexts.
- 41
- 42
- 43
- 44
- 45
- 46
- 47
- 48
- 49
- 50
- 51
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

Introduction

Health indicators offer valuable insight into the performance of health systems and the health of populations. As the discipline of health system performance (HSP) measurement has grown over the decades, so too have the number of available health measures. In Europe alone, journal publications related to performance indicators increased at a rate of ~20% annually between 2000 and 2009 [1]. However, continuing to increase the number of indicators reported runs contrary to, and inhibits, the provision of concise findings on the performance of health systems [2]. Health measure producers and users are constrained with finite resources, and must make important decisions on which indicators they deem important, have high utility, are valid, and are feasible. Periodic reviews of indicators and conceptual frameworks can ensure their continued relevance and efficacy [3].

Two national agencies, the Canadian Institute for Health Information (CIHI) and Statistics Canada, have collaborated for more than 15 years on developing and publicly reporting health measures for health regions, provinces and territories as part of the Health Information Roadmap [4]. Over the years, the number of indicators has increased from 13 in 2000 to more than 80 in 2014. This in part reflects the growing information needs of health care systems in general. For example, new indicators measuring outcomes, wait times, and patient safety were the areas of focus for development in recent years. CIHI also expanded its indicator reporting over the years by refining the granularity of public reporting, and in 2007 began public reporting of health indicators for acute care hospitals in Canada. The indicators were developed and reported on according to the CIHI–Statistics Canada Health Indicator Framework [5]. In 2012, the suite of publicly available indicators at the hospital level was expanded substantially and in 2015, was expanded again to include indicators for long-term care homes.

After a period of rapid growth in public reporting of indicators likely due to the rising demand for accountability and quality improvement data as well as increases in capacity-building activities across the country, health system managers identified that having too many indicators to monitor and respond to was not achieving the goal of helping understand how well the health care system was performing. In 2010, this phenomenon was coined “indicator chaos” [6], and initiated a new focus on streamlining indicator reporting and development activities.

Partly in response to this notion of indicator chaos, but also in efforts to ensure relevancy and efficiency, CIHI initiated a program of work aimed at streamlining health system reporting in Canada. As part of this work, CIHI developed a new Health System Performance Framework to better reflect the relationship between indicator measurement and health system goals [5]. CIHI also recognized the need to ensure that the indicators being produced and reported reflected these goals and contributed to a broader understanding of health system performance rather than continuing to add to the reporting and monitoring burden across the country. This required a systematic indicator evaluation process that could be repeated periodically to inform indicator reporting initiatives across the organization and possibly beyond.

Coincidentally, every five years (1999, 2004, 2009, 2014), CIHI and Statistics Canada invite stakeholders from across the country to a national Consensus Conference on Health Indicators to discuss priority-setting of indicator development and reporting for the next half decade [7,8,9,10]. The latest such conference (held in 2014) provided an opportunity to present the results of the internal evaluation of publicly reported indicators and to validate the results with stakeholders.

This paper describes CIHI’s approach to evaluating of a set of HSP indicators using a systematic criteria-based assessment tool and process. The results of the pilot — including achieving reconfirmation through a national consensus process — and possible next steps for broader implementation of the strategy are also presented in the paper.

EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS**Methods**

This project had four distinct components:

1. Process and criteria development for systematic evaluation of HSP indicators
2. Internal CIHI modified-Delphi sessions
3. Pre-conference survey of stakeholders on indicator recommendations
4. Presentation and ratification of results at the national Consensus Conference

Systematic Evaluation of HSP Indicators

The Institute of Medicine's (IOM's) *Recommendations for Measure Selection Criteria* [11] — Usability, Importance, Scientific Soundness and Feasibility — are consistently used in the evaluation and selection of health measures [12]. While many examples in the literature employ these four domains of criteria, we saw the need to expand the dimensions to include other criteria within three of the four domains. Through a process of research and testing, we arrived at a total of 18 criteria points organized around the four IOM domains (see Table 1) that were feasible to apply and that held meaning to our project objective regarding continued reporting of indicators. There is congruence between these criteria and CIHI's Data Quality Framework [13] dimensions of Accuracy, Timeliness, Comparability, Usability and Relevance. Over a period of two months, 56 of CIHI's suite of HSP indicators were assessed against these 18 criteria. To aid evaluators in their subsequent reviews, we created a one-page summary for each indicator denoting results for each evaluation criterion [14].

Internal CIHI modified-Delphi sessions

Two groups within CIHI participated in the evaluation. First, a technical group of experts (n=6) (comprised of epidemiologists, methodologists and statisticians) independently reviewed each indicator and criterion point, and provided a Likert Scale score between 1 and 9. Likert scores were assessed as follows: 7-9 was considered as robust strength for the indicator and agreement for continued reporting; 4-6 denoted equivocal evidence and further discussion at in-person Delphi session is required; and, 1-3 was considered as weak support for the indicator suggesting it should be retired. Respondents were instructed to produce a Likert score and disposition recommendation based on their assessment of all 18 evaluation criteria as a whole. We therefore forewent weighting evaluation criteria. This allowed for flexibility and context in instances where some criteria proved more informative than others.

Likert scores were averaged and presented at an internal in-person Delphi session as a basis for discussions, but were not automatically tied to a final result of continued indicator reporting. The mean was used to average scores as there were no outlier values across responses. Furthermore, all individual respondent ratings were shown alongside the mean score, thereby illustrating the level of concordance. Beginning with the lowest average scores, each indicator was discussed, pertinent commentary synthesized, and a final consensus reached on a disposition recommendation. Disposition options for indicators were Retain, Recommend Further Research and Development (R&D) or Consultation, or Retire.

Recommendations of the technical group's Delphi sessions were then presented to the CIHI HSP leadership group (n=6) (comprised of senior managers and researchers) who repeated the preceding exercise. First, they were asked to independently review all results to date (including indicator assessments and Likert scores, commentary and disposition recommendations). Results of their individual assessments were collated and presented at an in-person session. Disposition recommendations for each of the 56 indicators were consolidated and finalized based on group consensus. The RAND/UCLA Appropriateness Method [15] guided our internal iterative modified-Delphi sessions.

Pre-conference survey of stakeholders on indicator recommendations

A pre-Consensus Conference survey solicited initial feedback on recommendations. The online survey was available for a period of six weeks prior to the conference. Consensus Conference participants were chosen from an existing list of CIHI partners, stakeholders, and clients; participants were largely hospital/health region CEOs, academics and researchers, representatives from ministries of health, clinicians, and national collaboration partners involved in measuring and monitoring the performance of the health care system. An electronic survey was emailed to conference participants along with background documentation on the evaluation process, methodology, and recommendations. The survey asked respondents whether they Agreed, Disagreed or had No Opinion on the recommendation to retire select HSP indicators as per recommendations from CIHI's internal review.

EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

National Consensus Conference presentation

There were 61 participants at the invitational in-person Consensus Conference held in Toronto on October 16 and 17, 2014. Results of the pre-conference survey were presented. A threshold of 70% agreement by respondents was used to automatically pass recommendations or to otherwise hold further group discussion at the conference. An external moderator facilitated discussion and voting on final indicator dispositions.

For peer review only

EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS**Results****Systematic evaluation of HSP indicators**

The systematic evaluation of HSP indicators was a summative process considering 18 criteria points. Some criteria differentiated indicators more than others. For example, a small number of criteria resulted in mostly uniform findings for the suite of HSP indicators. However, when assessed alongside remaining criteria, important contextual considerations can be gleaned. Notable findings are summarized below by criterion.

Usability

The *Granularity of reporting* criterion identified nuances inherent within public reporting purposes. There are approximately 100 administrative health regions in Canada, and approximately 600 acute care hospitals. Twenty-nine indicators are reported at the regional level, and 27 are reported at the hospital/facility level. All indicators are reported at an aggregate provincial/territorial and national level.

With respect to *Pan-Canadian coverage*, 44 of 56 indicators provided complete pan-Canadian coverage (all provinces and territories). The province of Quebec does not have available or comparable data for a dozen indicators. Similar to the criterion of pan-Canadian coverage, the *Comprehensiveness* criterion assessed the inclusiveness of health services providers that submit data toward the indicator. For example, the mental illness hospitalization indicator includes data on mental health patients treated in general hospitals only, while hospitalizations at free-standing psychiatric institutions are not included due to the differences in data collection.

For the *Usage* criterion, we polled CIHI HSP staff responsible for interacting with clients on indicators and data requests. This provided a proxy for the level and extent of the indicator's usage by clients. The 56 indicators under evaluation were rated as High (n=33), Medium (n=15) or Low usage (n=8).

With regard to *Dimensionality*, breakdowns of indicator results by dimensions of sex and socio-economic status (SES) are available where applicable. Thus, 15 indicators are reportable by SES and 14 are reportable by sex.

In terms of *Timeliness, Reporting frequency and Accessibility*, all 56 indicators were publicly reported annually within 10 months of the relevant data being available for analysis. At the time of the evaluation, all HSP indicators were accessible publicly through online publications such as the Health Indicators e-Publication. Additionally, a majority of facility-level indicators are available to providers through private online tools to allow for more granular breakdowns of results and peer comparative reports.

For the *Trendability* criterion, it was found that time trends vary by indicator. For example, the set of facility-level indicators was largely first reported beginning with 2007 data. Results for select regional indicators dated back to 1997. Overall, regional indicators possessed almost twice as many available years of results compared with facility indicators, a nature of the timing of reporting programs.

Importance

As a proxy measure for *Relevance*, an environmental scan was conducted to understand stakeholder utilization of indicators. A total of 232 instances online were recorded. The top five indicators were Hospital Standardized Mortality Ratio (HSMR) (n=23), 30-Day Overall Readmission (n=18), Wait Times for Hip Fracture Repair (n=17), Ambulatory Care Sensitive Conditions (n=14), and Caesarean Section Rate (n=13).

Detailed statements on the *Actionability* of each indicator were provided to evaluation participants. Specifically, summations on the purpose of indicator, strengths, caveats, and scientific evidence in support were considered.

To measure the degree of *Stakeholder follow-up*, we reviewed all instances of patient-level data requests from providers. In 2013-14, there were 298 requests, with 11 facility-level readmission indicators accounting for 58% of all requests (n=173).

EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS

1 The criterion *Sufficient volumes* quantifies the proportion of indicator results that are suppressed per CIHI's data privacy
2 protocols. In general, indicator results with cell counts less than five are suppressed, and results based on less than 50
3 denominator cases per hospital are flagged as low volume and unstable rates. Facility-level indicators are particularly
4 affected by low volumes and suppressed results: 23 of 27 facility-level indicators had at least one-fifth of all results flagged
5 as low volume. A further 7 of these indicators had at least one-fifth of all results suppressed due to small cell counts. At the
6 extreme, we note the 28-Day Readmission After Stroke and Acute Myocardial Infarction (AMI) indicators with ~75% low
7 volume rates and one-third of all results suppressed.
8
9

10 We performed *Significance of variation* analysis to determine the variability within indicator results. For example, the Hip
11 Fracture Surgical Procedures Performed Within 48 Hours indicators (both Within One and Across facilities) had the lowest
12 relative standard deviation values of 16% and 17%, respectively, indicating minimal differences across indicator results.
13
14

Scientific Soundness

15 The criterion *Data quality* garnered the greatest discussion during Delphi reviews. Limitations of using administrative data
16 were considered. Examples of concern include the inability to assess indications for angiography for AMI patients for the
17 indicator Use of Coronary Angiography Following AMI, and the ability to properly identify denominator cases for the
18 Hysterectomy indicator.
19
20

21 The evaluation revealed that *Validity reviews* were performed for each indicator on an annual basis. These included
22 significance testing of risk factors, monitoring of diagnosis and procedure coding updates, and outlier and significant change
23 analyses. Indicators recommended for further consultation and R&D were identified as such mainly for the purpose of
24 seeking feedback from stakeholders on the validity and clinical relevance of current calculation methodologies.
25
26
27

28 The criterion *Participation bias* assessed whether data submission and participation in the calculation of indicator results
29 were a nature of voluntary participation. All but two indicators — Physician Specialists and General/Family Physicians per
30 100,000 Population — required mandatory participation. In other instances, such as indicators produced for long-term care
31 facilities, participation is not yet mandatory across the country, and therefore results published may contain a participation
32 bias.
33
34

Feasibility

35 *Production cost* was considered based on the extent of staff resources required to produce each indicator. Indicators with
36 complex linkages across multiple databases and those requiring building of episodes of care necessitate a larger degree of
37 resources.
38
39

Modified-Delphi sessions of CIHI technical and leadership groups

40 Mean Likert scores, recommendations and rationale are noted in tables 2, 3 and 4. Nine indicators were recommended as
41 candidates for retirement (Table 3), seven were identified as requiring additional consultation and three were
42 recommended to undergo further redevelopment (Table 2). Thirty-five indicators were recommended for retention (Table
43 4). The rationale to retain these HSP indicators was based on the assessment of all 18 evaluation criteria as a whole.
44 Although retained indicators correlate strongly with high mean Likert scores, this was only one contributor to the
45 recommendation. Ultimately, the discussion during the in-person Delphi sessions allowed for the most pertinent and
46 informative of the 18 evaluation criteria to be considered above others.
47
48
49

50 CIHI leadership and technical groups identified indicators for additional consultation and redevelopment. These indicator
51 recommendations were not forwarded to Consensus Conference participants, but were instead identified for internal R&D
52 efforts in the interim.
53
54

Pre-Consensus Conference survey

55 Forty-eight Consensus Conference participants completed the online survey (response rate of 79%). Eighty-five percent of
56 conference participants had more than 10 years of health care experience. Geographic distribution of respondents
57 correlated well with Canada's population across provinces/territories. Stakeholders from federal and provincial government
58
59
60

EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS

1 agencies accounted for three-quarters of survey respondents, followed by regional health authority executives, hospital
2 administrators and academic/research funding organizations. The mean survey agreement score (as a percentage of
3 responses) for all nine indicators proposed for retirement was 70%, and was used as a benchmark for automatic ratification.
4 The option to select No Opinion for each indicator under survey accounted for an average of 20% of responses (ranging
5 between 12% and 30% across indicators); such an option was made available in the event that respondents held insufficient
6 knowledge on the indicator or did not utilize the indicator within their setting; a response of Agreed, Disagreed or No
7 Opinion was mandatory in the survey.
8
9

National Consensus Conference

10
11 Of the nine indicators recommended for retirement, six received more than 70% agreement as a proportion of responses in
12 the pre-conference survey, and therefore were automatically accepted for retirement (Table 3). The remaining three
13 indicators were discussed as a group, and subsequently also ratified for retirement by conference participants. The majority
14 of indicators recommended for retirement were condition-specific readmission indicators. Ultimately, the decision to retire
15 these indicators was based on appropriateness for continued public reporting. While these indicators were ratified for
16 retirement over concerns of rate stability and small numbers, facilities can continue to calculate and monitor these
17 indicators through CIHI private reporting tools. Consensus on retiring these indicators was achieved with greater ease given
18 that a provider's capacity to continue to privately monitor performance would be maintained.
19
20

21
22 Two contextual health human resources indicators at the regional level — Physician Specialists and General/Family
23 Physicians per 100,000 Population — were also included in the modified Delphi review process, and rated low in Likert Scale
24 scoring (both received a mean score of 3.2). While these indicators provide some context on HSP characteristics, they are
25 already reported elsewhere within CIHI. It was agreed to continue reporting on these indicators but outside of the HSP
26 framework.
27
28

29 Table 4 lists 35 HSP indicators retained for continued public reporting. Although retained indicators correlate strongly with
30 high mean Likert scores, this was only one contributor to the recommendation. For example, the regional level Caesarean
31 Section Rate indicator received a mean Likert score of 4.8 from the technical group, but was retained for public reporting
32 after discussion by the leadership group due to continued concerns over high rates in Canada and therefore a need for
33 continued monitoring.
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

*EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS*Discussion

This exercise proved to be an informative, objective, systematic, transparent, inclusive, and likely repeatable process for evaluating and reconfirming a national set of HSP indicators. Overall, the approach of using 18 sub-criteria was manageable and informative, with feedback from participants that the added information and context made it easier to make a final disposition recommendation for each indicator. The overall timeline of the evaluation process from inception to completion was 18 months. Three distinct phases stand out, each requiring approximately six months to complete: initial R&D of the evaluation plan, executing the evaluation internally at CIHI, and achieving consensus across stakeholders.

An initial Likert score of indicators provides a baseline to proceed with group Delphi reviews. We found it beneficial to begin with the lowest scores and work our way to the highest rated indicators. We also found it operational to have our technical group first review indicators and to pass on recommendations to a leadership group that would consider these in addition to their knowledge and understanding of the use of HSP information in the field. The iterative process of having participants first review indicators independent of other Delphi members and to then convene as a group to discuss findings allowed for a balanced and participatory discussion among participants. These iterative methods ensured a summative process whereby findings were transparent and confirmed at each stage.

The national Consensus Conference provided an opportunity to pilot-test the results of a rigorous, mostly internal methodology for evaluating indicators produced by CIHI. Most recently, CIHI has been incorporating the learnings from this exercise into a broader “lifecycle” approach to indicator development, evaluation and retirement recognizing that all too often there is a tendency to add new indicators to the suite of those reported paying little attention to the utility of those reported in some instances for years. The internally developed evaluation process including the 18 criteria used for assessing previously reported indicators will also lend itself to mid-cycle reviews of suites of indicators that could be modified for such a process. The ability to affirm our internal process with external stakeholders at a national conference provided further confidence in the process. And, while stakeholders appreciated the opportunity to review and ratify our findings, going forward, they expressed comfort with CIHI implementing a systematic evaluation of the indicators and making decisions about reporting. There was congruence in opinion on the suitability of HSP indicators for public reporting throughout the evaluation process, beginning with Likert scores and assessments from CIHI technical staff, to CIHI leadership, and lastly with stakeholders.

Strengths and weaknesses of the study

We recognize that the overall evaluation process required considerable time and resources, there are important benefits to such a comprehensive approach. For example, we ensured a transparent and sequential evaluation, whereby discourse and findings were accumulated and presented in a summarized manner at each phase. We solicited feedback from a wide array of expertise including those responsible for monitoring the results of these indicators on a regular basis. An external moderator facilitating the discussion ensured independence during the consensus process. These processes have been described as favourable conceptual approaches to aid exercises of indicator development, maintenance, and evaluation [14].

One main weakness of this process was the lack of involvement of the “patient/public” voice in evaluating the utility of CIHI’s current suite of publicly reported health system performance indicators. Traditionally the approach to health system performance reporting has largely been targeted to system decision makers. With the growing recognition that health system performance includes measuring things that are important and relevant to the patient/public, it is clear that the patient/public perspective needs to be embedded in future aspects of this work. In 2013, CIHI solicited input from 3,000 Canadians (randomized, representative sample) through small group dialogues and online questionnaires about which types of indicators and domains of health system performance they would like to see publicly-reported. In an attempt to obtain broader input to the evaluation process discussed in this paper, the same survey sent to Consensus Conference participants was made available on CIHI’s website for public participation. The survey responses from the general public were highlighted and considered at the Consensus Conference. However, a more systematic approach to including the patient/public perspective within the “lifecycle” approach to development, evaluation and retirement is needed going forward.

EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS

1 Shekelle [16] notes there is little agreement on methodologies for developing performance indicators, and this can also be
2 said regarding their evaluation. Nonetheless, Stelfox and Straus [14] emphasize the importance of clearly establishing the
3 chosen evaluation criteria in advance of launching a consensus process. In the majority of the studies we reviewed and cite,
4 a smaller number of evaluation criteria were applied: most often, usability, importance, scientific soundness, and feasibility
5 (or a variation thereof that drew upon similar domains). Conversely, we found it helpful to apply multiple sub-criteria to
6 comprehensively reflect the evaluation of indicators for their suitability of ongoing public reporting. Furthermore, providing
7 a more granular evaluation schema for participants ensured more consistent definitions of domains and structured
8 evidence/results for evaluators' consideration. Nonetheless, while these evaluation criteria were informative and applicable
9 to this precise context, not all would apply for other evaluation purposes. Further efforts are necessary to determine the
10 level of customization required to ensure that the process and criteria are applicable to other sectors of care and types of
11 indicators.
12
13

14
15 In addition to convening an in-person consensus conference (or expert panel) to evaluate indicators, Santana et al [17]
16 forwarded their evaluation survey to 101 trauma centres across four countries involved in the use and assessment of injury-
17 care indicators. Moreover, a novel subsequent processes has been described by Bobrovitz et al [18] whereby the discussion
18 occurring throughout the consensus conference is transcribed and undergoes qualitative content analysis to identify key
19 themes raised throughout the deliberations. These additional activities can provide complementary evidence to the
20 evaluation process, such as qualitative findings to an otherwise objective and quantitative exercise, and reaching a broader
21 group of stakeholders and users of health measures.
22
23

24 There are certain characteristics of the Canadian health care system that are favourable for such an evaluation exercise. As
25 the national healthcare system information steward, CIHI receives data for virtually all hospitalizations across the country in
26 a standardized manner. All but two of the 56 HSP indicators are calculated using this standardized data source. Therefore,
27 the application of 18 evaluation criteria to these indicators can be done so in a systematic process so that objectivity is
28 maintained. A centralized healthcare information system is more conducive for cross-country analysis and reporting [19].
29 This also extends to the convening strength of CIHI to bring together stakeholders from all provinces and territories to agree
30 on a national agenda.
31
32

33 To balance the limiting aspects of a Delphi exercise on a set of existing indicators, the Consensus Conference also included
34 working group sessions on identifying priority areas for future indicator development (organized by health system
35 quadrants of Inputs and Characteristics, Outputs, Outcomes and Social Determinants of Health). From these discussions,
36 along with a cross-country consultation process, CIHI has embarked on a path to develop new indicators for the domains of
37 safety (e.g. infections), mental health and addictions (alcohol attributable hospitalizations), and others relating to recently
38 identified priority populations such as seniors and aging (e.g. palliative care), and children and youth.
39
40

Conclusion

41
42 The proliferation of health measures required to fulfill reporting gaps occurred with minimal consideration to alignment and
43 utility with pre-existing indicators. Not surprisingly, then, stakeholders were overwhelmingly in favour of implementing a
44 process that would result in a leaner, more applicable suite of HSP indicators.
45
46

47 CIHI will gradually expand this evaluation methodology to applicable sectors of care. We will also continue to work with
48 external partners to reduce indicator chaos and increase alignment with reporting requirements across the country [6].
49
50

51 This exercise generated identified analytical alignment actions that can be taken at CIHI throughout indicator production
52 and maintenance with a view to reduce indicator chaos. Furthermore, we gained new knowledge about how the HSP
53 indicators we produce are used by stakeholders both through an internet-based environmental scan and via discussions
54 held at the Consensus Conference [10].
55

56 In line with established practices of convening a Consensus Conference every five years, we feel that it is highly beneficial to
57 inform those discussions with a wholesale and systematic criteria-based review of indicators just prior. A broad consultation
58 process encompassing diverse public health stakeholders from across the country helps ensure the development and use of
59
60

EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

indicators most appropriately reflecting the health of populations and the performance of health systems [20]. Similarly, a retrospective exercise on national HSP practices can identify important lessons, of which the selection of indicators suitable for public reporting is an integral component [21].

For peer review only

EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS**Contributors**

OF participated in the design of the evaluation, performed data analysis, facilitated internal Delphi sessions, and drafted the manuscript. KL participated in the design of the evaluation, facilitated internal Delphi sessions, organized the national consensus conference, and drafted the manuscript. YG participated in the design of the evaluation, facilitated internal Delphi sessions, and drafted the manuscript. All authors reviewed and approve the final version.

Funding statement

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests

We have read and understood the BMJ policy on declaration of interests and declare that we have no competing interests.

Data sharing statement

No additional data are available.

Ethics approval and dissemination

Ethics approval was not required as the study did not utilize any sensitive data.

References

1. Klazinga N, Fischer C, ten Asbroek A. Health services research related to performance indicators and benchmarking in Europe. *J Health Serv Res Policy*, 2011;16:38-47. http://hsr.sagepub.com/content/16/suppl_2/38.long (accessed 16 October 2016).
2. Kelley E, Arispe I, Homes J. Beyond the initial indicators: lessons from the OECD Health Care Quality Indicators Project and the US National Healthcare Quality Report. *Int J Qual Health Care*, September 2006:45-51. http://intqhc.oxfordjournals.org/content/18/suppl_1/45.long (accessed 16 October 2016).
3. Carinci F, Van Gool K, Mainz J et al. Towards actionable international comparisons of health system performance: expert revision of the OECD framework and quality indicators. *Int J Qual Health Care*, 2015:1-10. <http://intqhc.oxfordjournals.org/content/27/2/137> (accessed 16 October 2016).
4. Canadian Institute for Health Information. Roadmap Initiative ... Launching the Process. Ottawa:2000. https://www.cihi.ca/en/profile_roadmap_launch_pdf_en.pdf (accessed 16 October 2016).
5. Canadian Institute for Health Information. A Performance Measurement Framework for the Canadian Health System. Ottawa:2013. https://secure.cihi.ca/free_products/HSP-Framework-ENweb.pdf (accessed 16 October 2016).
6. Saskatchewan Health Quality Council. Think big, start small, act now: tackling indicator chaos: a report on a national summit: Saskatoon, May 30-31, 2011. <http://hqc.sk.ca/Portals/0/documents/tracking-indicator-chaos.pdf> (accessed 16 October 2016).
7. Canadian Institute for Health Information. National Consensus conference on Population Health Indicators. Ottawa: 1999. https://secure.cihi.ca/free_products/phi.pdf (accessed 16 October 2016).
8. Canadian Institute for Health Information. The Health Indicators Project: The Next 5 Years. Report from the Second Consensus Conference on Population Health Indicators. Ottawa:2005. https://secure.cihi.ca/free_products/Consensus_Conference_e.pdf (accessed 16 October 2016).
9. Canadian Institute for Health Information. The Health Indicators Project: Report From the Third Consensus Conference on Health Indicators. Ottawa:2009. https://secure.cihi.ca/free_products/82-230-XWE_e.PDF (accessed 16 October 2016).
10. Canadian Institute for Health Information. Rethink, Renew, Retire: Report from the Fourth Consensus Conference on Evaluating Priorities for Canada's Health Indicators. Ottawa: 2015. https://secure.cihi.ca/free_products/Rethink_Renew_Retire.pdf (accessed 16 October 2016).
11. Hurtado M, Swift E, Corrigan J, editors, Committee on the National Quality Report on Health Care Delivery, Board on Health Care Services. Envisioning the National Health Care Quality Report, Washington, DC, 2001, National Academy press. http://www.ncbi.nlm.nih.gov/books/NBK223318/pdf/Bookshelf_NBK223318.pdf (accessed 16 October 2016).
12. Mattke S. When should measures be updated? Development of a conceptual framework for maintenance of quality-of-care measures. *Qual Saf Health Care* 2008;17:182-186. <http://qualitysafety.bmj.com/content/17/3/182.abstract> (accessed 16 October 2016).
13. Canadian Institute for Health Information. The CIHI Data Quality Framework, 2009. Ottawa:2009. https://www.cihi.ca/en/data_quality_framework_2009_en.pdf (accessed 16 October 2016).
14. Stelfox, H & Straus, S. Measuring quality of care: considering conceptual approaches to quality indicator development and evaluation. *J Clinical Epidemiology*. 2013; 66: 1328-1337.
15. Fitch K, Bernstein S, Aguilar M, Burnand B, LaCalle JR, Lazaro P, et al. The RAND/UCLA Appropriateness method user's manual. Santa Monica, CA: RAND Corporation, 2001. http://www.rand.org/pubs/monograph_reports/MR1269 (accessed 16 October 2016).
16. Shekelle P. Quality indicators and performance measures: methods for development need more standardization. *J Clinical Epidemiology*. 2013; 66: 1338-1339.
17. Santana MJ, Stelfox HT, Asbridge M, Ball CG, Cameron P, Dyer D et al. Development and evaluation of evidence-informed quality indicators for adult injury care. *Annals of Surgery*. 2014 Jan;259(1):186-192.
18. Bobrovitz N et al.: A qualitative analysis of a consensus process to develop quality indicators of injury care. *Implementation Science*. 2013 8:45.
19. OECD. Strengthening Health Information Infrastructure for Health Care Quality Governance: Good Practices, New Opportunities and Data Privacy Protection Challenges. Paris: 2013. DOI: <http://dx.doi.org/10.1787/9789264193505->

EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS

- 1 en (accessed 12 December 2016).
2
3 20. Klazinga N, Stronks K, Delnoij D, Verhoeff A. Indicators without a cause. Reflections on the development and use of
4 indicators in health care from a public health perspective. *Int J Qual Health Care*, 2001;13:433-438.
5 <http://intqhc.oxfordjournals.org/content/intqhc/13/6/433.full.pdf> (accessed 16 October 2016).
6 21. van den Berg, M et al.: The Dutch health care performance report: seven years of health care performance
7 assessment in the Netherlands. *Health Research Policy and Systems* 2014 12:1.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 – Evaluation criteria	
Domain/Criterion	Definition
<u>Usability</u>	
· Granularity of reporting	Reporting at national, provincial/territorial, regional and facility levels
· Pan-Canadian coverage	Extent of participation from all provinces and territories
· Comprehensiveness	Proportion of providers submitting data for the indicator
· Usage	Level and extent of usage
· Dimensionality	Ability to break down results by age, sex, socio-economic status, and other dimensions
· Timeliness	Latest year of available results
· Reporting frequency	Whether indicator is reported quarterly, annually or other
· Accessibility	Whether the indicator is publicly and/or privately reported
· Trendability	Number of years of available results for trending
<u>Importance</u>	
· Relevance	Environmental scan identified uses of indicator by stakeholders
· Actionability	Extent to which providers can meaningfully influence the indicator
· Stakeholder follow-up	Number of data and methodological requests within last fiscal year
· Sufficient volumes	Percentage of results suppressed (due to low counts)
· Significance of variation	Degree of variation across reported values
<u>Scientific Soundness</u>	
· Data quality	Strength of data quality, ability to validate results, based on standards
· Validity review	Extent and frequency of reviewing indicator's validity/methodology
· Participation bias	Mandatory or voluntary participation by providers
<u>Feasibility</u>	
· Production cost	Extent of staff/resources to produce indicator

EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS

Table 2 – Indicators identified for additional consultation and further redevelopment

Type	Indicator	Mean Likert Score	Rationale
ADDITIONAL CONSULTATION			
Region	Hip Replacement	5.0	There are concerns of utility and actionability for these indicators as they represent procedure counts per population.
	Knee Replacement	4.8	
	Coronary Artery Bypass Graft (CABG)	6.6	
	Percutaneous Coronary Intervention (PCI)	6.6	
	Cardiac Revascularization	6.6	
Facility	Vaginal Birth After Caesarean Section	4.4	There are concerns of validity and utility for these indicators.
	Birth Trauma	5.4	
FURTHER REDEVELOPMENT			
Region	Hysterectomy	4.4	R&D is required to improve identification of appropriate denominator cases.
Facility	Nursing Sensitive Adverse Events for Medical Patients	6.8	There is an opportunity for incorporation within newly developed Hospital Harm indicator.
	Nursing Sensitive Adverse Events for Surgical Patients	6.8	
<p>Mean Likert Scale Score:</p> <p>7-9 Robust indicator, recommending continued reporting.</p> <p>4-6 Equivocal indicator, further discussion at in-person Delphi session required.</p> <p>1-3 Weak indicator, recommending indicator retirement.</p>			

EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS

Table 3 – Indicators recommended for retirement

Type	Indicator	Mean Likert Score	Rationale	Pre-Consensus Conference Survey Agreement for retirement (as a % of responses)
Facility	28-Day Readmission After Prostatectomy	5.2	These indicators have low volumes of cases leading to unstable rates as well as to the suppression of a large number of results for public reporting. Furthermore, these cases are included in the Surgical/Medical Readmission indicators, and can still be derived through private reporting tools.	82%*
	28-Day Readmission After Hysterectomy	5.6		80%*
	90-Day Readmission After Knee Replacement	6.4		73%*
	90-Day Readmission After Hip Replacement	6.4		72%*
	28-Day Readmission After Stroke	6.2		58%
	Use of Coronary Angiography Following AMI	6.4	Angiography may not be indicated for every AMI patient, depending on his or her clinical history, and the clinical appropriateness of angiography is difficult to ascertain from the administrative hospitalization data. Therefore, it is challenging to interpret and compare the results for this indicator.	78%*
	Hip Fracture Surgical Procedures Performed Within One Facility (48 Hours)	6.4	This indicator does not measure the true proportion of surgeries performed within 48 hours of admission to an acute care hospital, since it does not account for transfers across hospitals. Many patients are transferred from their initial admitting acute care facility to another facility for surgery. The indicator Hip Fracture Surgical Procedures Performed Within 48 Hours, which measures total time across all acute care facilities, will continue to be produced and reported on.	72%*
	28-Day Readmission After AMI	6.4	Concerns have been raised regarding hospitals' ability to take action on this indicator. It is felt that with the regionalization of cardiac care, it is more appropriate to measure readmission after AMI at the regional level (by patient residence) than at the hospital level. In addition, having a low volume of cases leads to unstable rates and to the suppression of a large number of results for public reporting. Therefore, it was proposed to keep the Readmission After AMI indicator at the regional level and to retire the facility-level indicator. Furthermore, readmissions after AMI are included in the 30-Day Overall Readmission indicator at the facility level.	59%
	Primary Caesarean Section Rate	4.6	A new indicator (Low-Risk Caesarean Section) measures the rate of deliveries via Caesarean section among singleton term cephalic pregnancies for women without placenta previa or previous C-section. Since this new indicator is limited to women who have not had a previous C-section, it can take the place of Primary Caesarean Section Rate and be a better indicator of appropriateness.	57%

EVALUATION OF HEALTH SYSTEM PERFORMANCE INDICATORS

Mean Likert Scale Score:
 7-9 Robust indicator, recommending continued reporting.
 4-6 Equivocal indicator, further discussion at in-person Delphi session required.
 1-3 Weak indicator, recommending indicator retirement.
 * Passing the threshold (of 70% agreement among responses) for automatic ratification.

Table 4 – Indicators retained

Type	Indicator	Mean Likert Score
Region	30-Day AMI In-Hospital Mortality	8.8
	30-Day Stroke In-Hospital Mortality	8.8
	Hospital Standardized Mortality Ratio (HSMR)	8.8
	Ambulatory Care Sensitive Conditions	8.6
	Wait Times for Hip Fracture Repair	8.4
	30-Day Readmission for Mental Illness	7.8
	Repeat Hospital Stays for Mental Illness	7.8
	Self-Injury Hospitalization	7.6
	30-Day AMI Readmission	7.4
	Hospitalized Hip Fracture Event	7.2
	Hospitalized Strokes	7.2
	Hospitalized AMI Event	7.0
	Inflow/Outflow Ratio	7.0
	30-Day Readmission: Patients Age 19 and Younger	6.8
	30-Day Obstetric Readmission	6.8
	30-Day Medical Readmission	6.8
	30-Day Surgical Readmission	6.4
	Facility	Mental Illness Patient Days
Mental Illness Hospitalization		6.0
Injury Hospitalization		5.4
Caesarean Section Rate		4.8
30-Day AMI In-Hospital Mortality		8.8
30-Day Stroke In-Hospital Mortality		8.6
Hip Fracture Surgery Within 48 Hours		8.4
30-Day Overall Readmission		8.0
30-Day In-Hospital Mortality Following Major Surgery		8.0
30-Day Readmission: Patients Age 19 and Younger		7.8
30-Day Obstetric Readmission		7.8
30-Day Medical Readmission		7.6
30-Day Surgical Readmission	7.4	
In-Hospital Hip Fracture In Elderly (Age 65+) Patients	7.4	
Obstetric Trauma — Vaginal Delivery With Instrument	7.4	
Obstetric Trauma — Vaginal Delivery Without Instrument	7.4	
Caesarean Section Rate	6.8	
Low-Risk Caesarean Section	6.8	

Mean Likert Scale Score:
 7-9 Robust indicator, recommending continued reporting.
 4-6 Equivocal indicator, further discussion at in-person Delphi session required.
 1-3 Weak indicator, recommending indicator retirement.