S8 Figure:	A random effects meta-analysis of no further pregnancy up to 28 years after cesarean delivery compared wi vaginal delivery
oo rigure.	

	Odds Ratio	Odds Ratio
Study or Subgroup	M-H, Random, 95% CI	M-H, Random, 95% Cl
Elvander 2015	1.80 [1.78, 1.83]	
Fussing-Clausen 2014	1.40 [1.39, 1.41]	
Hall 1989	1.69 [1.54, 1.85]	-
Huttly 1990	1.39 [1.20, 1.62]	
Kjerulff 2013	1.35 [1.30, 1.41]	+
Mollison 2005	1.35 [1.26, 1.46]	-
Murphy 2002	1.96 [1.01, 3.81]	· · · · · · · · · · · · · · · · · · ·
O'Neill 2014	1.74 [1.72, 1.76]	· · · ·
Smith 2006	1.34 [1.29, 1.39]	+
Tollanes 2007a	2.23 [2.15, 2.31]	•
Tollanes 2007b	1.81 [1.77, 1.85]	-
Total (95% CI)	1.60 [1.45, 1.76]	•
Total events		
Heterogeneity: Tau <sup>2</sup> = 0.02; Chi <sup>2</sup> = 1877.79, df = 10 (P < 0.00001); l <sup>2</sup> = 99%		0.2 0.5 1 2 5
Test for overall effect: $Z = 9.56 (P < 0.00001)$		Favors cesarean delivery Favors vaginal delivery