

BMJ Open

BMJ Open is committed to open peer review. As part of this commitment we make the peer review history of every article we publish publicly available.

When an article is published we post the peer reviewers' comments and the authors' responses online. We also post the versions of the paper that were used during peer review. These are the versions that the peer review comments apply to.

The versions of the paper that follow are the versions that were submitted during the peer review process. They are not the versions of record or the final published versions. They should not be cited or distributed as the published version of this manuscript.

BMJ Open is an open access journal and the full, final, typeset and author-corrected version of record of the manuscript is available on our site with no access controls, subscription charges or pay-per-view fees (<http://bmjopen.bmj.com>).

If you have any questions on BMJ Open's open peer review process please email editorial.bmjopen@bmj.com

BMJ Open

Community pharmacists' services for women during pregnancy and breastfeeding in Kuwait: A cross-sectional study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-018980
Article Type:	Research
Date Submitted by the Author:	03-Aug-2017
Complete List of Authors:	Albassam, Abdullah ; Kuwait University - Faculty of Pharmacy, Pharmacy Practice Awad, Abdelmoneim; Kuwait University - Faculty of Pharmacy, Pharmacy Practice
Primary Subject Heading:	Public health
Secondary Subject Heading:	Health services research
Keywords:	Community pharmacists, pregnancy, breastfeeding, self-care, self-medication, Kuwait

SCHOLARONE™
Manuscripts

1 **Title: Community pharmacists' services for women during pregnancy and breastfeeding in**
2 **Kuwait: A cross-sectional study**

3
4 **Authors Information:**

5
6
7 **1. Abdullah Albassam, PhD**

8
9 Assistant Professor
10 Department of Pharmacy Practice, Faculty of Pharmacy, Kuwait University
11 E-mail: albassam@hsc.edu.kw
12

13
14 **2. Abdelmoneim Awad, PhD – Corresponding Author**

15
16 Professor
17 Department of Pharmacy Practice, Faculty of Pharmacy, Kuwait University, Kuwait
18 E-mail: amoneim@hsc.edu.kw
19 Tel: +965-24636914
20 Fax: +965-24636840
21 ORCID:0000-0002-2266-3687
22
23
24
25
26

27 **Sources of support:** This research received no specific grant from any funding agency in the
28 public, commercial or not-for-profit sectors.
29
30

31 **Word count for the paper's text:** 3993

32
33 **Word count for abstract:** 292

34
35 **Number of figures:** 3

36
37 **Number of tables:** 2

38
39 **Conflict of Interest declaration:** We have read and understood the BMJ policy on declaration of
40 interests and declare that we have no competing interests with regard to the data produced
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Objectives: This study was designed to identify the services provided by community pharmacists in Kuwait and their views regarding self-care in pregnancy and lactation. In addition, it determined the pharmacists' recommendations for treatment of pregnancy and breastfeeding related ailments.

Design: Cross-sectional survey.

Setting: Community pharmacies in Kuwait.

Participants: 207 pharmacies were randomly selected from the Ministry of Health database. One registered pharmacist was approached from each pharmacy. 192 (92.8%) pharmacists agreed to participate and completed a self-administered questionnaire.

Outcomes: Services most regularly offered by pharmacists to pregnant and lactating women, pharmacists' recommendations (services) for common and specific ailments during pregnancy and breastfeeding, and pharmacists' views about self-care in pregnancy and breastfeeding.

Results: The top services provided to pregnant and lactating women were recommending vitamins and food supplements (89.8%) and contraception advice (83.4), respectively. More than half of participants indicated that they would recommend medications for headache, constipation, cough, runny nose, sore throat, nausea/vomiting, indigestion, sore or cracked nipple and insufficient milk. Diarrhea, hemorrhoids, insomnia, varicose vein, swelling of the feet and legs, vaginal itching, back pain, fever, mastitis and engorgement were frequently referred to the physician. Recommendations on medication use were occasionally inappropriate in terms of unneeded drug therapy, off-label use, and safety. More than half of pharmacists agreed that they have sufficient knowledge (61.5%; 50.5%) and confidence (58.3%; 53.1%) about offering advice and solving medication and health problems of pregnant and lactating women, respectively. Most of the respondents (88.5%) agreed that a continuing education program on this topic would be of value and priority for their practice.

1 **Conclusion** The present findings show that respondents had different approaches towards
2
3 responding to pregnancy and lactation related ailments; and also highlight the need for multifaceted
4
5 interventions to enhance pharmacists' role in improving maternal health.
6
7

8
9
10 **Keywords:** Community pharmacists, pregnancy, breastfeeding, self-care, self-medication, Kuwait
11
12

13 **Strengths and limitations of this study**

- 14
- 15 ➤ The strength of this survey included the high response rate, which could indicate the importance
16
17 of this topic to community pharmacists and the length of time that they were willing to spend on
18
19 completing the questionnaire.
20
21
 - 22 ➤ Further strength was the proper sample size and sampling method to produce a representative
23
24 data regarding the study population; therefore, the present findings can be generalized at the
25
26 community pharmacists level in Kuwait.
27
28
 - 29 ➤ In addition, this study fills in a gap in the limited existing literature in the developing countries
30
31 and provides useful pieces of information for community pharmacists' services for pregnant and
32
33 lactating women in the Middle Eastern region.
34
35
 - 36 ➤ Limitations of this study include that it is a theoretical survey that did not truly assess real-life
37
38 situations; therefore, the extent of being definitely sure that respondents perform what they
39
40 declare when responding to the questionnaire is not possible and open to recall bias or error.
41
42
 - 43 ➤ Further limitation was the cross-sectional design of the study that represented one point in time;
44
45 hence, did not reflect any alterations in participants' opinions over time regarding responding to
46
47 ailments during pregnancy and lactation.
48
49
50
51
52
53
54
55
56
57
58
59
60

INTRODUCTION

1
2
3 Self-care is defined as “ *the action individuals take for themselves and their families to stay healthy*
4 *and take care of minor and long term conditions, based on their knowledge and the information*
5 *available, and working in collaboration with health and social care professionals where*
6 *necessary*”.¹ Self-medication is a part of self-care behaviors, which is mainly considered in
7
8 developed countries for minor illnesses by using over-the-counter (OTC) medications, while in
9
10 developing countries is for both minor and major illnesses as a wider spectrum of medications is
11
12 available from community pharmacies without a prescription.²
13
14
15
16
17
18
19
20
21

22 The prevalence of self-medication throughout pregnancy was found to be in the range between
23
24 25.1% and 68.3%.³⁻⁷ The most commonly used OTC medications were analgesics, cough and
25
26 common cold remedies, allergy products, laxatives, antacids, vitamins, antibiotics and herbal
27
28 products.^{4 6 8} The rates of self-medication among breastfeeding women ranged between 17% and
29
30 52.4%.^{9 10} The most used OTC medicines were analgesics, antispasmodics, laxatives, and nasal
31
32 decongestants.⁹ The effect of medication use during pregnancy and lactation is a major worry for
33
34 both women and healthcare practitioners.¹¹ Hence, there is a need for professional guidance for
35
36 selection of appropriate and safe OTC medicines for each ailment. In 2011, the International
37
38 Pharmaceutical Federation (FIP) Council approved a document on the valuable pharmacists’ roles
39
40 to improve maternal, newborn, and child health. These roles have been structured in accordance
41
42 with the FIP/WHO (World Health Organization) Guidelines on Good Pharmacy Practice.¹²
43
44
45
46
47
48
49
50

51 Previous studies have evaluated the role of community pharmacists in providing advice or
52
53 counselling regarding pregnancy and lactation related ailments. A study from Israel showed that
54
55 only 9% of community pharmacists reported asking the women to elucidate whether or not they are
56
57 pregnant or lactating.¹³ In Rhode Island, the USA 42% of community pharmacists indicated that
58
59
60

1 they asked women about breastfeeding before providing services.¹⁴ A study in Nebraska, the USA
2
3 reported a variation in community pharmacists' responses to whether they would recommend
4
5 medications for seven common OTC-treatable conditions in pregnancy and breastfeeding. Also,
6
7 some of the respondents recommended unsafe medications.¹⁵ The results of a study conducted in
8
9 France revealed that medications were often recommended by community pharmacists for pain,
10
11 fever, nose and oropharynx disorders, venous insufficiency, dyspepsia and constipation. Overall,
12
13 pharmacists sometimes provided inappropriate advice including medications that were potentially
14
15 harmful in pregnancy.¹⁶ A study that was performed in three countries: the Netherlands, Canada,
16
17 and Iceland showed that most of the pharmacists were unable to provide evidence-based
18
19 recommendation and 90% of them recommended referral to the physician.¹⁷ In Thailand, a study
20
21 reported that about 75% of community pharmacists treated headache, runny nose and sore throat
22
23 with medicines. Over half of the respondents indicated that they were confident and have adequate
24
25 knowledge about providing advice and solving medication and health problems for women during
26
27 pregnancy and breastfeeding.¹⁸ A study conducted in Serbia and Norway reported that several
28
29 respondents' recommendations on medications use were inappropriate.¹⁹
30
31
32
33
34
35
36
37
38

39 Additional studies regarding community pharmacists' services for pregnant and lactating women
40
41 still necessary to be performed, particularly in developing countries where most of medications can
42
43 be obtained from the pharmacy without prescription. To our knowledge, there are no published data
44
45 that evaluate the role of community pharmacists in advising and recommending medications for
46
47 pregnant and lactating women in the Eastern Mediterranean region, including Kuwait. Such study is
48
49 highly warranted since community pharmacists should have a crucial role in sustaining positive
50
51 behaviors in self-care by providing the proper evidence-based information and supporting the public
52
53 to maintain a healthy lifestyle. Hence, this study was designed to identify the services provided by
54
55 community pharmacists regarding self-care in pregnancy and lactation, determine the pharmacists'
56
57
58
59
60

1 recommendations (services) for treatment of pregnancy and lactation related ailments, and identify
2
3 their views about self-care during pregnancy and lactation.
4
5
6
7

8 **METHODS**

9
10 The study design was descriptive and cross-sectional. It was performed in Kuwait, a Middle-
11 Eastern country with an area of 17,820 km² and an approximate population of 3,065,850 individuals
12 (2011 estimate).²⁰ It was conducted during the period from March to December 2015. The study
13 population were employed community pharmacists in Kuwait. Ethical approval was received from
14 the “Ministry of Health Ethical Committee, Kuwait”.
15
16
17
18
19
20
21
22
23

24 PS power and sample size calculator V.3.05 was used to determine the sample size.²¹ One hundred
25 and eight six pharmacists would be needed to determine a 20% difference in proportion between
26 two groups (e.g., male vs. female) with an 80% power and a 5% significance level. Presuming a
27 response rate of 90%, a sample size of 207 community pharmacies were randomly selected from the
28 six governorates using stratified and systematic random sampling.²² Due to the lack of lists with the
29 names and addresses of community pharmacists in Kuwait, lists of community pharmacies at the
30 various governorates were acquired from the Ministry of Health. The lists included a total of 348
31 pharmacies distributed among the six governorates of Kuwait. Only one full-licensed pharmacist
32 was approached from pharmacies hiring more than one pharmacist. The aim of the survey was
33 concisely explained to the pharmacist on duty (face-to face). Pharmacists were free to refuse to take
34 part in the study. Those who agreed to participate in the survey were handed the questionnaires and
35 then were gathered from them anonymously after being completed within one to two weeks. The
36 study participants offered written consent to take part in the survey.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 The study survey was adapted from validated questionnaires that were previously used in Thailand
2 and France.^{16 18} A research group at Kuwait University established the content validity of the
3 adapted survey. Its face validity was assessed with 5 community pharmacists for clarity of
4 questions. Then the survey was pretested on 10 community pharmacists, and refinements were
5 made as needed so that the survey was simple to comprehend and answer.
6
7
8
9
10
11
12
13

14 The pre-tested survey contained four sections. Demographic and other characteristics of
15 respondents were included in the first section (Table 1). Section two contained eleven questions to
16 provide information about the services provided by the community pharmacists regarding self-care
17 in pregnancy and lactation. These questions were about the availability of information leaflet or
18 brochure to promote health for pregnant and breastfeeding women, experience with pregnant and
19 breastfeeding women, number of pregnant and breastfeeding women who visited the pharmacy per
20 week, the three services most commonly provided for pregnant and breastfeeding women, the two
21 symptoms and/or questions that both pregnant and breastfeeding women most frequently consulted
22 the pharmacists in the past, and how do they know that women are pregnant or breastfeeding. Seven
23 of the above questions were close-ended and their options were presented in the results section. The
24 third section included 16 common symptoms in pregnancy and 12 common symptoms in
25 breastfeeding for which pregnant and breastfeeding women often seek advice from pharmacists.
26 They were asked to indicate the recommendations (services) that they will provide for each
27 symptom if being consulted by a pregnant or breastfeeding women. They were needed to select
28 from three options for each ailment as they would be in real life situations: refer to a doctor,
29 dispense medicine, and provide only advice without medicine dispensation. If they decided to
30 dispense medications, they were asked to indicate the names of the medications. The results for this
31 section were presented in three parts: eight common minor symptoms in both pregnant and
32 breastfeeding women, eight specific pregnancy symptoms and four specific breastfeeding symptoms
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 (Figures 1, 2 and 3, respectively). The final section included twelve statements to identify the
2
3 pharmacists' views about self-care in pregnancy and breastfeeding (Table 2). The responses were
4
5 measured using a 5-point Likert scale (strongly disagree, disagree, neither agree nor disagree
6
7 (neutral), agree, and strongly agree). In addition to three questions to determine the pharmacists
8
9 need for continuing education about self-care in pregnancy and breastfeeding, the convenient
10
11 method of delivering the continuing education for them, and the most common source of
12
13 information used by them to prepare themselves for responding to symptoms during pregnancy and
14
15 lactation.
16
17
18
19
20
21

22 Data analysis was conducted using the Statistical Package for Social Sciences (SPSS, version 23,
23
24 SPSS, Chicago, IL, U.S.A.). Pharmacists' responses were presented as percentages (95%
25
26 confidence intervals; CI) and medians (interquartile ranges; IQR). To simplify the results'
27
28 presentation in the text, those who answered "strongly agree" or "agree" were classified as
29
30 "agreed", and those who answered "strongly disagree" or "disagree" as having disagreed. The
31
32 internal consistency for the sections to determine the pharmacists' views about self-care in
33
34 pregnancy and breastfeeding was assessed using Cronbach's α test. The test results were as follows:
35
36 four statements of support self-care, 0.84; two statements about the safety of OTC medicines, 0.91,
37
38 four statements about knowledge and confidence about pregnancy and breastfeeding, 0.79, and two
39
40 statements about undergraduate training in self-care for both pregnant and breastfeeding women,
41
42
43
44
45
46 0.97.
47
48
49
50

51 The univariate logistic regression was used first to evaluate the association of respondents'
52
53 characteristics with the dependent variables. All variables with $p < 0.25$ in the univariate analysis
54
55 were included in the multivariate logistic regression analysis to determine the factors that are
56
57 independently associated with each of the dependent variables. Only the results of multivariate
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

logistic analysis are reported showing odds ratio (OR) and 95% CI. Statistical significance was accepted at $p < 0.05$.

For peer review only

RESULTS

One hundred and ninety-two (92.8%) pharmacists agreed to participate in the study. Their median (IQR) age and experience as practitioners were 35 (11) years and 11 (7) years, respectively. Table (1) shows the respondents' characteristics.

Above two-fifths (n=85; 44.3%; 95% CI: 37.2- 51.6) of participants have information leaflets or brochures to promote health for pregnancy and lactation, and most of these were from pharmaceutical companies (66%). Most of the respondents had experience with pregnant (n =186; 96.9%; 95% CI: 93.0-98.7) and breastfeeding (n=181; 94.3%; 95% CI: 89.7-97.0) women. The median (IQR) numbers of pregnant women and breastfeeding women who visited the community pharmacy per week were 10 (7) and 6 (3), respectively.

Community pharmacists who reported to have experience with pregnant and breastfeeding women were asked to indicate the most frequently provided services. The top three services provided for pregnant women were recommending vitamins and food supplements (n=167; 89.8%; 95% CI: 84.3-93.6), referral to a doctor (n=126; 67.7%; 95% CI: 60.5-74.3), and providing advice about suitable behavior such as lifestyle and exercise (n=115; 61.8%; 95% CI: 54.4-68.8). Other offered services, but to a lesser extent, were diagnosis of symptoms and dispensing of medicines (n=100; 53.8%; 95% CI: 46.3-61.0) and herbal products (n=80; 43.0%; 95% CI: 35.9-50.5). The three services most frequently provided to breastfeeding women were contraception advice (n=151; 83.4%; 95% CI: 77.0-88.4), recommending vitamins and food supplements (n=101; 55.8%; 95% CI: 48.3-63.1), and weight control advice (n=92; 50.8%; 95% CI: 43.3-58.3). Other offered services, but to a lesser extent, were diagnosis of symptoms and dispensing of herbal products (n=88; 48.6%; 95% CI: 41.2-56.1) and medicines (n=69; 38.1%; 95% CI: 31.1- 45.7), and referral to a doctor (n=58; 32.0%; 95% CI: 25.4-39.4). About three-fifths (n=106; 58.6%; 95% CI: 51.0-

1 65.8) of respondents stated that they knew women are pregnant or breastfeeding by asking them,
2
3 while 41.4% (n=75; 95% CI: 34.3-49.0) reported that women inform them before asking about the
4
5 services.
6
7

8
9
10 Moreover, respondents indicated the symptoms and/or questions that most frequently been
11
12 consulted about the past. They were most frequently consulted by pregnant women regarding
13
14 gastrointestinal symptoms (nausea/vomiting, constipation, and stomach cramp), respiratory
15
16 symptoms (common cold and cough), safety of medicine use in pregnancy, and back pain.
17
18 Breastfeeding women most frequently consulted them about a medicine to increase breast milk,
19
20 contraceptive pills, safety of medicine use in breastfeeding, and respiratory symptoms (common
21
22 cold and cough).
23
24
25
26
27

28
29 Figure (1) presents the distribution of pharmacists' responses to the most commonly treated
30
31 symptoms during pregnancy and breastfeeding. Most of the pharmacists recommended medicines or
32
33 referral to a doctor rather than providing advice only without medicine dispensation for treatment of
34
35 these symptoms. More than half of participants recommended medications for treatment of
36
37 headache, constipation, cough, sore throat, and runny nose. In relation to diarrhea and hemorrhoids,
38
39 about three-fifths and half of respondents recommended referral to a doctor rather than dispensing
40
41 medicines or providing only advice, respectively. There were significant associations between the
42
43 recommendation to dispense medicines for treatment of diarrhea or constipation in breastfeeding
44
45 women and the respondents' experience as practitioners ($p < 0.05$). It was found to be more common
46
47 among those with experience of > 10 years compared to those with experience of < 10 years (for
48
49 diarrhea: $p = 0.02$; OR = 2.0; 95% CI: 1.1-3.5) and (for constipation: $p = 0.02$; OR = 2.2; 95% CI: 1.1-
50
51 4.2).
52
53
54
55
56
57
58
59
60

1 Figure (2) presents the distribution of pharmacists' responses to the specific symptoms in
2 pregnancy. Over half of respondents recommended referral of pregnant women to the doctor for
3 swelling of the feet and legs, varicose vein, insomnia, vaginal itching, back pain, and fever and
4 aches. More than two-thirds of participants recommended dispensing of medications for treatment
5 of nausea, vomiting, and indigestion. There was a significant association between the
6 recommendation to dispense medicines for nausea/vomiting and the pharmacists' experience as
7 practitioners ($p < 0.05$). It was found to be more common among those with experience of > 10
8 years compared to those with experience of < 10 years ($p = 0.03$; OR = 2.6; 95% CI: 1.1-6.5). The
9 recommendation to dispense medicines for treatment of vaginal itching was significantly more
10 common among females compared to males ($p = 0.01$; OR = 2.3; 95% CI: 1.2-4.4).
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 Figure (3) presents the distribution of pharmacists' responses to the specific symptoms in
28 breastfeeding. Pharmacists mainly recommended dispensing of medications for sore or cracked
29 nipple and to increase the breast milk, and referral to the doctor for mastitis. There were significant
30 associations between the recommendation to dispense medicine to relieve engorgement or increase
31 the breast milk and gender. It was found to be more prevalent among females compared to males
32 (for engorgement: $p = 0.01$; OR = 2.2; 95% CI: 1.1-4.1) and (for insufficient milk: $p = 0.03$; OR = 2.5;
33 95% CI: 1.1-5.9).
34
35
36
37
38
39
40
41
42
43
44
45

46 The medications that respondents recommended for each of the symptoms during pregnancy and
47 breastfeeding are presented in the supplementary file. Most medicines that were recommended are
48 not detrimental to the mother, fetus and infant. However, the respondents' recommendations on
49 medicine use were sometimes inappropriate. Respondents sometimes recommended ibuprofen for
50 headache, antibiotics for sore throat and productive cough, herbal products for cough, loratadine for
51 runny nose and productive cough, stimulant laxatives for constipation, loperamide for diarrhea, St
52
53
54
55
56
57
58
59
60

1 John's wort and melatonin for insomnia, topical products with no evidence to heal a nipple,
2
3 paracetamol, moisturizing cream and domperidone for engorgement, and herbal products that
4
5 contain fenugreek for insufficient milk.
6

7
8 Table (2) shows the respondents' views regarding self-care of pregnant and breastfeeding women.
9
10 In the first dimension about self-care support, above half of respondents agreed that pharmacists are
11
12 qualified to provide advice and an OTC therapy to treat symptoms in pregnant (58.9%) and
13
14 breastfeeding (65.1%) women. However, less than half of participants agreed that pharmacists
15
16 should recommend an OTC therapy to treat symptoms in pregnant (39.1%) and breastfeeding
17
18 (46.9%) women. In the second dimension about safety, more than two-fifths of participants
19
20 disagreed about the safety of OTC medicines for pregnant (51.6%) and breastfeeding (43.8%)
21
22 women. In the third dimension about knowledge and confidence, more than half of pharmacists
23
24 agreed that they are confident about providing advice and counselling to pregnant (58.3%) and
25
26 breastfeeding (53.1%) women; and that they have adequate knowledge to solve medication and
27
28 health problems of pregnant (61.5%) and breastfeeding (50.5%) women. In the last dimension about
29
30 undergraduate training, four-in ten responders agreed that pharmacy schools provided appropriate
31
32 training to provide advice and an OTC therapy for pregnant (44.3%) and breastfeeding (42.2%)
33
34 women. The agreement that OTC medicines are safe was significantly higher among females
35
36 compared to males ($p=0.04$; OR = 2.1; 95% CI: 1.0-4.3).
37
38
39
40
41
42
43
44
45

46 Most of the respondents ($n=170$; 88.5%; 95% CI: 83.0-92.5) agreed that a continuing education
47
48 program regarding this topic would be of value and priority for their practice. Attending
49
50 lectures/workshops/seminars ($n=80$; 47.1%) was the most convenient method of delivering
51
52 continuing education for them, followed by receiving regular newsletters ($n=61$; 35.9%), and
53
54 receiving distant learning packages ($n=34$; 20%). There was no significant association between the
55
56 respondents' need for continuing education and the independent variables ($p > 0.05$). The most
57
58
59
60

1 commonly indicated sources of information used by respondents to prepare themselves for
2
3 responding to symptoms in pregnancy and breastfeeding was the websites (n=133; 69.2%),
4
5 followed by books (n=99; 51.5 %), and journal articles (n=19; 9.9%).
6
7

8 **DISCUSSION**

9
10 The present results showed that community pharmacists in Kuwait are frequently consulted by
11
12 pregnant and breastfeeding women. Most of the pharmacists recommended medicines or referral to
13
14 a doctor rather than providing advice only for the treatment of pregnancy and breastfeeding related
15
16 ailments. Recommendations on medication use were occasionally inappropriate in terms of
17
18 unneeded drug therapy, off-label use, and safety. Some community pharmacists still lack confidence
19
20 and knowledge to provide advice and resolve health and medication problems of pregnant and
21
22 lactating women. To our knowledge, this is the first study to be performed in Kuwait, and likely in
23
24 the Middle Eastern area, and it contributes to the limited amount of existing literature in the
25
26 developing countries about the services provided by community pharmacists for women during
27
28 pregnancy and lactation. The present results present a baseline quantitative data of these services
29
30 that will aid in the assessment of the current pharmacy practices towards self-care during pregnancy
31
32 and lactation, and offer additional insight in designing future multifaceted interventions to improve
33
34 the community pharmacists' role to deliver the proper advice and resolve the healthcare matters of
35
36 pregnant and lactating women in Kuwait.
37
38
39
40
41
42
43
44
45
46

47 The present findings show that the number of pregnant and lactating women who visited the
48
49 pharmacy per week was 10 and 6, respectively. These results are higher than that reported in the
50
51 USA and Thailand, where pharmacists provided advice for 2.2 to 2.8 pregnant and lactating women
52
53 per week.^{15 18} The current results reveal that the services most regularly offered to pregnant and
54
55 lactating women were recommending vitamin or food supplements, referral to a doctor,
56
57 contraception advice, and weight control advice. These results are close to that indicated in the Thai
58
59
60

1 study, with the exception that symptom diagnosis and medicine dispensing was a commonly
2 provided service in Thailand.¹⁸ The study population indicated that pregnant and lactating women
3 most frequently consulted them regarding gastrointestinal symptoms, respiratory symptoms, and
4 safety of medicine use. It was reported that most people considered these gastrointestinal and
5 respiratory symptoms as minor ailments that can be treated by using OTC medicines that they tend
6 to purchase from a community pharmacy.^{18 29} The finding that respondents were also frequently
7 consulted about the safety of medicines use in pregnancy and lactation is in accordance with a
8 previous report, which has shown that women require information regarding medication use during
9 pregnancy and indicated pharmacists amongst the three mainly utilized sources of information.³⁰
10 These findings indicate that women with pregnancy and lactation related ailments were more
11 tending to visit a pharmacist than a physician. This could be partly explained by the easy access to
12 the community pharmacies, it is evident that community pharmacies are recognized as the utmost
13 reachable healthcare settings due to the high volume of the public that utilize their services.²³
14 Furthermore, it may be due to the observation of the pharmacists by the pregnant and breastfeeding
15 women as the self-care consultant who provide them with enough time to discuss their health
16 problems and prefer to get advice from a pharmacist rather than from a physician when they have
17 non-serious condition.²⁴⁻²⁸ These findings indicate the relevance of maternal-fetal medication as a
18 crucial area for pharmacy practice, which needs the pharmacists to have adequate knowledge and a
19 responsible framework in paying particular attention when these women request advice to improve
20 maternal health. This is confirmed by the finding that 88.5% of respondents agreed that a continuing
21 education regarding self-care for pregnant and breastfeeding women would be of value and priority
22 for their practice.

23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54 Respondents were also asked about the services that they would recommend for pregnancy and
55 lactation related ailments. More than half of pharmacists indicated that they would recommend
56 medications for headache, constipation, cough, runny nose, sore throat, as well as, nausea/vomiting,
57
58
59
60

1 indigestion, sore or cracked nipple and insufficient milk. Diarrhea, hemorrhoids, insomnia, varicose
2
3 vein, swelling of the feet and legs, as well as, vaginal itching/simple discharge, back pain, fever and
4
5 aches, mastitis and engorgement were frequently referred to the physician. In comparison to
6
7 previous reports, a study conducted in France revealed that medications were often recommended
8
9 by pharmacists for pain, fever, nose and oropharynx disorders, venous insufficiency, dyspepsia and
10
11 constipation. They recommended referral to the doctor for nausea/vomiting and back pain.¹⁶ In
12
13 Thailand, about 75% of pharmacists treated headache, runny nose and sore throat with medicines.
14
15 Providing only advice without the dispensation of medicine was mainly recommended for
16
17 constipation, backache, indigestion, varicose vein, insomnia and engorgement. They mainly referred
18
19 women with vaginal itching and simple discharge, and swelling of the feet and legs to the doctor.¹⁸
20
21 A recent study reported that 52% of pharmacists in Serbia recommended medication use for
22
23 treatment of back pain, heavy legs, nausea, common cold and constipation during pregnancy, while
24
25 62% of respondents in Norway recommended non-pharmacological advice as well as referral to a
26
27 doctor.¹⁹ These findings illustrate the large differences in community pharmacists practices between
28
29 countries regarding the services recommended for treatment of pregnancy and lactation related
30
31 ailments. This could be partly explained by the differences in regulatory environments, types of
32
33 undergraduate programs, and the availability of products at the local pharmacies. The finding that
34
35 less than one-tenth of the participants recommended only advice for most of the symptoms
36
37 underscores the need for pharmacists to have sufficient knowledge and information about self-care
38
39 practices which are crucial to alleviate some ailments without medications. The pharmacist must
40
41 have adequate information to reach a conclusion about the risk: benefit ratio of treatment for the
42
43 women to be able to counsel them effectively.
44
45
46
47
48
49
50
51
52
53
54

55 In the present study, recommendations on medication use were occasionally inappropriate in terms
56
57 of unneeded drug therapy, off-label use, and safety. Previous studies showed that community
58
59
60

1 pharmacists were incapable to offer adequate evidence-based information about use of medicines
2
3 during pregnancy.^{15 17} These results demonstrate that respondents have different knowledge levels
4
5 in the subject of maternal-fetal medicine. This is confirmed by the findings that about two-fifths of
6
7 respondents did not agree that they have confidence and aknowledge about giving advice and
8
9 resolving medication and health problems of pregnant and lactating women. In addition, over half of
10
11 participants reported that pharmacy schools did not provide appropriate training regrading advice
12
13 and OTC therapy for pregnant and lactating women. These results underscore the need for
14
15 continuing professional development and the revision of the undergraduate pharmacy curriculum to
16
17 fill the knowledge gaps of pharmacy students and practitioners in maternal-fetal medicine and to
18
19 support pharmacists to deliver the proper care for pregnant and lactating women.
20
21
22
23
24
25

26 The current findings reveal that pharmacists with experience of more than 10 years recommended to
27
28 dispense medicines for diarrhea, constipation, and nausea/vomiting more than those with less
29
30 experience. This might be due to that their knowledge base in the area of pregnancy and
31
32 breastfeeding related ailments relies on experience gained in practice. Also, it was found that female
33
34 respondents recommended to dispense medications for vaginal itching/simple discharge,
35
36 engorgement, and insufficient milk more than males. This could be explained by the fact that female
37
38 pharmacists have more exposure to pregnancy-related issues, either personal or work experience for
39
40 pregnant or lactating women tend to discuss their concerns more comfortable with female than male
41
42 pharmacists. Another possible reason demonstrated by this study was their agreement that OTC
43
44 medicines are safe, which was significantly higher than males. Further qualitative research is
45
46 needed for describing and understanding these predictors.
47
48
49
50
51
52
53
54

55 **Conclusions**

56
57
58
59
60

1 The present results reveal that community pharmacists in Kuwait are frequently consulted by
2 pregnant and breastfeeding women and that the pharmacists had different approaches towards
3 responding to pregnancy and lactation related ailments. Also highlight the need for multifaceted
4 interventions, including continuing professional development and the revision of the undergraduate
5 pharmacy curriculum to fill the knowledge gaps of pharmacy students and practitioners in maternal-
6 fetal medicine and to enhance pharmacists' role in improving maternal health.
7
8
9
10
11
12
13

14 **Acknowledgements**

15 We appreciate the participation of the community pharmacists in this study. We gratefully
16 acknowledge the contribution of pharmacist Moodi Al-Asfour in pretesting the study questionnaire
17 and data collection.
18
19
20
21
22
23
24
25
26
27
28

29 **Authors' contributions** AB contributed in data collection, analysis and interpretation, and wrote
30 the manuscript. AA designed and supervised the study, performed the data analysis and reviewed
31 the manuscript critically for important intellectual content. Both authors read and approved the final
32 manuscript.
33
34
35
36
37
38
39
40

41 **Data sharing statement** The raw data of the present study are available from the corresponding
42 author on reasonable request.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
REFERENCES

1. World self-medication industry (WSMI). WSMI declaration on self-care and self-medication, 2006. <http://www.wsmi.org/wsmi-declaration-on-self-care-and-self-medication-2/>. (accessed 07 Feb 2017).
2. Geissler PW, Nokes K, Prince RJ, *et al*. Children and medicines: Self-treatment of common illnesses among Luo schoolchildren in western Kenya. *Soc Sci Med* 2000;50:1771-83.
3. Buitendijk S, Bracken MB. Medication in early pregnancy: Prevalence of use and relationship to maternal characteristics. *Am J Obstet Gynecol* 1991;165:33-40.
4. Aviv RI, Chubb K, Lindow SW. The prevalence of maternal medication ingestion in the antenatal period. *S Afr Med J* 1993;83:657-60.
5. Abeje G, Admasie C, Wasie B. Factors associated with self medication practice among pregnant mothers attending antenatal care at governmental health centers in Bahir Dar city administration, Northwest Ethiopia, a cross sectional study. *Pan Afr Med J* 2015;20:276.
6. Yusuff KB, Omarusehe LD. Determinants of self medication practices among pregnant women in Ibadan, Nigeria. *Int J Clin Pharm* 2011;33:868-75.
7. Kulkarni PK. IM, Chandrasekhar A Selfmedication practice among urban slum dwellers in south Indian city. *IJPBS* 2012;3:81-7.
8. Westfall RE. Herbal healing in pregnancy: women's experiences. *J Herb Pharmacother* 2003;3:17-39.
9. Chaves RG, Lamounier JA, César CC. Self-medication in nursing mothers and its influence on the duration of breastfeeding. *J Pediatr (Rio J)* 2009;85:129-34.
10. Al-Sawalha NA, Tahaineh L, Sawalha A, *et al*. Medication Use in Breastfeeding Women: A National Study. *Breastfeed Med* 2016;11:386-91.
11. Lagoy CT, Joshi N, Cragan JD, *et al*. Medication use during pregnancy and lactation: an urgent call for public health action. *J Womens Health (Larchmt)* 2005;14:104-9.


- 1 12. International Pharmaceutical Federation (FIP). International Pharmaceutical Federation
2 report on effective use of pharmacists in improving maternal, newborn and child health
3 (MNCH), 2011. http://www.who.int/pmnch/media/news/2012/20120223_fip_paper/en/.
4
5
6 (accessed 11 Feb 2017).
7
8
- 9 13. Merlob P, Stahel B, Kaplan B. Drug use in pregnancy and breast feeding: the role of the
10 pharmacist. *Int J Risk Saf Med* 1998;11:45-7.
11
- 12 14. Ronai C, Taylor JS, Dugan E, *et al*. The identifying and counseling of breastfeeding women
13 by pharmacists. *Breastfeed Med* 2009;4:9-15.
14
- 15 15. Schrempp S, Ryan-Haddad A, Gait KA. Pharmacist counseling of pregnant or lactating
16 women. *J Am Pharm Assoc (Wash)* 2001;41:887-90.
17
- 18 16. Damase-Michel C, Vié C, Lacroix I, *et al*. Drug counselling in pregnancy: an opinion survey
19 of French community pharmacists. *Pharmacoepidemiol Drug Saf* 2004;13:711-5.
20
- 21 17. Lyszkiewicz DA, Gerichhausen S, Björnsdóttir I, *et al*. Evidence based information on drug
22 use during pregnancy: a survey of community pharmacists in three countries. *Pharm World*
23 *Sci* 2001;23:76-81.
24
- 25 18. Sathon B. Self-care in pregnancy and breastfeeding: views of women and community
26 pharmacists in Thailand. . Dissertation Nottingham: University of Nottingham, 2010.
27 <http://eprints.nottingham.ac.uk/11605/>. (accessed 26 Dec 2014).
28
- 29 19. Odalović M, Milanković S, Holst L *et al*. Pharmacists counselling of pregnant women: Web-
30 based, comparative study between Serbia and Norway. *Midwifery* 2016;40:79-86.
31
- 32 20. State of kuwait: General Statistics Bureau. Statistical Review. 39 ed, 2011.
33 https://www.csb.gov.kw/Socan_Statistic.aspx?ID=6. (accessed 02 Mar 2017).
34
- 35 21. Dupont WD, Plummer WD Jr. Power and sample size calculations. version 3.1.2, 2014.
36 <http://biostat.mc.vanderbilt.edu/wiki/Main/PowerSampleSize>. (accessed 12 Jan 2015).
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
22. World Health Organization. How to investigate drug use in health facilities: Selected drug use indicators. Report WHO/DAP/ 93.1, 1993.
 23. Jackson JK SBSM, Duncan GJ, Spinks, JM. Public health - recognising the role of Australian pharmacists. *J Pharm Pract Res* 2004;34:290–2.
 24. Chen J, Britten. Strong medicine': an analysis of pharmacist consultations in primary care. *Fam Pract* 2000;17:480-3.
 25. SA B. Consumer attitudes towards community pharmacy services in Saudi Arabia. *Int J Pharm Pract* 2004;12:83-9.
 26. Wazaify M, Al-Bsoul-Younes A, Abu-Gharbieh E, *et al.* Societal perspectives on the role of community pharmacists and over-the-counter drugs in Jordan. *Pharm World Sci* 2008;30:884-91.
 27. Simoens S, Lobeau M, Verbeke K, *et al.* Patient experiences of over-the-counter medicine purchases in Flemish community pharmacies. *Pharm World Sci* 2009;31:450-7.
 28. Wilbur K, Salam SE, Mohammadi E. Patient perceptions of pharmacist roles in guiding self-medication of over-the-counter therapy in Qatar. *Patient Prefer Adherence* 2010;4:87-93.
 29. Boardman H. Lewis M, Croft P, *et al.* Trinder P, Rajaratnam G. Use of community pharmacies: a population-based survey. *J Public Health (Oxf)* 2005;27:254-62.
 30. Hämeen-Anttila K1, Jyrkkä J, Enlund H *et al.* Medicines information needs during pregnancy: a multinational comparison. *BMJ Open* 2013;3:4.

Table 1 Characteristics of respondents (n=192)


	Frequency	Percentage (%)
Gender		
Male	127	66.1
Female	65	33.9
Age (years)		
20-39	132	68.8
≥ 40	60	31.2
Basic qualification in pharmacy		
B. Pharm	177	92.2
M. Pharm	9	4.7
Pharm D	6	3.1
Postgraduate qualification(s) in pharmacy		
Diploma	15	7.8
Master degree	12	6.2
PhD	3	1.6
Experience as practitioners (Years)		
≤ 10	92	47.9
> 10	100	52.1
Location of pharmacy (Governorates)		
Hawalli	57	29.7
Al-Farwaniyah	43	22.4
Al-Ahmadi/Mubarak Alkabeer	43	22.4
Capital	25	13.0
Al-Jahra	24	12.5

Figure 1 Pharmacists' responses to eight common symptoms in pregnancy and breastfeeding (n=192)




er review only

Figure 2 Pharmacists' responses to specific symptoms in pregnancy (n=192)



review only

Figure 3 Pharmacists' responses to specific symptoms in breastfeeding (n=192)



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49**Table 2 Respondents views about self-care of pregnant and breastfeeding women (n=192)**

	Strongly disagree n (%)	Disagree n (%)	Neutral n (%)	Agree n(%)	Strongly agree n (%)	Median [^] (IQR)
Support self-care						
Community pharmacists are qualified to provide advice and an over- the-counter (OTC) therapy to treat common and minor symptoms in pregnant women	6 (3.1)	29 (15.1)	44 (22.9)	84 (43.8)	29 (15.1)	4.0 (1.0)
Community pharmacists are qualified to provide advice and an OTC therapy to treat common and minor symptoms in breastfeeding women	4 (2.1)	25 (13.0)	38 (19.8)	90 (46.9)	35 (18.2)	4.0 (1.0)
Community pharmacists should recommend OTC therapy and counseling to treat common and minor symptoms in pregnant women	14 (7.3)	49 (25.5)	54 (28.1)	55 (28.7)	20 (10.4)	3.0 (2.0)
Community pharmacists should recommend OTC therapy and counseling to treat common and minor symptoms in breastfeeding women	13 (6.8)	40 (20.8)	49 (25.5)	72 (37.5)	18 (9.4)	3.0 (2.0)
Overall Scale						4.0 (1.0)
Safety of OTC medicine						
OTC medicines are safe for pregnancy	29 (15.1)	70 (36.5)	52 (27.1)	34 (17.7)	7 (3.6)	2.0 (1.0)
OTC medicines are safe for breastfeeding	19 (9.9)	65 (33.9)	56 (29.1)	44 (22.9)	8 (4.2)	3.0 (2.0)
Overall Scale						3.0 (1.0)
Knowledge and confidence about pregnancy and breastfeeding						
I am confident about giving advice and counselling to pregnant women	8 (4.2)	27 (14.1)	45 (23.4)	97 (50.5)	15 (7.8)	4.0 (1.0)
I have sufficient knowledge to solve medication and health problems of pregnant women	7 (3.6)	19 (9.9)	48 (25.0)	95 (49.5)	23 (12.0)	3.0 (2.0)
I am confident about giving advice and counselling to breastfeeding women	11 (5.8)	25 (13.0)	54 (28.1)	83 (43.2)	19 (9.9)	4.0 (1.0)
I have sufficient knowledge to solve medication and health problems of breastfeeding women	14 (7.2)	36 (18.8)	45 (23.4)	75 (39.1)	22 (11.5)	4.0 (1.0)
Overall Scale						4..0 (1.0)
Undergraduate Training						
Pharmacy school provided appropriate training regarding advice and OTC therapy for pregnant women	15 (7.8)	40 (20.8)	52 (27.1)	70 (36.5)	15 (7.8)	3.0 (2.0)
Pharmacy school provided appropriate training regarding advice and OTC therapy for breastfeeding women	16 (8.4)	40 (20.8)	55 (28.6)	65 (33.9)	16 (8.3)	3.0 (2.0)
Overall Scale						3.0 (2.0)

Names of medications recommended by pharmacists for eight common symptoms in pregnancy and breastfeeding

Common symptoms	Frequency (%) of pharmacists recommended medicines for pregnant women	Frequency (%) of pharmacists recommended medicines for breastfeeding women
4 Headache	*n=154	*n=169
5 Paracetamol tablet	151 (98.1%)	145 (85.8)
6 Ibuprofen tablet	3 (1.9%)	24 (14.2)
7 Sore throat**	*n= 106	*n=124
8 Lozenges	90 (84.9)	89 (71.8)
9 Paracetamol tablet	16 (15.1)	29 (23.4)
10 Vitamin C tablet	14 (13.2)	16 (12.9)
11 Amoxicillin capsule	4 (3.8)	6 (4.8)
12 Azithromycin tablet	2 (1.9)	2 (1.6)
13 Runny nose**	*n=113	*n=124
14 Normal saline spray	69 (61.1)	33 (26.6)
15 Loratadine tablet	17 (15.0)	30 (24.2)
16 Paracetamol and Pseudoephedrine tablet	19 (16.8)	22 (17.7)
17 Vitamin C tablet	21 (18.6)	25 (20.2)
18 Vitamin C tablet	7 (6.2)	17 (13.7)
19 Cetirizine tablet	2 (1.8)	3 (1.6)
20 Corticosteroid nasal spray		
21 Dry cough	*n=110	*n=126
22 Prospan [®] syrup	86 (78.2)	75 (59.5)
23 Codilar [®] syrup	12 (10.9)	20 (15.9)
24 Cloperastine syrup	8 (7.3)	22 (17.5)
25 Butamirate syrup	4 (3.6)	9 (7.1)
26		
27 Productive cough**	*n=129	*n=131
28 Prospan [®] syrup	113 (87.6)	85 (64.9)
29 Mucolytic syrup	15 (11.6)	9 (6.9)
30 Amoxicillin capsule	2 (1.6)	2 (1.5)
31 Loratadine tablet	3 (2.3)	24 (18.3)
32 Butamirate syrup	-	10 (7.6)
33 Cloperastine syrup	-	8 (6.1)
34 Constipation	*n=138	*n=134
35 Sterculia granules	63 (45.7)	43 (32.1)
36 Lactulose syrup	60 (43.5)	63 (47.0)
37 Glycerine suppository	6 (4.3)	11 (8.2)
38 Sennalax [®] tablet [®]	9 (6.5)	17 (12.7)
39 Diarrhea**	n=75	n=90
40 Kaolin + Pectin syrup	23 (30.7)	18 (20.0)
41 Nifuroxazide tablet	18 (24.0)	32 (35.6)
42 Loperamide tablet	16 (21.3)	26 (28.9)
43 Oral rehydration salts	25 (33.3)	21 (23.3)
44 Activated charcoal tablet	3 (4.0)	-
45 Hyoscine tablet	-	3 (3.3)
46 Hemorrhoids	n=84	n=95
47 Neo-healar [®] cream	35 (41.6)	25 (26.3)
48 Pilex [®] ointment [®]	24 (28.6)	17 (17.9)
49 Procto-Glyvenol [®] cream	15 (17.9)	24 (25.3)
50 Procto-Glyvenol [®] suppository	10 (11.9)	15 (15.8)
51 Daflon [®] tablet	-	2 (2.1)
52 Proctosedyl [®] suppository	-	12 (12.6)
53 *Number of pharmacists recommended dispensation of a medicine; ** May total > 100% because of recommended combination of medicines; Prospan [®] (herbal medicinal product containing ivy leaf extract); Codilar [®] (Chlorpheniramine + Dextromethorphan + Phenylephrine); Sennalax [®] (Sennosides + docusate); Neo-healar [®] (four medicinal plants: Lupinus Albus, Vateria Indica, Mentha Piperita, Aloe Vera); Pilex [®] (a mixture of herbs [Mimosa pudica, thistles, Vitex negundo, Calendula flowers, camphor], Zinc, and Borax); Procto-Glyvenol [®] (tribenosid + lidocaine); Daflon [®] (diosmin + hesperidin); Proctosedyl [®] (hydrocortisone + cinchocaine).		

Names of medications recommended by respondents for specific symptoms in pregnancy	
Specific pregnancy symptoms	Frequency (%) of pharmacists recommended medicines
Nausea and vomiting	*n=139
Navidoxine® tablet	110 (79.1)
Ondasetron tablet	27 (19.4)
Domperidone tablet	1 (0.7)
Multivitamin capsule	1 (0.7)
Indigestion (heartburn)	*n=131
Gaviscon® suspension	89 (67.9)
Ranitidine tablet	20 (15.3)
Zymogen® tablet	22 (16.8)
Back pain**	*n=92
Paracetamol tablet	42 (45.7)
Diclofenac gel	27 (29.3)
Reparil® gel	7 (7.6)
Relaxnova® cream	21 (22.8)
Insomnia	*n=20
St John's wort®	5 (25.0)
Dream water®	10 (50.0)
Melatonin tablet	5 (25.0)
Fever and aches	*n=93
Paracetamol tablet	93 (100)
Vaginal itching and simple discharge	*n=67
Clotrimazole ovule	20 (29.9)
Clotrimazole cream	15 (22.4)
Clotrimazole wash	9 (13.4)
Miconazole cream	11 (16.4)
Miconazole ovule	12 (17.9)
Varicose veins	*n=28
Venoruton® gel	21 (75.0)
Reparil® gel	7 (25.0)
Swelling feet and legs	*n=15
Reparil® gel	11 (73.3)
Varixinal® gel	4 (26.7)

*Number of pharmacists recommended dispensation of a medicine; ** May total > 100% because of recommended combination of medicines; Zymogen® (Lipase + Proteas +Pepsin + Hemicellulase +Ox bile extract +Dimethylpolysiloxane +Vitamin B1); Reparil® (Aescinum +Diethylamini salicylas); Relaxnova® (Idrocilamide) ; St John's Wort® (Hypericum perforatum); Dream water® (Melatonin + 5-HTP + GABA); Venorutin® (Rutoside); Varixinal® (Ruscus Aculeatus Extract, Aesculus Hippocastanum Extract, Centella Asiatica Extract, Vaccinium Myrtillus Extract).

STROBE Statement—checklist of items that should be included in reports of observational studies

	Item No	Recommendation
Title and abstract	1	(a) Indicate the study's design with a commonly used term in the title or the abstract Pages 1 and 2 (paragraph 2) of the manuscript. (b) Provide in the abstract an informative and balanced summary of what was done and what was found Pages 2 and 3 (paragraph 1) of the manuscript.
Introduction		
Background/rationale	2	Explain the scientific background. Pages 4 and 5 (paragraph 1) of the manuscript. and rationale for the investigation being reported. Page 5 (paragraph 2) of the manuscript.
Objectives	3	State specific objectives, including any prespecified hypotheses. Page 5 (paragraph 2) and page 6 (paragraph 1) of the manuscript.
Methods		
Study design	4	Present key elements of study design early in the paper. Page 6 (paragraph 2) of the manuscript.
Setting	5	Describe the setting, locations, and relevant dates, including periods of recruitment, exposure, follow-up, and data collection. Page 6 (paragraph 2) of the manuscript.
Participants	6	(a) <i>Cohort study</i> —Give the eligibility criteria, and the sources and methods of selection of participants. Describe methods of follow-up <i>Case-control study</i> —Give the eligibility criteria, and the sources and methods of case ascertainment and control selection. Give the rationale for the choice of cases and controls <i>Cross-sectional study</i> —Give the eligibility criteria, and the sources and methods of selection of participants. Page 6 (paragraph 3) of the manuscript. (b) <i>Cohort study</i> —For matched studies, give matching criteria and number of exposed and unexposed <i>Case-control study</i> —For matched studies, give matching criteria and the number of controls per case
Variables	7	Clearly define all outcomes, exposures, predictors, potential confounders, and effect modifiers. Give diagnostic criteria, if applicable. Pages 7 and 8 (paragraph 1) of the manuscript.
Data sources/ measurement	8*	For each variable of interest, give sources of data and details of methods of assessment (measurement). Describe comparability of assessment methods if there is more than one group. Pages 7 and 8 (paragraph 1) of the manuscript.
Bias	9	Describe any efforts to address potential sources of bias. Page 3 of the manuscript.
Study size	10	Explain how the study size was arrived at. Page 6 (paragraph 2) of the manuscript.
Quantitative variables	11	Explain how quantitative variables were handled in the analyses. If applicable, describe which groupings were chosen and why. Page 8 (paragraphs 2 and 3) of the manuscript.
Statistical methods	12	(a) Describe all statistical methods, including those used to control for confounding (b) Describe any methods used to examine subgroups and interactions (c) Explain how missing data were addressed (d) <i>Cohort study</i> —If applicable, explain how loss to follow-up was addressed <i>Case-control study</i> —If applicable, explain how matching of cases and controls was

1
2 addressed

3 *Cross-sectional study*—If applicable, describe analytical methods taking account of
4 sampling strategy. **Page 8 (paragraphs 2 and 3) of the manuscript.**

5 (e) Describe any sensitivity analyses
6

7 Continued on next page
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

Results

Participants	13*	(a) Report numbers of individuals at each stage of study—eg numbers potentially eligible, examined for eligibility, confirmed eligible, included in the study, completing follow-up, and analysed. Page 9 (paragraph 1) of the manuscript and Table 1 (page 21). (b) Give reasons for non-participation at each stage (c) Consider use of a flow diagram
Descriptive data	14*	(a) Give characteristics of study participants (eg demographic, clinical, social) and information on exposures and potential confounders. Page 9 (paragraph 1) of the manuscript and Table 1 (page 21) (b) Indicate number of participants with missing data for each variable of interest There were no missing data for variables of interest since the questionnaires were checked for being completed at the stage of their collection in hand from the study participants. (c) <i>Cohort study</i> —Summarise follow-up time (eg, average and total amount)
Outcome data	15*	<i>Cohort study</i> —Report numbers of outcome events or summary measures over time <i>Case-control study</i> —Report numbers in each exposure category, or summary measures of exposure <i>Cross-sectional study</i> —Report numbers of outcome events or summary measures Pages 9 to 12 of the manuscript + Table 1 (page 21), Figures 1, 2 and 3 (pages 22-24) and table 2 (page 25) + Supplementary file
Main results	16	(a) Give unadjusted estimates and, if applicable, confounder-adjusted estimates and their precision (eg, 95% confidence interval). Make clear which confounders were adjusted for and why they were included (b) Report category boundaries when continuous variables were categorized Pages 9 to 12 of the manuscript (c) If relevant, consider translating estimates of relative risk into absolute risk for a meaningful time period
Other analyses	17	Report other analyses done—eg analyses of subgroups and interactions, and sensitivity analyses Page 8 (paragraph 2) of the manuscript

Discussion

Key results	18	Summarise key results with reference to study objectives Page 13 (paragraph 1) of the manuscript
Limitations	19	Discuss limitations of the study, taking into account sources of potential bias or imprecision. Discuss both direction and magnitude of any potential bias. Page 3 of the manuscript
Interpretation	20	Give a cautious overall interpretation of results considering objectives, limitations, multiplicity of analyses, results from similar studies, and other relevant evidence Pages 13 to 16 (paragraph 2) of the manuscript
Generalisability	21	Discuss the generalisability (external validity) of the study results Page 3 of the manuscript

Other information

Funding	22	Give the source of funding and the role of the funders for the present study and, if applicable, for the original study on which the present article is based Page 1 of the manuscript
---------	----	---

*Give information separately for cases and controls in case-control studies and, if applicable, for exposed and unexposed groups in cohort and cross-sectional studies.

Note: An Explanation and Elaboration article discusses each checklist item and gives methodological background and published examples of transparent reporting. The STROBE checklist is best used in conjunction with this article (freely

1
2 available on the Web sites of PLoS Medicine at <http://www.plosmedicine.org/>, Annals of Internal Medicine at
3 <http://www.annals.org/>, and Epidemiology at <http://www.epidem.com/>). Information on the STROBE Initiative is
4 available at www.strobe-statement.org.
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

BMJ Open

Community pharmacists' services for women during pregnancy and breastfeeding in Kuwait: A cross-sectional study

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2017-018980.R1
Article Type:	Research
Date Submitted by the Author:	26-Sep-2017
Complete List of Authors:	Albassam, Abdullah ; Kuwait University - Faculty of Pharmacy, Pharmacy Practice Awad, Abdelmoneim; Kuwait University - Faculty of Pharmacy, Pharmacy Practice
Primary Subject Heading:	Public health
Secondary Subject Heading:	Health services research
Keywords:	Community pharmacists, pregnancy, breastfeeding, self-care, self-medication, Kuwait

SCHOLARONE™
Manuscripts

1 **Title: Community pharmacists' services for women during pregnancy and breastfeeding in**
2 **Kuwait: A cross-sectional study**

3
4 **Authors Information:**

5
6
7 **1. Abdullah Albassam, PhD**

8
9 Assistant Professor
10 Department of Pharmacy Practice, Faculty of Pharmacy, Kuwait University
11 E-mail: albassam@hsc.edu.kw
12

13
14 **2. Abdelmoneim Awad, PhD – Corresponding Author**

15
16 Professor
17 Department of Pharmacy Practice, Faculty of Pharmacy, Kuwait University, Kuwait
18 E-mail: amoneim@hsc.edu.kw
19 Tel: +965-24636914
20 Fax: +965-24636840
21 ORCID:0000-0002-2266-3687
22
23
24
25
26

27 **Sources of support:** This research received no specific grant from any funding agency in the
28 public, commercial or not-for-profit sectors.
29
30

31 **Word count for the paper's text:** 3594

32
33 **Word count for abstract:** 300

34
35 **Number of figures:** 3

36
37 **Number of tables:** 2

38
39 **Conflict of Interest declaration:** We have read and understood the BMJ policy on declaration of
40 interests and declare that we have no competing interests with regard to the data produced
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT

Objectives: This study was designed to identify the services provided by community pharmacists in Kuwait and their views regarding self-care in pregnancy and lactation. In addition, it determined the pharmacists' recommendations for treatment of pregnancy and breastfeeding related ailments.

Design: Cross-sectional questionnaire-based survey.

Setting: Community pharmacies in Kuwait.

Participants: 207 pharmacies were randomly selected from the Ministry of Health database. One registered pharmacist was approached from each pharmacy. One hundred and ninety-two (92.8%) pharmacists agreed to participate and completed a self-administered questionnaire.

Outcomes: The proportions of pharmacists offering particular advice for health conditions in pregnancy and lactation, pharmacists' recommendations for common and specific ailments during pregnancy and breastfeeding, and pharmacists' views about self-care in pregnancy and breastfeeding.

Results: The top services provided to pregnant and lactating women were recommending vitamins and food supplements (89.8%) and contraception advice (83.4%), respectively. More than half of participants indicated that they would recommend medications for headache, constipation, cough, runny nose, sore throat, nausea/vomiting, indigestion, sore or cracked nipple and insufficient milk. Diarrhea, hemorrhoids, insomnia, varicose vein, swelling of the feet and legs, vaginal itching, back pain, fever, mastitis and engorgement were frequently referred to the physician. Recommendations on medication use were occasionally inappropriate in terms of unneeded drug therapy, off-label use, and safety. In relation to offering advice and solving medication and health problems of pregnant and lactating women, more than half of pharmacists indicated that they have sufficient knowledge (61.5%; 50.5%) and confidence (58.3%; 53.1%), respectively. Most of the respondents (88.5%) agreed that a continuing education program on this topic would be of value for their practice.

1 **Conclusion** The present findings show that respondents had different recommendations for
2
3 treatment of pregnancy and lactation related ailments; and also highlight the need for interventions,
4
5 including continuing professional development and revision of the undergraduate pharmacy
6
7 curriculum.
8
9

10
11
12 **Keywords:** Community pharmacists, pregnancy, breastfeeding, lactation, feto-maternal self-care,
13
14 self-medication, Kuwait
15
16

17 18 19 **Strengths and limitations of this study**

- 20
21
22 ➤ The strength of this survey included the high response rate, which could indicate the importance
23
24 of this topic to community pharmacists and the length of time that they were willing to spend on
25
26 completing the questionnaire.
27
28
29 ➤ Further strength was the adequate sample size and sampling method to produce a representative
30
31 data regarding the study population; therefore, the present findings can be generalized at the
32
33 community pharmacists level in Kuwait.
34
35
36 ➤ In addition, this study fills in a gap in the limited existing literature in the developing countries
37
38 and provides useful pieces of information for community pharmacists' services for pregnant and
39
40 lactating women in the Middle Eastern region.
41
42
43 ➤ Limitations of this study include that the survey did not truly assess real-life situations;
44
45 therefore, the extent of being definitely sure that respondents perform what they declare is not
46
47 possible and open to recall bias or error.
48
49
50 ➤ Further limitation is the social desirability bias that the respondents might have offered favorable
51
52 answers to conform to the more socially accepted view.
53
54
55
56
57
58
59
60

INTRODUCTION

Self-care is defined as “*the action individuals take for themselves and their families to stay healthy and take care of minor and long term conditions, based on their knowledge and the information available, and working in collaboration with health and social care professionals where necessary*”.¹ Self-medication is a part of self-care behaviors, which is mainly considered in developed countries for minor illnesses by using over-the-counter (OTC) medications, while in developing countries is for both minor and major illnesses as a wider spectrum of medications is available from community pharmacies without a prescription.²

The prevalence of self-medication throughout pregnancy was found to be in the range between 25% and 68%.³⁻⁷ The most commonly used OTC medications were analgesics, cough and common cold remedies, allergy products, laxatives, antacids, vitamins, antibiotics and herbal products.^{4 6 8} The rates of self-medication among breastfeeding women ranged between 17% and 52%.^{9 10} The most used OTC medicines were analgesics, antispasmodics, laxatives, and nasal decongestants.⁹ The effect of medication use during pregnancy and lactation is a major worry for both women and healthcare practitioners.¹¹ Hence, there is a need for professional guidance for selection of appropriate and safe OTC medicines for each ailment.

In 2011, the International Pharmaceutical Federation (FIP) Council approved a document on the valuable pharmacists' roles to improve maternal, newborn, and child health. These roles have been structured in accordance with the FIP/WHO (World Health Organization) Guidelines on Good Pharmacy Practice.¹²

1 Previous studies have evaluated the role of community pharmacists in providing advice or
2
3 counselling regarding pregnancy and lactation related ailments. in the USA, France, the
4
5 Netherlands, Canada, Iceland, Serbia, Norway, Thailand, Uganda and Qatar.¹³⁻¹⁸ The main findings
6
7 of these studies included a variation in community pharmacists' responses for treatment of common
8
9 ailments during pregnancy and breastfeeding, the recommendations of unsafe medications during
10
11 pregnancy and lactation, and the different levels of pharmacists' knowledge in the area of maternal-
12
13 fetal medicine.
14
15

16
17
18
19
20 Additional studies regarding community pharmacists' services for pregnant and lactating women
21
22 still need to be performed, particularly in developing countries where most of medications can be
23
24 obtained from the pharmacy without prescription. To our knowledge, only one study has explored
25
26 the pharmacists' knowledge and perceptions of maternal-fetal medicine in the Eastern
27
28 Mediterranean region in Qatar.¹⁸ Hence, this study was designed to identify the services provided
29
30 by community pharmacists regarding self-care in pregnancy and lactation, determine the
31
32 pharmacists' recommendations (services) for treatment of pregnancy and lactation related ailments,
33
34 and identify their views about self-care during pregnancy and lactation. Secondary objectives were
35
36 to determine factors associated with pharmacists' (a) recommendations for treatment of pregnancy
37
38 and lactation related ailments, (b) views about self-care of pregnant and breastfeeding women, and
39
40 (c) need for continuing education.
41
42
43
44
45
46
47

48 **METHODS**

49
50 The study design was a cross-sectional questionnaire-based survey. It was performed in Kuwait, a
51
52 Middle-Eastern country with an area of 17,820 km² and an approximate population of 3,065,850
53
54 individuals (2011 estimate).¹⁹ It was conducted during the period from March to December 2015.
55
56
57
58
59
60

1 The study population comprised of employed community pharmacists in Kuwait. Ethical approval
2
3 (No. 1593) was received from the “Ministry of Health Ethical Committee, Kuwait” on 02/12/2014
4
5 PS power and sample size calculator V.3.05 was used to determine the sample size.²⁰ One hundred
6
7 and eight six pharmacists would be needed to determine a 20% difference in proportion between
8
9 two groups (e.g., male vs. female) with an 80% power and a 5% significance level. Presuming a
10
11 response rate of 90%, a sample size of 207 community pharmacies were randomly selected from the
12
13 six governorates using stratified and systematic random sampling.²¹ Due to the lack of lists with the
14
15 names and addresses of community pharmacists in Kuwait, lists of community pharmacies at the
16
17 various governorates were acquired from the Ministry of Health. The lists included a total of 348
18
19 pharmacies distributed across the six governorates of Kuwait. Only one full-licensed pharmacist
20
21 was approached from pharmacies hiring more than one pharmacist. The aim of the survey was
22
23 concisely explained to the pharmacist on duty (face-to face). Pharmacists were free to refuse to take
24
25 part in the study. Those who agreed to participate in the survey were handed the questionnaires and
26
27 then were gathered from them anonymously after being completed within one to two weeks. They
28
29 signed a consent form to participate in the study.
30
31
32
33
34
35
36
37
38

39 The study survey was adapted from validated questionnaires that were previously used in Thailand
40
41 and France.^{14 16} A research group at Kuwait University established the content validity of the
42
43 adapted survey. Its face validity was assessed with 5 community pharmacists for clarity of
44
45 questions. Then the survey was pretested on 10 community pharmacists, and refinements were
46
47 made as needed so that the survey was simple to comprehend and answer.
48
49
50
51
52

53 The pre-tested survey contained four sections (Appendix-1). Demographic and other characteristics
54
55 of respondents were included in the first section . Section two contained eleven questions to provide
56
57 information about the services provided by the community pharmacists regarding self-care in
58
59
60

1 pregnancy and lactation. These questions were about the availability of information leaflet or
2
3 brochure to promote health for pregnant and breastfeeding women, experience in providing services
4
5 for pregnant and breastfeeding women, number of pregnant and breastfeeding women who visited
6
7 the pharmacy per week, the three services most commonly provided for pregnant and breastfeeding
8
9 women, the two symptoms and/or questions that both pregnant and breastfeeding women most
10
11 frequently consulted the pharmacists in the past, and how do they know that women are pregnant or
12
13 breastfeeding. Seven of the above questions were close-ended. . The third section included 16
14
15 common symptoms in pregnancy and 12 common symptoms in breastfeeding for which pregnant
16
17 and breastfeeding women often seek advice from pharmacists. They were asked to indicate the
18
19 recommendations (services) that they will provide for each symptom if being consulted by a
20
21 pregnant or breastfeeding women. They were needed to select from three options for each ailment
22
23 as they would be in real life situations: refer to a doctor, dispense medicine, and provide only advice
24
25 without dispensing medicine. If they decided to dispense medications, they were asked to indicate
26
27 the names of the medications. . The final section included twelve statements to identify the
28
29 pharmacists' views about self-care in pregnancy and breastfeeding. The responses were measured
30
31 using a 5-point Likert scale (strongly disagree, disagree, neither agree nor disagree (neutral), agree,
32
33 and strongly agree). In addition to three questions to determine the pharmacists need for continuing
34
35 education about self-care in pregnancy and breastfeeding, the convenient method of delivering the
36
37 continuing education for them, and the most common source of information used by them to
38
39 prepare themselves for responding to symptoms during pregnancy and lactation were asked
40
41
42
43
44
45
46
47
48
49

50 Data analysis was conducted using the Statistical Package for Social Sciences (IBM SPSS Statistics
51
52 for Windows, version 23, Armonk, NY: IBM Corp) . Pharmacists' responses were presented as
53
54 percentages (95% confidence intervals; CI) and medians (interquartile ranges; IQR). To simplify the
55
56 results' presentation in the text, those who answered "strongly agree" or "agree" were classified as
57
58
59
60

1 “agreed”, and those who answered “strongly disagree” or “disagree” as having disagreed. The
2
3 internal consistency for the sections to determine the pharmacists’ views about self-care in
4
5 pregnancy and breastfeeding was assessed using Cronbach’s α test.
6
7
8
9

10 The univariate logistic regression was used first to evaluate the association of respondents’
11
12 characteristics with the dependent variables. All variables with $p < 0.25$ in the univariate analysis
13
14 were included in the multivariate logistic regression analysis to determine the factors that are
15
16 independently associated with each of the dependent variables. Only the results of multivariate
17
18 logistic analysis are reported showing odds ratio (OR) and 95% CI. Statistical significance was
19
20 accepted at $p < 0.05$.
21
22
23
24
25
26

27 RESULTS

28
29 One hundred and ninety-two (92.8%) pharmacists agreed to participate in the study. Their median
30
31 (IQR) age and experience as practitioners were 35 (11) years and 11 (7) years, respectively. Table
32
33 (1) shows the respondents’ characteristics.
34
35
36
37
38

39 Above two-fifths ($n=85$; 44.3%; 95% CI: 37.2- 51.6) of participants have information leaflets or
40
41 brochures to promote health for pregnancy and lactation, and most of these were from
42
43 pharmaceutical companies (66%). Most of the respondents had experience in providing services for
44
45 pregnant ($n =186$; 96.9%; 95% CI: 93.0-98.7) and breastfeeding ($n=181$; 94.3%; 95% CI: 89.7-
46
47 97.0) women. The median (IQR) numbers of pregnant women and breastfeeding women who
48
49 visited the community pharmacy per week were 10 (7) and 6 (3), respectively.
50
51
52
53
54

55 Community pharmacists who reported to have experience in providing services for pregnant and
56
57 breastfeeding women were asked to indicate the most frequently provided services. The top three
58
59
60

1 services provided for pregnant women were recommending vitamins and food supplements (n=167;
2 89.8%; 95% CI: 84.3-93.6), referral to a doctor (n=126; 67.7%; 95% CI: 60.5-74.3), and providing
3 advice about suitable behavior such as lifestyle and exercise (n=115; 61.8%; 95% CI: 54.4-68.8).
4
5 Other offered services, but to a lesser extent, were diagnosis of symptoms and dispensing of
6 medicines (n=100; 53.8%; 95% CI: 46.3-61.0) and herbal products (n=80; 43.0%; 95% CI: 35.9-
7 50.5). The three services most frequently provided for breastfeeding women were contraception
8 advice (n=151; 83.4%; 95% CI: 77.0-88.4), recommending vitamins and food supplements (n=101;
9 55.8%; 95% CI: 48.3-63.1), and weight control advice (n=92; 50.8%; 95% CI: 43.3-58.3). The other
10 offered services, but to a lesser extent, were diagnosis of symptoms and dispensing of herbal
11 products (n=88; 48.6%; 95% CI: 41.2-56.1) and medicines (n=69; 38.1%; 95% CI: 31.1- 45.7), and
12 referral to a doctor (n=58; 32.0%; 95% CI: 25.4-39.4). About three-fifths (n=106; 58.6%; 95% CI:
13 51.0-65.8) of respondents stated that they knew women are pregnant or breastfeeding by asking
14 them, while 41.4% (n=75; 95% CI: 34.3-49.0) reported that women inform them before asking
15 about the services.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32

33
34
35
36 Moreover, respondents indicated that they were most often consulted by pregnant women regarding
37 gastrointestinal symptoms (nausea/vomiting, constipation, and stomach cramp), respiratory
38 symptoms (common cold and cough), safety of medication use , and back pain. Breastfeeding
39 women most frequently consulted them about a medicine to increase breast milk, contraceptive
40 pills, safety of medication use , and respiratory symptoms (common cold and cough).
41
42
43
44
45
46
47
48
49

50 Figure (1) presents the distribution of pharmacists' responses to the most commonly treated
51 symptoms during pregnancy and breastfeeding. Most of the pharmacists recommended medicines or
52 referral to a doctor rather than providing advice only without dispensing medicine for treatment of
53 these symptoms. More than half of participants recommended medications for treatment of
54
55
56
57
58
59
60

1 headache, constipation, cough, sore throat, and runny nose. In relation to diarrhea and hemorrhoids,
2
3 about three-fifths and half of respondents recommended referral to a doctor rather than dispensing
4
5 medicines or providing only advice, respectively. There were significant associations between the
6
7 recommendation to dispense medicines for treatment of diarrhea or constipation in breastfeeding
8
9 women and the respondents' experience as practitioners ($p < 0.05$). It was found to be more common
10
11 among those with experience of > 10 years compared to those with experience of < 10 years (for
12
13 diarrhea: $p = 0.02$; OR = 2.0; 95% CI: 1.1-3.5) and (for constipation: $p = 0.02$; OR = 2.2; 95% CI: 1.1-
14
15 4.2).
16
17
18
19

20
21
22 Figure (2) presents the distribution of pharmacists' responses to the specific symptoms in
23
24 pregnancy. Over half of respondents recommended referral of pregnant women to the doctor for
25
26 swelling of the feet and legs, varicose vein, insomnia, vaginal itching, back pain, and fever and
27
28 aches. More than two-thirds of participants recommended dispensing of medications for treatment
29
30 of nausea, vomiting, and indigestion. There was a significant association between the
31
32 recommendation to dispense medicines for nausea/vomiting and the pharmacists' experience as
33
34 practitioners ($p < 0.05$). It was found to be more common among those with experience of > 10
35
36 years compared to those with experience of < 10 years ($p = 0.03$; OR = 2.6; 95% CI: 1.1-6.5). The
37
38 recommendation to dispense medicines for treatment of vaginal itching was significantly more
39
40 common among females compared to males ($p = 0.01$; OR = 2.3; 95% CI: 1.2-4.4).
41
42
43
44
45
46
47

48
49 Figure (3) presents the distribution of pharmacists' responses to the specific symptoms in
50
51 breastfeeding women. Pharmacists mainly recommended dispensing of medications for sore or
52
53 cracked nipple and to increase the breast milk, and referral to the doctor for mastitis. There were
54
55 significant associations between the recommendation to dispense medicine to relieve engorgement
56
57 or increase the breast milk and gender. It was found to be more prevalent among females compared
58
59
60

1 to males (for engorgement: $p=0.01$; OR = 2.2; 95% CI: 1.1-4.1) and (for insufficient milk: $p=0.03$;
2
3 OR = 2.5; 95% CI: 1.1-5.9).
4
5
6
7

8 The medications that respondents recommended for each of the symptoms during pregnancy and
9 breastfeeding are presented in the supplementary file (Appendix 2). Most medicines that were
10 recommended are not detrimental to the mother, fetus and infant. However, the respondents'
11 recommendations on medicine use were sometimes inappropriate in terms of unneeded drug
12 therapy, off-label use, and safety. Respondents sometimes recommended ibuprofen for headache,
13 antibiotics for sore throat and productive cough, herbal products for cough, loratadine for runny
14 nose and productive cough, stimulant laxatives for constipation, loperamide for diarrhea, St John's
15 wort and melatonin for insomnia, topical products with no evidence to heal a nipple, paracetamol,
16 moisturizing cream and domperidone for engorgement, and herbal products that contain fenugreek
17 for insufficient milk.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33

34 The Cronbach's α test results for the sections to determine the pharmacists' views about self-care
35 in pregnancy and breastfeeding were as follows: four statements of support self-care, 0.84; two
36 statements about the safety of OTC medicines, 0.91, four statements about knowledge and
37 confidence about pregnancy and breastfeeding, 0.79, and two statements about undergraduate
38 training in self-care for both pregnant and breastfeeding women, 0.97. Table (2) shows the
39 respondents' views regarding self-care of pregnant and breastfeeding women. In the first dimension
40 about self-care support, above half of respondents agreed that pharmacists are qualified to provide
41 advice and an OTC therapy to treat symptoms in pregnant (58.9%) and breastfeeding (65.1%)
42 women. However, less than half of participants agreed that pharmacists should recommend an OTC
43 therapy to treat symptoms in pregnant (39.1%) and breastfeeding (46.9%) women. In the second
44 dimension, s, more than two-fifths of participants disagreed about the safety of OTC medicines for
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 pregnant (51.6%) and breastfeeding (43.8%) women. In the third dimension about knowledge and
2 confidence, more than half of pharmacists agreed that they are confident about providing advice and
3 counselling to pregnant (58.3%) and breastfeeding (53.1%) women; and that they have adequate
4 knowledge to solve medication and health problems of pregnant (61.5%) and breastfeeding (50.5%)
5 women. In the last dimension about undergraduate training, four-in ten responders agreed that
6 pharmacy schools provided appropriate training to provide advice and an OTC therapy for pregnant
7 (44.3%) and breastfeeding (42.2%) women. The agreement that OTC medicines are safe was
8 significantly higher among females compared to males ($p=0.04$; OR = 2.1; 95% CI: 1.1-4.3).
9
10
11
12
13
14
15
16
17
18
19
20
21

22 Most of the respondents ($n=170$; 88.5%; 95% CI: 83.0-92.5) agreed that a continuing education
23 program regarding this topic would be of value and priority for their practice. Attending
24 lectures/workshops/seminars ($n=80$; 47.1%) was the most convenient method of delivering
25 continuing education for them, followed by receiving regular newsletters ($n=61$; 35.9%), and
26 receiving distant learning packages ($n=34$; 20%). There was no significant association between the
27 respondents' need for continuing education and the independent variables (age: $p=0.44$; gender:
28 $p=0.47$; experience: $p=0.49$). The most commonly indicated sources of information used by
29 respondents to prepare themselves for responding to symptoms in pregnancy and breastfeeding was
30 the websites ($n=133$; 69.2%), followed by books ($n=99$; 51.5 %), and journal articles ($n=19$; 9.9%).
31
32
33
34
35
36
37
38
39
40
41
42
43

44 **DISCUSSION**

45
46 The present results showed that community pharmacists in Kuwait are frequently consulted by
47 pregnant and breastfeeding women. Most of the pharmacists recommended medicines or referral to
48 a doctor rather than providing advice only for the treatment of pregnancy and breastfeeding related
49 ailments. Recommendations on medication use were occasionally inappropriate in terms of
50 unneeded drug therapy, off-label use, and safety. Some community pharmacists still lack confidence
51 and knowledge to provide advice and resolve health and medication problems of pregnant and
52
53
54
55
56
57
58
59
60

1 lactating women. To our knowledge, this is the first study to be performed in Kuwait, and the
2
3 second in the Middle Eastern area, and it contributes to the limited amount of existing literature in
4
5 the developing countries about the services provided by community pharmacists for women during
6
7 pregnancy and lactation. The present results present a baseline quantitative data of these services
8
9 that will aid in the assessment of the current pharmacy practices towards self-care during pregnancy
10
11 and lactation, and offer additional insight in designing future multifaceted interventions to improve
12
13 the community pharmacists' role to deliver the proper advice and resolve the healthcare matters of
14
15 pregnant and lactating women in Kuwait.
16
17
18
19
20
21

22 The current results reveal that the services most regularly offered to pregnant and lactating women
23
24 were recommending vitamin or food supplements, referral to a doctor, contraception advice, and
25
26 weight control advice. These results are close to those indicated in the Thai study, with the
27
28 exception that symptom diagnosis and medicine dispensing was a commonly provided service in
29
30 Thailand.¹⁶ The study participants indicated that pregnant and lactating women most frequently
31
32 consulted them regarding gastrointestinal symptoms, respiratory symptoms, and safety of medicine
33
34 use. It was reported that most people considered these gastrointestinal and respiratory symptoms as
35
36 minor ailments that can be treated by using OTC medicines that they tend to purchase from a
37
38 community pharmacy.^{16 22} The finding that respondents were also frequently consulted about the
39
40 safety of medicines use in pregnancy and lactation is in accordance with a previous report, which
41
42 has shown that women require information regarding medication use during pregnancy and
43
44 indicated pharmacists amongst the three mainly utilized sources of information.²³ These findings
45
46 indicate that women with pregnancy and lactation related ailments were more tending to visit a
47
48 pharmacist than a physician. This could be partly explained by the easy access to the community
49
50 pharmacies, it is evident that community pharmacies are recognized as the utmost reachable
51
52 healthcare settings due to the high volume of the public that utilize their services.²⁴ Furthermore, it
53
54
55
56
57
58
59
60

1 may be due to the observation of the pharmacists by the pregnant and breastfeeding women as the
2 self-care consultant who provide them with enough time to discuss their health problems and prefer
3 to get advice from a pharmacist rather than from a physician when they have non-serious
4 condition.²⁵⁻²⁹ These findings indicate the relevance of maternal-fetal medication as a crucial area
5 for pharmacy practice, which needs the pharmacists to have adequate knowledge and a responsible
6 framework in paying particular attention when these women request advice to improve maternal
7 health.
8
9

10 Respondents were also asked about the services that they would recommend for pregnancy and
11 lactation related ailments. More than half of pharmacists indicated that they would recommend
12 medications for headache, constipation, cough, runny nose, sore throat, as well as, nausea/vomiting,
13 indigestion, sore or cracked nipple and insufficient milk. In comparison to previous reports, a study
14 conducted in France revealed that medications were often recommended by pharmacists for pain,
15 fever, nose and oropharynx disorders, venous insufficiency, dyspepsia and constipation.¹⁴ In
16 Thailand, about 75% of pharmacists treated headache, runny nose and sore throat with medicines. .
17 A recent study reported that 52% of pharmacists in Serbia recommended medication use for
18 treatment of back pain, heavy legs, nausea, common cold and constipation during pregnancy, while
19 62% of respondents in Norway recommended non-pharmacological advice as well as referral to a
20 doctor.¹⁷ These findings illustrate the large differences in community pharmacists practices between
21 countries regarding the services recommended for treatment of pregnancy and lactation related
22 ailments. This could be partly explained by the differences in regulatory environments, types of
23 undergraduate programs, and the availability of products at the local pharmacies. The finding that
24 less than one-tenth of the participants recommended only advice for most of the symptoms
25 underscores the need for pharmacists to have sufficient knowledge and information about self-care
26 practices which are crucial to alleviate some ailments without medications. The pharmacist must
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 have adequate information to reach a conclusion about the risk: benefit ratio of treatment for the
2
3 women to be able to counsel them effectively.
4
5
6
7

8 In the present study, recommendations on medication use were occasionally inappropriate in terms
9
10 of unneeded drug therapy, off-label use, and safety. Previous studies showed that community
11
12 pharmacists were incapable to offer adequate evidence-based information about use of medicines
13
14 during pregnancy.^{13 15} These results demonstrate that respondents have different knowledge levels
15
16 in the subject of maternal-fetal medicine. and underscore the need for continuing professional
17
18 development and the revision of the undergraduate pharmacy curriculum to fill the knowledge gaps
19
20 of pharmacy students and practitioners in maternal-fetal medicine and to support pharmacists to
21
22 deliver the proper care for pregnant and lactating women.
23
24
25
26
27

28 The current findings reveal that pharmacists with experience of more than 10 years recommended to
29
30 dispense medicines for diarrhea, constipation, and nausea/vomiting more than those with less
31
32 experience. This might be due to that their knowledge base in the area of pregnancy and
33
34 breastfeeding related ailments relies on experience gained in practice. Also, it was found that female
35
36 respondents recommended to dispense medications for vaginal itching/simple discharge,
37
38 engorgement, and insufficient milk more than males. This could be explained by the fact that female
39
40 pharmacists have more exposure to pregnancy-related issues, either personal or work experience for
41
42 pregnant or lactating women tend to discuss their concerns more comfortable with female than male
43
44 pharmacists. Another possible reason demonstrated by this study was their agreement that OTC
45
46 medicines are safe, which was significantly higher than males. Further qualitative research is
47
48 needed for describing and understanding these predictors.
49
50
51
52
53
54
55
56

57 **Conclusions**

58
59
60

1 The present results reveal that community pharmacists in Kuwait are frequently consulted by
2
3 pregnant and breastfeeding women and that the pharmacists had different approaches towards
4
5 responding to pregnancy and lactation related ailments. Also highlight the need for multifaceted
6
7 interventions, including continuing professional development and the revision of the undergraduate
8
9 pharmacy curriculum to fill the knowledge gaps of pharmacy students and practitioners in maternal-
10
11 fetal medicine and to enhance pharmacists' role in improving maternal health.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

29 **Acknowledgements**

30
31 We appreciate the participation of the community pharmacists in this study. We gratefully
32
33 acknowledge the contribution of pharmacist Moodi Al-Asfour in pretesting the study questionnaire
34
35 and data collection.
36
37
38
39

40
41 **Authors' contributions** AB contributed in data collection, analysis and interpretation, and wrote
42
43 the manuscript. AA designed and supervised the study, performed the data analysis and reviewed
44
45 the manuscript critically for important intellectual content. Both authors read and approved the final
46
47 manuscript.
48
49

50
51
52
53 **Data sharing statement** The raw data of the present study are available from the corresponding
54
55 author on reasonable request.
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For peer review only

REFERENCES

1. World self-medication industry (WSMI). WSMI declaration on self-care and self-medication, 2006. <http://www.wsmi.org/wsmi-declaration-on-self-care-and-self-medication-2/>. (accessed 07 Feb 2017).
2. Geissler PW, Nokes K, Prince RJ, *et al*. Children and medicines: Self-treatment of common illnesses among Luo schoolchildren in western Kenya. *Soc Sci Med* 2000;50:1771-83.
3. Buitendijk S, Bracken MB. Medication in early pregnancy: Prevalence of use and relationship to maternal characteristics. *Am J Obstet Gynecol* 1991;165:33-40.
4. Aviv RI, Chubb K, Lindow SW. The prevalence of maternal medication ingestion in the antenatal period. *S Afr Med J* 1993;83:657-60.
5. Abeje G, Admasie C, Wasie B. Factors associated with self medication practice among pregnant mothers attending antenatal care at governmental health centers in Bahir Dar city administration, Northwest Ethiopia, a cross sectional study. *Pan Afr Med J* 2015;20:276.
6. Yusuff KB, Omarusehe LD. Determinants of self medication practices among pregnant women in Ibadan, Nigeria. *Int J Clin Pharm* 2011;33:868-75.
7. Kulkarni PK. IM, Chandrasekhar A Selfmedication practice among urban slum dwellers in south Indian city. *IJPBS* 2012;3:81-7.
8. Westfall RE. Herbal healing in pregnancy: women's experiences. *J Herb Pharmacother* 2003;3:17-39.
9. Chaves RG, Lamounier JA, César CC. Self-medication in nursing mothers and its influence on the duration of breastfeeding. *J Pediatr (Rio J)* 2009;85:129-34.
10. Al-Sawalha NA, Tahaine L, Sawalha A, *et al*. Medication Use in Breastfeeding Women: A National Study. *Breastfeed Med* 2016;11:386-91.
11. Lagoy CT, Joshi N, Cragan JD, *et al*. Medication use during pregnancy and lactation: an urgent call for public health action. *J Womens Health (Larchmt)* 2005;14:104-9.

- 1 12. International Pharmaceutical Federation (FIP). International Pharmaceutical Federation
2 report on effective use of pharmacists in improving maternal, newborn and child health
3 (MNCH), 2011. http://www.who.int/pmnch/media/news/2012/20120223_fip_paper/en/.
4
5
6
7 (accessed 11 Feb 2017).
8
- 9
10 13. Schrempp S, Ryan-Haddad A, Gait KA. Pharmacist counseling of pregnant or lactating
11 women. *J Am Pharm Assoc (Wash)* 2001;41:887-90.
12
- 13 14. Damase-Michel C, Vié C, Lacroix I, *et al.* Drug counselling in pregnancy: an opinion survey
14 of French community pharmacists. *Pharmacoepidemiol Drug Saf* 2004;13:711-5.
15
- 16 15. Lyszkiewicz DA, Gerichhausen S, Björnsdóttir I, *et al.* Evidence based information on drug
17 use during pregnancy: a survey of community pharmacists in three countries. *Pharm World*
18
19
20
21
22
23
24
25
26
27 16. Sathon B. Self-care in pregnancy and breastfeeding: views of women and community
28 pharmacists in Thailand. . Dissertation Nottingham: University of Nottingham, 2010.
29
30
31
32
33 <http://eprints.nottingham.ac.uk/11605/>. (accessed 26 Dec 2014).
34
- 35 17. Odalović M, Milanković S, Holst L *et al.* Pharmacists counselling of pregnant women: Web-
36 based, comparative study between Serbia and Norway. *Midwifery* 2016;40:79-86.
37
- 38 18. Bains S, Kitutu FE, Rahhal A *et al.* Comparison of pharmacist knowledge, perceptions and
39 training opportunities regarding maternal/feral medicine in Canada, Qatar, and Uganda. *Can*
40
41
42
43
44
45
46 19. State of kuwait: General Statistics Bureau. Statistical Review. 39 ed, 2011.
47
48
49
50
51
52
53
54
55
56
57
58
59
60
20. Dupont WD, Plummer WD Jr. Power and sample size calculations. version 3.1.2, 2014.
<http://biostat.mc.vanderbilt.edu/wiki/Main/PowerSampleSize>. (accessed 12 Jan 2015).
21. World Health Organization. How to investigate drug use in health facilities: Selected drug
use indicators. Report WHO/DAP/ 93.1, 1993.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
22. Boardman H, Lewis M, Croft P, *et al.* Trinder P, Rajaratnam G. Use of community pharmacies: a population-based survey. *J Public Health (Oxf)* 2005;27:254-62.
23. Hämeen-Anttila K1, Jyrkkä J, Enlund H *et al.* Medicines information needs during pregnancy: a multinational comparison. *BMJ Open* 2013;3(4) pii: e002594.
24. Jackson JK SBSM, Duncan GJ, Spinks, JM. Public health - recognising the role of Australian pharmacists. *J Pharm Pract Res* 2004;34:290–2.
25. Chen J, Britten. Strong medicine': an analysis of pharmacist consultations in primary care. *Fam Pract* 2000;17:480-3.
26. Bawazir SA . Consumer attitudes towards community pharmacy services in Saudi Arabia. *Int J Pharm Pract* 2004;12:83-9.
27. Wazaify M, Al-Bsoul-Younes A, Abu-Gharbieh E, *et al.* Societal perspectives on the role of community pharmacists and over-the-counter drugs in Jordan. *Pharm World Sci* 2008;30:884-91.
28. Simoens S, Lobeau M, Verbeke K, *et al.* Patient experiences of over-the-counter medicine purchases in Flemish community pharmacies. *Pharm World Sci* 2009;31:450-7.
- 29 . Wilbur K, Salam SE, Mohammadi E. Patient perceptions of pharmacist roles in guiding self-medication of over-the-counter therapy in Qatar. *Patient Prefer Adherence* 2010;4:87-93.

Table 1 Characteristics of respondents (n=192)

	Frequency	Percentage (%)
Gender		
Male	127	66.1
Female	65	33.9
Age (years)		
20-29	28	14.6
30-39	104	54.2
≥ 40	60	31.2
Basic qualification in pharmacy		
B. Pharm	177	92.2
M. Pharm	9	4.7
*Pharm D	6	3.1
Postgraduate qualification(s) in pharmacy		
Diploma	15	7.8
Master degree	12	6.2
PhD	3	1.6
Experience as practitioners (Years)		
≤ 10	92	47.9
> 10	100	52.1
Location of pharmacy (Governorates)		
Hawalli	57	29.7
Al-Farwaniyah	43	22.4
Al-Ahmadi/Mubarak Alkabeer	43	22.4
Capital	25	13.0
Al-Jahra	24	12.5
*Doctor of Pharmacy degree		

Table 2 Respondents views about self-care of pregnant and breastfeeding women (n=192)

	Strongly disagree n (%)	Disagree n (%)	Neutral n (%)	Agree n(%)	Strongly agree n (%)	Median [^] (IQR)
Support self-care						
Community pharmacists are qualified to provide advice and an over-the-counter (OTC) therapy to treat common and minor symptoms in pregnant women	6 (3.1)	29 (15.1)	44 (22.9)	84 (43.8)	29 (15.1)	4.0 (1.0)
Community pharmacists are qualified to provide advice and an OTC therapy to treat common and minor symptoms in breastfeeding women	4 (2.1)	25 (13.0)	38 (19.8)	90 (46.9)	35 (18.2)	4.0 (1.0)
Community pharmacists should recommend OTC therapy and counseling to treat common and minor symptoms in pregnant women	14 (7.3)	49 (25.5)	54 (28.1)	55 (28.7)	20 (10.4)	3.0 (2.0)
Community pharmacists should recommend OTC therapy and counseling to treat common and minor symptoms in breastfeeding women	13 (6.8)	40 (20.8)	49 (25.5)	72 (37.5)	18 (9.4)	3.0 (2.0)
Overall Scale						4.0 (1.0)
Safety of OTC medicine						
OTC medicines are safe for pregnancy	29 (15.1)	70 (36.5)	52 (27.1)	34 (17.7)	7 (3.6)	2.0 (1.0)
OTC medicines are safe for breastfeeding	19 (9.9)	65 (33.9)	56 (29.1)	44 (22.9)	8 (4.2)	3.0 (2.0)
Overall Scale						3.0 (1.0)
Knowledge and confidence about pregnancy and breastfeeding						
I am confident about giving advice and counselling to pregnant women	8 (4.2)	27 (14.1)	45 (23.4)	97 (50.5)	15 (7.8)	4.0 (1.0)
I have sufficient knowledge to solve medication and health problems of pregnant women	7 (3.6)	19 (9.9)	48 (25.0)	95 (49.5)	23 (12.0)	3.0 (2.0)
I am confident about giving advice and counselling to breastfeeding women	11 (5.8)	25 (13.0)	54 (28.1)	83 (43.2)	19 (9.9)	4.0 (1.0)
I have sufficient knowledge to solve medication and health problems of breastfeeding women	14 (7.2)	36 (18.8)	45 (23.4)	75 (39.1)	22 (11.5)	4.0 (1.0)
Overall Scale						4.0 (1.0)
Undergraduate Training						
Pharmacy school provided appropriate training regarding advice and OTC therapy for pregnant women	15 (7.8)	40 (20.8)	52 (27.1)	70 (36.5)	15 (7.8)	3.0 (2.0)
Pharmacy school provided appropriate training regarding advice and OTC therapy for breastfeeding women	16 (8.4)	40 (20.8)	55 (28.6)	65 (33.9)	16 (8.3)	3.0 (2.0)
Overall Scale						3.0 (2.0)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49

Figure Legends

Figure 1 Pharmacists' responses to eight common symptoms in pregnancy and breastfeeding (n=192)

Figure 2 Pharmacists' responses to specific symptoms in pregnancy (n=192)

Figure 3 Pharmacists' responses to specific symptoms in breastfeeding (n=192)

For peer review only




Figure 1 Pharmacists' responses to eight common symptoms in pregnancy and breastfeeding (n=192)

164x70mm (300 x 300 DPI)

Peer review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2 Pharmacists' responses to specific symptoms in pregnancy (n=192)

111x51mm (300 x 300 DPI)




Figure 3 Pharmacists' responses to specific symptoms in breastfeeding (n=192)

90x48mm (300 x 300 DPI)

review only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Part 1. Demographic and other Characteristics**PLEASE FILL IN OR TICK (✓) THE APPROPRIATE ANSWER**

1.1 Gender: [1] Male [2] Female

1.2. Age in years:

1.3. Basic qualification in pharmacy: [1] BSC (Pharm) or B Pharm [2] M Pharm [3] Pharm D

1.4. Do you have any postgraduate qualification(s) in pharmacy? [1] Yes [2] No

1.5. If yes to question (1.4), please tick below all that apply:

[1] Diploma [2] Master [3] Ph.D [4] Others (Please state)

1.6. Number of years you have practiced as pharmacist since initial licensure

1.7. Location of pharmacy: [1] Hawalli [2] Al-Farwaniyah [3] Al-Ahmadi/ Mubarak Al-kabeer
[4] Al-Asima (Capital) [5] Al-Jahra

PART 2: Services provided regarding self-care in pregnancy and lactation

2.1 Do you have information leaflet or brochure to promote health for pregnant and breastfeeding women? [1] Yes [2] No

2.2 If yes to question (2.1), please tick the appropriate options

[1] Make it by myself [2] Drug company [3] Organization (please state).....

[4] Others (please state).....

2.3 Do you have experience in providing services for pregnant women? [1] Yes [2] No
(If no experience, please go to question 2.7)

2.4. If yes to question (2.3), how many pregnant women do receive your services in this pharmacy per week?

2.5. If yes to question (2.3), please rank the three services that you give most frequently for pregnant women. (i.e., first, second, third)

..... Advice about a suitable behavior such as lifestyle, exercise

..... Recommend vitamin and food supplements

..... Dispensing herbal medicine

..... Dispensing medicine

..... Refer to a doctor

..... Others (Please state).....

2.6. If yes to question (2.3), please indicate two symptoms and/or questions that pregnant women most frequently consulted you about in your pharmacy in the past.

1.....

2.....

1 **2.7. Do you have experience in providing services for breastfeeding women?** [1] Yes [2] No
 2 (If no experience, please go to PART 3)

3
 4 **2.8. If yes to question (2.7), how many breastfeeding women do receive your services in this pharmacy per**
 5 **week?**

6
 7 **2.9. If yes to question (2.7), how do you mostly know that women who receive your services are pregnant**
 8 **or breastfeeding? (Please tick one option)**

9
 10 [1] Asking by Pharmacist [2] Women tell before get the services

11
 12 [3] Others (Please state)

13
 14 **2.10. If yes to question (2.7), please rank the three services that you give most frequently for breastfeeding**
 15 **mothers. (i.e., first, second, third)**

- 16Advice about a suitable behavior such as lifestyle, exercise
 17Advice about contraception such as contraceptive pill
 18Advice about weight control or lose weight
 19Dispensing herbal medicine
 20Recommend vitamin and food supplement
 21Dispensing medicine
 22Refer to doctor
 23Others (Please state).....

24
 25 **2.11. If yes to question (2.7), please indicate two symptoms and/or questions that breastfeeding women**
 26 **most frequently consulted you about in your pharmacy in the past.**

27
 28
 29
 30
 31
 32
 33
 34

35
 36 **PART 3: Recommendations when counseling women on self-care in pregnancy and lactation.**

37
 38 **3.1. If pregnant women consult you about their symptoms as listed below. Please tick one**
 39 **recommendation (service) that would be your response for each symptom. If you choose to dispense**
 40 **medicines or vitamin or herbal medicines, please write the name in the table.**

Symptom	Refer to doctor	Provide only advice without dispensing a medicine	Dispense medicines or vitamin or herbal medicines <u>Please write the name</u>
1. Nausea and vomiting			
2. Indigestion (heartburn)			
3. Headache			
4. Back pain			
5. Insomnia (difficulty in sleeping)			

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34	6. Sore throat			
	8. Dry cough			
	9. Productive cough			
	10. Fever and aches			
	11. Constipation			
	12. Diarrhea			
	13. Hemorrhoids			
	14. Vaginal itching and simple discharge			
	15. Varicose vein			
	16. Swelling feet and legs			

3.2 If breastfeeding women consult you about their symptoms as listed below. Please tick one recommendation (service) that would be your response for each symptom. If you choose to dispense medicines or vitamin or herbal medicines, please write the name in the table.

Symptom	Refer to doctor	Provide only advice without dispensing a medicine	Dispense medicines or vitamin or herbal medicines <u>Please write the name</u>
1. Sore or cracked nipple (small ulcerations (cuts) develop on the nipple)			
2. Engorgement (painful overfilling of the breasts with milk)			
3. Insufficient milk			
4. Mastitis (inflammation of a breast)			
5. Diarrhea			

6. Constipation			
7. Hemorrhoids			
8. Headache			
9. Sore throat			
10. Runny nose			
11. Dry cough			
12. Productive cough			

PART 4: Views about self-care in pregnancy and lactation

Please tick one option that best describes your own view

	1. Strongly disagree	2. Disagree	3. Neither disagree or agree	4. Agree	5. Strongly agree
4.1 Community pharmacists are qualified to provide advice and an over-the-counter (OTC) therapy to treat common and minor symptoms in pregnant women					
4.2 Community pharmacists are qualified to provide advice and an OTC therapy to treat common and minor symptoms in breastfeeding women					
4.3 Community pharmacists should recommend OTC therapy and counseling to treat common and minor symptoms in pregnant women					
4.4 Community pharmacists should recommend OTC therapy and counseling to treat common and minor symptoms in breastfeeding women					
4.5 OTC medicines are safe for pregnancy.					
4.6 OTC medicines are safe for breastfeeding.					
4.7 I am confident about giving advice and counselling to pregnant women					
4.8 I have sufficient knowledge to solve medication and health problems of pregnant women					
4.9 I am confident about giving advice and counselling to breastfeeding women					

1 2 3 4 5 6 7 8 9	4.10 I have sufficient knowledge to solve medication and health problems of breastfeeding women					
10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	4.11 Pharmacy school provided appropriate training regarding advice and OTC therapy for pregnant women.					
25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60	4.12 Pharmacy school provided appropriate training regarding advice and OTC therapy for breastfeeding women.					

4.14 Do a continuing education program regarding this topic would be of value/priority for your practice?

[1] Strongly disagree [2] disagree [3] Neutral [4] Agree [5] strongly agree

4.15 If you agree or strongly agree to participate in continuing education program, what would be the most convenient method of delivering continuing education to you? *(Please tick one option)*

[1] Attending lectures and workshops/seminars [2] Receiving regular news letters
 [3] Receiving distant learning packages [4] Others (Please state)

4.16 What is /are the most common source(s) of information do you use for responding to symptoms during pregnancy and breastfeeding and /or searching about medicines use in pregnancy and breastfeeding? *(You may tick more than one option)*

[1] Books [2] Journal articles [3] Handout [4] Website [5] Others (please state).....

Thank you for your time and cooperation in completing this questionnaire.

Names of medications recommended by pharmacists for eight common symptoms in pregnancy and breastfeeding

Common symptoms	Frequency (%) of pharmacists recommended medicines for pregnant women	Frequency (%) of pharmacists recommended medicines for breastfeeding women
4 Headache	*n=154	*n=169
5 Paracetamol tablet	151 (98.1%)	145 (85.8)
6 Ibuprofen tablet	3 (1.9%)	24 (14.2)
7 Sore throat**	*n= 106	*n=124
8 Lozenges	90 (84.9)	89 (71.8)
9 Paracetamol tablet	16 (15.1)	29 (23.4)
10 Vitamin C tablet	14 (13.2)	16 (12.9)
11 Amoxicillin capsule	4 (3.8)	6 (4.8)
12 Azithromycin tablet	2 (1.9)	2 (1.6)
13 Runny nose**	*n=113	*n=124
14 Normal saline spray	69 (61.1)	33 (26.6)
15 Loratadine tablet	17 (15.0)	30 (24.2)
16 Paracetamol and Pseudoephedrine tablet	19 (16.8)	22 (17.7)
17 Vitamin C tablet	21 (18.6)	25 (20.2)
18 Cetrizine tablet	7 (6.2)	17 (13.7)
19 Corticosteroid nasal spray	2 (1.8)	3 (1.6)
21 Dry cough	*n=110	*n=126
22 Prospan® syrup	86 (78.2)	75 (59.5)
23 Codilar® syrup	12 (10.9)	20 (15.9)
24 Cloperastine syrup	8 (7.3)	22 (17.5)
25 Butamirate syrup	4 (3.6)	9 (7.1)
27 Productive cough**	*n=129	*n=131
28 Prospan® syrup	113 (87.6)	85 (64.9)
29 Mucolytic syrup	15 (11.6)	9 (6.9)
30 Amoxicillin capsule	2 (1.6)	2 (1.5)
31 Loratadine tablet	3 (2.3)	24 (18.3)
32 Butamirate syrup	-	10 (7.6)
33 Cloperastine syrup	-	8 (6.1)
34 Constipation	*n=138	*n=134
35 Sterculia granules	63 (45.7)	43 (32.1)
36 Lactulose syrup	60 (43.5)	63 (47.0)
37 Glycerine suppository	6 (4.3)	11 (8.2)
38 Sennalax® tablet®	9 (6.5)	17 (12.7)
39 Diarrhea**	n=75	n=90
40 Kaolin + Pectin syrup	23 (30.7)	18 (20.0)
41 Nifuroxazide tablet	18 (24.0)	32 (35.6)
42 Loperamide tablet	16 (21.3)	26 (28.9)
43 Oral rehydration salts	25 (33.3)	21 (23.3)
44 Activated charcoal tablet	3 (4.0)	-
45 Hyoscine tablet	-	3 (3.3)
46 Hemorrhoids	n=84	n=95
47 Neo-healar® cream	35 (41.6)	25 (26.3)
48 Pilex® ointment®	24 (28.6)	17 (17.9)
49 Procto-Glyvenol® cream	15 (17.9)	24 (25.3)
50 Procto-Glyvenol® suppository	10 (11.9)	15 (15.8)
51 Daflon® tablet	-	2 (2.1)
52 Proctosedyl® suppository	-	12 (12.6)

53 *Number of pharmacists recommended dispensation of a medicine; ** May total > 100% because of recommended combination of medicines; Prospan® (herbal medicinal product containing ivy leaf extract); Codilar® (Chlorpheniramine + Dextromethorphan + Phenylephrine); Sennalax® (Sennosides + docusate); Neo-healar® (four medicinal plants: Lupinus Albus, Vateria Indica, Mentha Piperita, Aloe Vera); Pilex® (a mixture of herbs [Mimosa pudica, thistles, Vitex negundo, Calendula flowers, camphor], Zinc, and Borax); Procto-Glyvenol® (tribenosid + lidocaine); Daflon® (diosmin + hesperidin); Proctosedyl® (hydrocortisone + cinchocaine).

Names of medications recommended by respondents for specific symptoms in pregnancy	
Specific pregnancy symptoms	Frequency (%) of pharmacists recommended medicines
Nausea and vomiting	*n=139
Navidoxine® tablet	110 (79.1)
Ondasetron tablet	27 (19.4)
Domperidone tablet	1 (0.7)
Multivitamin capsule	1 (0.7)
Indigestion (heartburn)	*n=131
Gaviscon® suspension	89 (67.9)
Ranitidine tablet	20 (15.3)
Zymogen® tablet	22 (16.8)
Back pain**	*n=92
Paracetamol tablet	42 (45.7)
Diclofenac gel	27 (29.3)
Reparil® gel	7 (7.6)
Relaxnova® cream	21 (22.8)
Insomnia	*n=20
St John's wort®	5 (25.0)
Dream water®	10 (50.0)
Melatonin tablet	5 (25.0)
Fever and aches	*n=93
Paracetamol tablet	93 (100)
Vaginal itching and simple discharge	*n=67
Clotrimazole ovule	20 (29.9)
Clotrimazole cream	15 (22.4)
Clotrimazole wash	9 (13.4)
Miconazole cream	11 (16.4)
Miconazole ovule	12 (17.9)
Varicose veins	*n=28
Venoruton® gel	21 (75.0)
Reparil® gel	7 (25.0)
Swelling feet and legs	*n=15
Reparil® gel	11 (73.3)
Varixinal® gel	4 (26.7)

*Number of pharmacists recommended dispensation of a medicine; ** May total > 100% because of recommended combination of medicines; Zymogen® (Lipase + Proteas + Pepsin + Hemicellulase + Ox bile extract + Dimethylpolysiloxane + Vitamin B1); Reparil® (Aescinum + Diethylamini salicylas); Relaxnova® (Idrocilamide) ; St John's Wort® (Hypericum perforatum); Dream water® (Melatonin + 5-HTP + GABA); Venorutin® (Rutoside); Varixinal® (Ruscus Aculeatus Extract, Aesculus Hippocastanum Extract, Centella Asiatica Extract, Vaccinium Myrtillus Extract).

Names of medications recommended by respondents for specific symptoms in breastfeeding	
Common breastfeeding symptoms	Frequency (%) of pharmacists recommended medicines
Sore or cracked nipple	*n=167
Panthenol cream	53 (31.7)
Lanolin cream	45 (26.9)
Avalon Organics® moisturizer	35 (21.0)
Mustela® Nipple Cream	18 (10.8)
Mebo® ointment	12 (7.2)
Cocoa butter cream	4 (2.4)
Engorgement	*n=17
Moisturizing cream	6 (35.3)
Cabergolin tablet	5 (29.4)
Paracetamol tablet	4 (23.5)
Fucidin cream	2 (11.8)
Insufficient milk	*n=138
Fitolat® tablet	118 (85.5)
Domperidone tablet	10 (7.2)
Multivitamins Capsule	10 (7.2)
Mastitis	*n=17
Local antibiotic cream	3 (17.6)
Panthenol cream	6 (35.3)
Amoxicillin capsule	3 (17.6)
Paracetamol tablet	2 (11.8)
Diclofenac tablet	2 (11.8)
Fitolat® tablet	1 (5.9)
*Number of pharmacists recommended dispensation of a medicine; Avalon® (Lavender + prebiotics); Mebo® (Sesame oil + Beeswax + other edible herbs); Mustela® (Avocado peptides + Bisabolol + Shea butter + Lupeol + Vitamin B5); Fitolat® (herbal extracts from plants, such as Fennel, Fenugreek, Hops and Verbena)	