

Article

Feature Fusion of ICP-AES, UV-Vis and FT-MIR for Origin Traceability of *Boletus Edulis* Mushrooms in Combination with Chemometrics

Luming Qi ^{1,2}, Honggao Liu ³, Jieqing Li ³, Tao Li ^{4,*} and Yuanzhong Wang ^{1,*}

¹ Institute of Medicinal Plants, Yunnan Academy of Agricultural Sciences, Kunming 650200, China; 18669326801@163.com

² State Key Laboratory Breeding Base of Systematic Research, Development and Utilization of Chinese Medicine Resources, Chengdu University of Traditional Chinese Medicine, Chengdu 611137, China

³ College of Agronomy and Biotechnology, Yunnan Agricultural University, Kunming 650201, China; honggaoliu@126.com (H.L.); lijieqing2008@126.com (J.L.)

⁴ College of Resources and Environment, Yuxi Normal University, Yuxi 653100, China

* Correspondence: ltyx_1976@126.com (T.L.); boletus@126.com (Y.W.)




Figure S1: Eleven microhabitats of *Boletus edulis* in Yunnan Province



Figure S2: The real product of *Boletus edulis*




Figure S3: The UV-Vis fingerprints of mushroom chloroform extract from other ten geographical origins




Figure S4: The FT-MIR fingerprints of the mushroom samples from other ten geographical origins




Figure S5: Score plots of the GS-SVM model of mushroom cap




Figure S6: Score plots of the GS-SVM model of mushroom stipe