

Supplementary Materials

β -HPV Infection Correlates with Early Stages of Carcinogenesis in Skin Tumors and Patient-Derived Xenografts from a Kidney Transplant Recipient Cohort

Cinzia Borgogna^{1*}, Carlotta Olivero^{1,2*}, Simone Lanfredini², Federica Calati¹, Marco De Andrea^{1,3}, Elisa Zavattaro⁴, Paola Savoia⁴, Elena Trisolini⁵, Renzo Boldorini⁵, Girish Patel², and Marisa Gariglio¹.

¹ Virology Unit, Department of Translational Medicine, Novara Medical School, Italy

² European Cancer Stem Cell Research Institute, School of Biosciences, Cardiff University, UK

³ Virology Unit, Department of Public Health and Pediatric Sciences, Turin Medical School, Italy


⁴ Dermatology Unit, Department of Health Sciences, Novara Medical School, Italy

⁵ Pathology Unit, Department of Health Sciences, Novara Medical School, Italy

*These authors contributed equally to this work.

Corresponding Author: marisa.gariglio@med.uniupo.it

Figure 1S. Percentage of Δ Np63+ or MCM7+ epithelial cells in skin lesions from the KTR cohort (n= 24, FFPE n=36). Positive cells were counted on three random fields (20X magnification) for each area: Non-pathological epithelium, HPV-infected area, Tumor. Error bars, SD. *, $p < 0.05$.



Supplementary Materials and Methods

Statistical Analysis

For Δ Np63+ and MCM7+ cell quantification, mean value \pm one standard deviation was presented. The statistical significance between two experimental groups is indicated in the figure by an asterisk. Comparisons were made using the Student's t test. Calculated p -values less than or equal to 0.05 were considered to be of statistical significance.