

***Nemopilema nomurai* jellyfish venom exerts an anti-metastatic effect by inhibiting Smad- and NF- κ B-mediated epithelial–mesenchymal transition in HepG2 cells**

Hyunkyung Lee^{1,3}, Min Jung Pyo¹, Seong Kyeong Bae¹, Yunwi Heo⁴, Indu Choudhary¹, Duhyeon Hwang³, Hyeryeon Yang¹, Je-hein Kim⁴, Jinho Chae⁵, Chang Hoon Han², Changkeun Kang¹, Seungshic Yum^{6,7*}, Euikyung Kim^{1,3*}

Fig.2A





Fig.2B




E-cadherin
 β -catenin




Vimentin



Snail




Slug




N-caherin


Fig.4A



E-cadherin
β-catenin



GAPDH




Vimentin



Slug

Snail


Same gel blot in different exposure time



N-cadherin

Snail


Fig.4B



N-cadherin

Vimentin

Same gel blot in different exposure time




Slug

GAPDH


gap

n-cad

vi




Snail



β-catenin

Same gel blot in different exposure time



E-cadherin

E-CAD

Fig.5A




Fig.5B





Fig.5C



Smad4



p-Smad3



smad3

GAPDH

Fig.6A




Fig.6D



Fig.6E








Fig.6F



Same gel blot in different exposure time