

Table S4. Ultrasound findings at three months follow-up

	<i>Transvaginal ultrasonography</i>			<i>Sonohysterography</i>		
	Intervention n=51	Control n=44	P-value	Intervention n=51	Control n=44	P-value
Performed in number of women	36 (70.6%)	23 (52.3%)		24 (47.1%)	13 (29.5%)	
<i>Sagittal measurement</i>						
Depth (mm)	4.5 (3.6 – 5.9)	4.5 (3.5-7.0)	0.64	5.0 (2.9-7.8)	7.1 (4.0-8.0)	0.32
Residual myometrium (mm)	4.0 (3.0 – 5.9)	4.4 (3.4 – 6.9)	0.30	4.4 (3.1-6.0)	4.0 (3.1-5.9)	0.70
<i>Transversal measurement</i>						
Depth (mm)	4.1 (2.7-6.7)	5.0 (3.2 – 7.0)	0.26	4.5 (3.4-7.0)	6.8 (4.6-8.7)	0.23
Residual myometrium (mm)	5.4 (4.3-6.6)	4.1 (3.3 – 6.9)	0.39	4.8 (3.8-7.0)	6.0 (3.3-8.0)	0.53
Intrauterine fluid	6 (16.2%)	6 (26.1%)	0.35	n.a.	n.a.	

Values are in mm. Data are reported as 'n (valid percentage) or as 'median (IQR)'. Analyses are by intention to treat.