## Orthopaedic Survey

How would you mostly describe your orthopaedic practice?
Adult foot ankle surgery only $\square$
Paediatric orthopaedics only
Both adult foot & ankle surgery and paediatric orthopaedics $\square$
Lower limb surgery □
General orthopaedic surgery
General orthopaedic surgery with an interest in foot and ankle surgery $\Box$
General orthopaedic surgery with an interest in paediatric orthopaedics  General orthopaedic surgery with an interest in foot and ankle surgery and paediatric
orthopaedics
How many new referrals of patients with SEVERE pes cavus do you see per year?
<2 🗆
2-5 🗆
6-10 🗆
11-15 🗆
>15 🗆
How many patients with Charcot-Marie-tooth (HMSN) do you see per year?
<2 🗆
2-5 🗆
6-10 🗆
11-15□
>15 🗆
How many corrections of the hindfoot in patients with SEVERE pes cavus do you perform per year (please do not include forefoot surgery)?
<2 🗆
2-5 🗆
6-10 🗆
11-15 🗆
<b>√15</b> □

Consider a 13 year old child with Charcot-Marie-tooth with a typical pes cavus who has failed conservative management. What are the procedures that you would commonly carry out?

Gastrocnemius/Achilles lengthening ☐
Plantar fascia release
Robert Jones Procedure
Soft tissue procedure lesser toes □
Bony procedure lesser toes
Calcaneal osteotomy □
First metalarsal osteotomy□
Midfoot osteotomy □
Midfoot fusion +/- correction □
Triple arthrodesis □
Tibialis posterior tendon transfer $\square$
Peroneal tendon surgery
Consider a typical 24 year old patient with Charcot –Marie – Tooth, without any arthritic changes on radiographs and who has failed conservative treatment. What are the procedures that you would commonly carry out? (more than one response allowed)
Gastrocnemius/Achilles lengthening □
Plantar fascia release
Robert Jones Procedure
Soft tissue procedure lesser toes
Bony procedure lesser toes
Calcaneal osteotomy □
First metalarsal osteotomy
Midfoot osteotomy □
Midfoot osteotomy □  Midfoot fusion +/- correction □
Midfoot fusion +/- correction □