

Supplemental Table 4. Scores on psychosocial developmental domains of survivors of childhood DTC versus survivors of other childhood cancers (diagnosed at all ages)

	DTC survivors	Childhood Cancer Survivors (CCS)							
	n = 39	Total CCS n = 350		Leukemia/ lymphoma n = 175		Solid tumors n = 150		Brain tumors n = 25	
			<i>P</i> Value		<i>P</i> Value		<i>P</i> Value		<i>P</i> Value
Social development [†]	22 (20.3, 23)	21 (18, 22)	0.008	20 (18, 22)	0.004	21 (19, 23)	0.068	20 (17, 22)	0.003
Autonomy development [‡]	9 (8, 10)	9 (8, 10)	0.734	9 (8, 10)	0.693	9 (8, 10)	0.974	9 (8, 10)	0.278
Psychosexual development [§]	8 (6.5, 8)	7 (5, 8)	0.036	7 (5, 8)	0.042	7 (6, 8)	0.097	5 (4, 8)	0.007

Scores are shown as median (p25, p27). Comparisons between DTC survivors and other groups were performed using Mann Whitney U tests. Higher score indicates achievement of earlier or more psychosocial developmental milestones; †, scale ranges 12-24; ‡, scale ranges 6-12; §, scale ranges 4-8. Missing values were excluded for statistical testing. *P* values in bold indicate a significant value ($P < 0.01$). Abbreviation: DTC, differentiated thyroid carcinoma; CCS, childhood cancer survivors.