

Supporting Information

New biosourced AA and AB monomers from 1,4:3,6-dianhydrohexitols, Isosorbide, Isomannide and Isoidide

Asma Saadaoui^{a,c,d}, Raouf Medimagh^{a*}, Sylvain Marque^b, Damien Prim^b, Saber Chatti^c, Herve Casabianca^c& Mongia Said Zina^d

^aInstitut National de Recherche et d'Analyse Physico-chimique (INRAP), Laboratoire des Substances Naturelles (LR10INRAP02), Sidi ThabetBiotechpole, Ariana 2020, (Tunisia).

^bUniversité de Versailles-St-Quentin (UVSQ), Institut Lavoisier de Versailles (ILV) UMR CNRS 8180, 45 Avenue des Etats-Unis, 78035 Versailles Cedex, France

^cUniversité de Lyon 1, Institut des Sciences Analytiques, UMR5280, CNRS, ENS-Lyon, 5 rue de la Doua, F-69100 Villeurbanne (France).

^dFaculté des Sciences de Tunis, Université de Tunis El Manar 2092 Tunis (Tunisia).




Figure S1. 300 MHz 2D correlation spectra (A) COSY and (B) HSQC of **11**




Figure S2. ¹H spectra of **11** (A), **12** (B) and **13** (C).




Figure S3. ^{13}C spectra of 11 (A), 12 (B) and 13 (C)




Figure S4. 300 MHz correlation spectrum (COSY) of **1b**.




Figure S5. 300 MHz correlation spectrum (HSQC) of **1b**




Figure S6. Zoom of 500 MHz correlation spectrum (HSQC) of **1h**




Figure S7. ¹H spectra of **1g** (A) and **Pe** (B) [DMSO-*d*6, 500 MHz]