

Supplementary Online Content

Marzec LN, Peterson PN, Bao H, et al. Use of cardiac resynchronization therapy among eligible patients receiving an implantable cardioverter defibrillator: insights from the National Cardiovascular Data Registry Implantable Cardioverter Defibrillator Registry. *JAMA Cardiol.* doi:10.1001/jamacardio.2016.5388

eTable. Patient Criteria Based on Strength of Guideline Recommendation for CRT.

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Patient Criteria Based on Strength of Guideline Recommendation for CRT

2008 Guidelines	Patients implanted between April 1, 2010 and August 31, 2012
Class I	LVEF \leq 35%, QRS duration \geq 0.12 seconds, sinus rhythm, NYHA III or ambulatory IV
Class IIa	LVEF \leq 35%, QRS duration \geq 0.12 seconds, atrial fibrillation, NYHA III or ambulatory IV
2012 Focused Update	Patients implanted between September 1, 2012 and June 30, 2014
Class I	LVEF \leq 35%, LBBB, QRS duration \geq 0.15 seconds, sinus rhythm, NYHA II, III, or ambulatory IV
Class IIa	LVEF \leq 35%, LBBB, QRS duration 0.12-0.149 seconds, sinus rhythm, NYHA II, III, or ambulatory IV
	LVEF \leq 35%, non-LBBB, QRS duration \geq 0.15 seconds, sinus rhythm, NYHA III or ambulatory IV
	LVEF \leq 35%, atrial fibrillation, LBBB, QRS duration \geq 0.15 seconds, NYHA II, III, or ambulatory IV

Abbreviations: LVEF, left ventricular ejection fraction; NYHA, New York Heart Association; LBBB, left bundle branch block