

Fig. S1. Effect of dimethylformamide (DMF) on chloroplast auto-fluorescence.
CLSM micrographs of *Zygnema* S (A, B) and *Klebsormidium crenulatum* (C, D) filaments before (A, C) and after (B, D) incubation in DMF. Bars = 10 μ m.




Fig. S2. Immunolabelling of whole cells and semithin sections (CLSM micrographs).
Zygnema S (A-D) and *Klebsormidium crenulatum* (E, F) filaments labelled with the monoclonal antibodies LM15 or 400-4. Whole cell labelling (A, B), labelling of semithin sections (C-F). Red autofluorescence is shown. (A) Weak fluorescence in outer and cross cell walls. (B) Recently fragmented filament with fluorescence close to terminal cross cell walls. (C-F) Labelling in outer and cross cell walls. Bars 10 = μ m




Fig. S3. Hemicellulose distribution in filamentous charophytes. Schemes are based on immuno-localisations of hemicellulose epitopes (antibody names in brackets) in young (1 month) and old (12 months) filaments of (A, B) *Zygnema S* and (C, D) *Klebsormidium crenulatum*.




Fig. S4. Photosynthetic performance of *Zygnema S* and *Klebsormidium crenulatum*. NIR and corresponding $Y(II)$ images (false colour) of young (A, C) and old (B, D) filaments of *Zygnema S* (A, B) and *Klebsormidium* (C, D). The colour bar at the top indicates the relative $Y(II)$ as a percentage. (A) Vacuolated cells containing two stellate chloroplasts and high $Y(II)$ values (~0.64). (B) Akinete cells dominate old filaments and exhibit a lower $Y(II)$ (i.e. ~0.41) than younger (i.e. vacuolated, thin-walled, growing) cells within the same filament (~0.6). (C) Cells of young filaments with a higher $Y(II)$ (~0.48-0.59) than cells of old filaments (D) (~0.3). Bars = 10 μ m.

Tab. S1 Evidence for major cell wall polysaccharides in Charophytes.

Cell wall component	Klebsormidiophyceae	Charaphyceae	Zygnematophyceae
Cellulose	A ^{1,5} B ⁷ C ⁶	A ⁵ B ^{8,7}	A ⁵ B ^{3,5,7}
Callose	A ⁵ B ^{3,5} C ⁶	A ⁵ B ^{5,8}	A ⁵ B ³
Pectins			
HG	A ^{1,5*} B ⁵ C ⁶	A ^{5,7} B ⁸	A ^{5,7} B ^{4,9,10}
RGI/RGII	-	A ^{5,7†}	A ^{7†}
Hemicelluloses			
Xyloglucan	B ⁵ C ⁶	A ⁵ B ^{5,8}	B ^{5,8}
MLG	B ⁵	B ^{5,8}	A ^{4,5} B ^{4,5}
Mannans	B ⁵	B ^{5,8}	B ⁵
Xylans	B ⁵	B ^{5,8}	B ⁵
(Glyco)Proteins			
Arabinogalactan proteins	-	B ⁵	B ⁵
Expansins	C ⁶	C ¹¹	C ¹¹
Extensins	C ¹²	-	B ⁵

Key: A biochemical, B immunological/histochemical, C genetically (genomic/transcriptomic data), '-' no evidence.

*Biochemical evidence for HG in Klebsormidiophyceae is ambiguous. While Domozych et al. (1980) and Sorensen et al. (2011) detected galacturonic acid in *K. flaccidum*, O'Rourke et al. 2015 suggested its absence in *K. flaccidum* and *K. subtile*.

†Suggested by the co-occurrence of 1→5-Ara f, 1→4-Gal p and 2→4-Rha p.

¹Domozych et al. 1980, ²Popper and Fry 2003, ³Herburger and Holzinger 2015, ⁴Eder et al. 2008, ⁵Sørensen et al. 2011, ⁶Hori et al. 2011, ⁷O'Rourke et al. 2015, ⁸Domozych et al. 2009, ⁹Domozych et al. 2006, ¹⁰Eder and Lütz-Meindl 2010,

¹¹Vannerum et al. 2011, ¹²Liu et al. 2016

Referneces not in the main document:

- Domozych, D. S., Sørensen, I. and Willats, W. G. (2009). The distribution of cell wall polymers during antheridium development and spermatogenesis in the Charophycean green alga, *Chara corallina*. Ann. Bot. 104, 1045-1056.
- Liu, X., Wolfe, R., Welch, L. R., Domozych, D. S., Popper, Z. A. and Showalter, A. M. (2016). Bioinformatic identification and analysis of extensins in the plant kingdom. PLoS ONE 11, e0150177.
- Popper, Z. A. and Fry, S. C. (2003). Primary cell wall composition of bryophytes and charophytes. Ann. Bot. 91, 1-12.
- Vannerum, K., Huysman, M. J., De Rycke, R., Vuylsteke, M., Leliaert, F., Pollier, J., Lütz-Meindl, U., Gillard, J., De Veylder, L., Goossens, A. et al. (2011). Transcriptional analysis of cell growth and morphogenesis in the unicellular green alga *Micrasterias* (Streptophyta), with emphasis on the role of expansin. BMC Plant Biol. 11, 128.