



SUPPLEMENTARY INFORMATION

Data on HBV immunization coverage among 1-year-olds, prevalence of obesity (percentage of adult population with a body mass index (BMI) ≥ 30 kg/m²), and alcohol per capita (APC) consumption were obtained from the WHO Indicator and Measurement Registry.¹ APC consumption (total amount of alcohol consumed per adult (15+ years) over a calendar year in litres of pure alcohol) was available yearly by country from 1980-2014, and projections available for 2015, 2020 and 2025.


Annex Figure 1. Percentage of defined adult population with a body mass index (BMI) of 30 Kg/m² or higher in 2010 for 30 countries obesity (WHO Indicator and Measurement Registry)¹



Annex Figure 2. Recorded alcohol per capita (for 15+ year olds) consumption (in litres of pure alcohol – all types) for 1980-2013 (solid lines) and predicted alcohol consumption (dashed lines) for 2015, 2020 and 2025 for 29 countries (projections data for Estonia is missing; WHO Indicator and Measurement Registry)¹



Annex Figure 3. Scatterplot of age-standardised incidence rates of PLC per 100,000, both sexes, in 2005 and recorded alcohol per capita (for 15+ year olds) consumption (in litres of pure alcohol – all types) for 1995* (WHO Indicator and Measurement Registry)¹



* $r=0.625$, $p\text{-value}<0.01$ when excluding Asian countries; $r=0.270$ $p\text{-value}=0.149$ including Asia

1. World Health Organization. The World Health Organization Indicator and Measurement Registry http://apps.who.int/gho/indicatorregistry/App_Main/indicator_registry.aspx (accessed 2 June 2016).