

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK

Supplementary Data – DNA fingerprinting gel images




Figure S1 PCR products of eight durian types at locus DZ01 on 8% (w/v) polyacrylamide gel




Figure S2 PCR products of eight durian types at locus DZ01 on 8% (w/v) polyacrylamide gel

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK




Figure S3 PCR products of seven durian types at locus DZ01 on 8% (w/v) polyacrylamide gel




Figure S4 PCR products of three durian types at locus DZ01 on 8% (w/v) polyacrylamide gel

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK




Figure S5 PCR products of two durian types and three durian types at loci DZ01 and DZ04 respectively on 8% (w/v) polyacrylamide gel




Figure S6 PCR products of eight durian types at locus DZ02 on 8% (w/v) polyacrylamide gel

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK



Figure S7 PCR products of eight durian types at locus DZ02 on 8% (w/v) polyacrylamide gel




Figure S8 PCR products of eight durian types at locus DZ02 on 8% (w/v) polyacrylamide gel

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK




Figure S9 PCR products of three durian types at locus DZ02 on 8% (w/v) polyacrylamide gel




Figure S10 PCR products of eight durian types at locus DZ03 on 8% (w/v) polyacrylamide gel

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK




Figure S11 PCR products of eight durian types at locus DZ03 on 8% (w/v) polyacrylamide gel




Figure S12 PCR products of eight durian types at locus DZ03 on 8% (w/v) polyacrylamide gel

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK




Figure S13 PCR products of three durian types at locus DZ03 on 8% (w/v) polyacrylamide gel




Figure S14 PCR products of eight durian types at locus DZ04 on 8% (w/v) polyacrylamide gel

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK




Figure S15 PCR products of eight durian types at locus DZ04 on 8% (w/v) polyacrylamide gel




Figure S16 PCR products of eight durian types at locus DZ04 on 8% (w/v) polyacrylamide gel

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK




Figure S17 PCR products of eight durian types at locus DZ05 on 8% (w/v) polyacrylamide gel




Figure S18 PCR products of five durian types at locus DZ05 on 8% (w/v) polyacrylamide gel

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK




Figure S19 PCR products of eight durian types at locus DZ05 on 8% (w/v) polyacrylamide gel




Figure S20 PCR products of six durian types at locus DZ05 on 8% (w/v) polyacrylamide gel

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK




Figure S21 PCR products of eight durian types at locus DZ07 on 8% (w/v) polyacrylamide gel




Figure S22 PCR products of seven durian types at locus DZ07 on 8% (w/v) polyacrylamide gel

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK




Figure S23 PCR products of eight durian types at locus DZ07 on 8% (w/v) polyacrylamide gel




Figure S24 PCR products of four durian types at locus DZ07 on 8% (w/v) polyacrylamide gel

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK




Figure S25 PCR products of eight durian types at locus DZ08 on 8% (w/v) polyacrylamide gel




Figure S26 PCR products of eight durian types at locus DZ08 on 8% (w/v) polyacrylamide gel

Genetic variation and DNA fingerprinting of durian types in Malaysia using simple sequence repeat (SSR) markers

Siew GY, Ng WL, Tan SW, Alitheen NB, Tan SG, Yeap SK




Figure S27 PCR products of eight durian types at locus DZ08 on 8% (w/v) polyacrylamide gel




Figure S28 PCR products of three durian types at locus DZ08 on 8% (w/v) polyacrylamide gel