

Supplementary Online Content

Sampaio-Junior B, Tortella G, Borrione L, et al. Efficacy and safety of transcranial direct current stimulation as an add-on treatment for bipolar depression: a randomized clinical trial. *JAMA Psychiatry*. Published online December 27, 2017.

doi:10.1001/jamapsychiatry.2017.4040

eTable. Mood Scores Over Time

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Mood Scores Over Time

	Group	Baseline	Week 2	Week 4	Week 6
Intention to treat, mean (SD)					
HDRS	Active	23.1 (3.9)	11.4 (5.5)	10.7 (5.3)	10.3 (5.6)
	Sham	23.5 (4.7)	15.8 (8.3)	14.7 (7.4)	16.2 (7.7)
MADRS	Active	28.7 (5.3)	15.1 (6.9)	14.2 (7.3)	13.6 (8)
	Sham	27.9 (5)	19.5 (10.4)	18.3 (10)	19 (10.3)
CGI	Active	3.5 (0.8)	2 (1)	2 (1.1)	1.8 (1.2)
	Sham	3.4 (0.7)	2.5 (1.3)	2.2 (1.2)	2.3 (1.3)
YMRS	Active	2.4 (2)	1.8 (2.1)	2.3 (2.8)	2.3 (3)
	Sham	2 (1.9)	2 (2.8)	2.9 (3.3)	2.3 (2.7)
Per protocol, mean (SD)					
HDRS	Active	23.1 (3.9)	10.5 (4.2)	9.8 (4)	9.2 (4.3)
	Sham	23.5 (4.7)	15.3 (8.3)	13.4 (6.6)	15.2 (7.2)
MADRS	Active	28.7 (5.3)	14.8 (5.5)	12.9 (6.4)	12 (7.1)
	Sham	27.9 (5)	18.9 (10.4)	16.8 (9.2)	17.5 (9.8)
CGI	Active	3.5 (0.8)	1.9 (1)	1.8 (1.1)	1.6 (1.1)
	Sham	3.4 (0.7)	2.4 (1.3)	2.1 (1.1)	2.1 (1.3)
YMRS	Active	2.4 (2)	1.6 (2.1)	2.2 (3)	2.3 (3.2)
	Sham	2 (1.9)	2 (2.9)	3.1 (3.3)	2.4 (2.8)

HDRS-17, Hamilton Depression Rating Scale-17 item version. MADRS, Montgomery-Åsberg Depression Rating Scale. YMRS, Young Manic Rating Scale. CGI, Clinical Global Impression scale. For the intention-to-treat scores, mean (SD) values were estimated using the last observation carried forward approach. For the per protocol scores, mean (SD) values were estimated based in the completer sample.