

Table S2. Death before discharge of infants 22-28 weeks' gestation by antenatal corticosteroid treatment

Gestational age	Any ANS	No ANS	ARR (95% CI) ^a
	N=9827	N=2028	
	n/total n (%)	n/total n (%)	
22 wk	80/106 (75.5)	519/536 (96.8)	0.79 (0.70 – 0.90) ^{b,c}
23 wk	494/904 (54.6)	438/519 (84.4)	0.68 (0.63 – 0.74) ^{b,c}
24 wk	696/1947 (35.7)	134/236 (56.8)	0.58 (0.50 – 0.66) ^{b,c}
25 wk	496/2183 (22.7)	78/260 (30.0)	0.78 (0.61 – 1.00)
26 wk	294/2011 (14.6)	55/251 (21.9)	0.60 (0.45 – 0.79) ^{b,c}
27 wk	189/1644 (11.5)	29/134 (21.6)	0.59 (0.40 – 0.89) ^{b,c}
28 wk	101/1032 (9.8)	13/92 (14.1)	0.69 (0.39 – 1.22) ^c

Includes data on 883 infants who died within 12 h without receiving delivery room resuscitation.

ANS, antenatal corticosteroid; ARR, adjusted relative risk; CI, confidence interval.

^a ARR and 95% CI are estimated with no ANS (not exposed to ANS) group used as referent category-models adjust for birthweight, sex, multiple births, small for gestational age, maternal variables (age, marital status, race, diabetes, rupture of membranes >24 h, antepartum hemorrhage, and mode of delivery), center, and epoch; ^b Significant with P value <.05; ^c Model does not adjust for center-model did not converge with center included.