

SUPPLEMENTARY TABLE S2. UPPER BOUNDS OF THE COEFFICIENT OF VARIATION (CV) FOR EACH QUARTILE
FROM INTERQUARTILE ANALYSIS OF GLYCEMIC VARIABILITY DATA IN ADULTS (>18 y/o)
WITHOUT DIABETES, WITH TYPE 2 DIABETES AND WITH TYPE 1 DIABETES

| <i>CV metric for glycemic variability</i> | <i>Adults without diabetes (%)</i> | <i>Adults with type 2 diabetes (%)</i> | <i>Adults with type 1 diabetes (%)</i> | <i>Children with type 1 diabetes (%)</i> | <i>Adolescents with type 1 diabetes (%)</i> |
|---|------------------------------------|--|--|--|---|
| 0th Pct. (min) | 11.8 | 15.6 | 24.5 | 23.0 | 22.4 |
| 2.5th Pct. | 12.3 | 16.5 | 28.1 | 25.4 | 26.4 |
| 25th Pct. | 15.0 | 25.4 | 36.2 | 34.8 | 35.9 |
| 50th Pct. | 16.7 | 30.4 | 40.3 | 37.7 | 39.0 |
| 75th Pct. | 18.1 | 35.4 | 44.4 | 40.5 | 42.5 |
| 97.5th Pct. | 20.9 | 42.1 | 52.5 | 44.5 | 50.5 |
| 100th Pct. (max) | 22.0 | 43.7 | 56.9 | 44.6 | 53.7 |

Also shown is the interquartile analysis for children (<13 y/o) and adolescents (13–18 y/o) with type 1 diabetes. CV, coefficient of variation.