

S1 Table. Sensitivity Analysis for 2005 Sample (n=337) Compared With Total Cohort (n=1021).

		[†] 2005 SAMPLE n=337	^{††} TOTAL COHORT n=1021	<i>p</i>
TOTAL	n	337	1021	
GENDER	Male	260 (77.2%)	764 (74.8%)	0.265 ^a
AGE	Mean years	41.7	40.4	0.134 ^b
ETHNICITY	White	184 (54.6%)	512 (50.1%)	0.407 ^a
BMI	Mean kg/m ²	25.8	25.5	0.835 ^b
WAIST	Mean cm	91.3	90.1	0.132 ^b
HYPERTENSION	n	66 (19.6%)	192 (18.8%)	0.891 ^a
LIPIDS (mean)	Total Cholesterol	4.8	4.7	0.794 ^b
	Triglycerides	1.8	1.7	0.237 ^b
CURRENT SMOKER	n	120 (35.6%)	378 (37.0%)	0.326 ^a
CVD	n	9 (2.7%)	32 (3.1%)	0.563 ^a
STATIN USE	n	54 (16.0%)	129 (12.6%)	0.268 ^a
HIV DURATION	Mean years	6.3	6.5	0.249 ^b
ARV NAIVE	n	66 (19.6%)	223 (21.8%)	0.426 ^a
LIPODYSTROPHY	n	92 (27.3%)	216 (21.2%)	0.336 ^a
HEPATITIS B	n	15 (4.5%)	50 (4.9%)	0.981 ^a
HEPATITIS C	n	12 (3.6%)	34 (3.3%)	0.831 ^a

Notes:

[†] Participants where glycaemia was measured

^{††} Total sample including where glycaemia was not measured

^a Significance of difference by Chi-squared

^b Significance of difference by ANOVA