

Supplementary Table SVI Age-adjusted and multivariable odds ratios of early menopause by BMI in 1989 excluding women who had ulcerative colitis, Crohn's disease, rheumatoid arthritis or systemic lupus before age 45, Nurses' Health Study II, 1989–2011^a.

Adiposity measure	Total	Cases	Age-adjusted model		Multivariable-adjusted model ^b	
			OR	95% CI	OR	95% CI
BMI, 1989 (kg/m ²)						
<18.5	2546	117	1.29	1.06, 1.57	1.30	1.07, 1.58
18.5–22.4	34 454	1233	ref		ref	
22.5–24.9	17 007	596	0.98	0.89, 1.09	0.94	0.85, 1.04
25.0–27.4	9402	288	0.86	0.75, 0.98	0.80	0.70, 0.92
27.5–29.9	4236	112	0.74	0.61, 0.90	0.68	0.56, 0.83
30.0–34.9	4937	162	0.92	0.78, 1.09	0.83	0.70, 0.99
≥35.0	3073	129	1.19	0.99, 1.44	1.03	0.85, 1.25

^an = 75 655.

^bAdjusted for age, race/ethnicity, physical activity, pack-years cigarette smoking, alcohol consumption, oral contraceptive use, parity, age at menarche and breastfeeding. See Table II footnote for categorization of covariates.