

Supplementary Table I Baseline characteristics of study participants and women not included in the study (2007–2014).

	Total cohort	Not included	P value
N	100	132	
Cycles	136	218	
		Mean (SD) or n (%)	
Personal characteristics			
Age, years	34.8 (3.8)	35.8 (4.0)	0.09
BMI, kg/m ²	24.4 (4.0)	24.0 (4.3)	0.13
Ever smoker, n (%)	29 (29.0)	37 (28.0)	0.88
White/Caucasian, n (%)	82 (82.0)	110 (83.3)	0.86
Moderate-to-vigorous exercise, h/week	4.3 (4.7)	3.3 (3.7)	0.19
Multivitamin, n (%)	91 (91.0)	113 (87.6)	0.52
Reproductive characteristics			
Infertility diagnosis, n (%)			0.35
Male factor	35 (35.0)	46 (34.8)	
Female factor	24 (24.0)	42 (31.8)	
Unexplained	41 (41.0)	44 (33.3)	
Treatment protocol, n (%)			0.40
Follicular phase GnRH antagonist	10 (10.0)	12 (9.1)	
Follicular phase GnRH agonist	9 (9.0)	20 (15.2)	
Luteal phase GnRH agonist	81 (81.0)	100 (75.8)	
Day 3 FSH, IU/L	7.0 (2.0)	7.6 (3.2)	0.27

P values were based on Kruskal–Wallis test (for continuous variable) or Fisher's exact test (for categorical variables).