

Village		TPN												
Survey of blood collection	Total population	Characteristic of blood spot						Antibody prevalence % , (n/N) , (95%CI)						
		Blood Collection date	Number of dry blood spot	Male	Female	Median age (range) (y)		Sex			Age			
								Male	Female	Total	0-4	5-15	15-59	60 up
M0	373	20/05/2013	258	146	112	27.3 (1-66)	% (95% CI) (n/N)	45,2 (37.0-53.6) (66/146)	42,9 (33.7-52.6) (48/112)	44,2 (38.1-50.0) (114/258)	33,3 (6.0-75.9) (2/6)	41,0 (30.2-52.8) (32/78)	45,2 (37.6-53.1) (76/168)	66,7 (24.1-94.0) (4/6)
M3	406	23/08/2013	283	154	129	24.8 (1-66)	% (95% CI) (n/N)	57,8 (49.6-65.6) (89/154)	60,5 (51.5-68.9) (78/129)	59,0 (53.0-64.8) (167/283)	51,7 (32.9-70.1) (15/29)	53,7 (42.4-64.6) (44/82)	63,0 (55.1-70.3) (104-165)	57,1 (20.2-88.2) (4/7)
M6	434	07/11/2013	300	158	142	24.8 (0-66)	% (95% CI) (n/N)	49,4 (41.4-57.4) (78/158)	49,3 (40.9-57.8) (70/142)	49,3 (43.6-55.1) (148/300)	43,6 (28.2-60.2) (17/39)	57,8 (46.5-68.4) (48/83)	45,6 (38.0-53.4) (78/171)	71,4 (30.3-94.9) (5/7)
M9	464	28/01/2014	307	160	147	23.9 (0-66)	% (95% CI) (n/N)	50,0 (42.0-58.0) (80/160)	36,1 (28.4-44.4) (53/147)	43,3 (37.7-49.1) (133/307)	34,8 (21.8-50.3) (16/46)	47,7 (36.9-58.7) (41/86)	43,4 (35.8-51.3) (72/166)	44,4 (15.3-77.3) (4/9)
M12	473	23/04/2014	260	135	125	22.8 (0-66)	% (95% CI) (n/N)	60,7 (51.9-68.9) (82/135)	68,0 (59.0-75.9) (85/125)	64,2 (58.0-70.0) (167/260)	65,1 (49.0-78.6) (28/43)	63,4 (52.0-73.6) (52/82)	65,9 (57.0-73.9) (85/129)	33,3 (6.0-75.9) (2/6)
M15	492	14/07/2014	233	127	106	24.3 (0-66)	% (95% CI) (n/N)	80,3 (72.1-86.6) (102/127)	76,4 (67.0-83.9) (81/106)	78,5 (72.6-83.5) (183/233)	70,6 (52.3-84.3) (24/34)	83,1 (71.3-90.9) (54/65)	77,2 (68.7-84.0) (98/127)	100,0 (56.1-100.0) (7/7)
M18	501	08-10-14	265	139	126	23.0 (0-66)	% (95% CI) (n/N)	83,5 (76.0-89.0) (116/139)	81,7 (73.7-87.9) (103-126)	82,6 (77.4-86.9) (219/265)	82,2 (67.4-91.5) (37/45)	78,2 (67.2-86.5) (61/78)	85,3 (78.0-90.6) (116/136)	83,3 (36.5-99.1) (5/6)

Village		TOT												
Survey of blood collection	Total population	Characteristic of blood spot					Antibody prevalence % , (n/N) , (95%CI)							
		Blood Collection date	Number of dry blood spot	Male	Female	Median age (range) (y)	Sex			Age				
							Male	Female	Total	0-4	5-15	15-59	60 up	
M0	740	14/06/2013	404	192	212	26.8 (0-80)	% (95% CI) (n/N)	68,2 (61.1-74.7) (131/192)	62,7 (55.8-69.2) (133/212)	65,3 (60.5-69.9) (264/404)	42,9 (18.8-70.4) (6/14)	58,1 (49.3-66.4) (79/136)	72,4 (66.1-78.0) (168/232)	50,0 (28.8-71.2) (11/22)
M3	785	12/09/2013	322	149	173	22.2 (0-70)	% (95% CI) (n/N)	77,2 (69.5-83.5) (115/149)	71,7 (28.7-38.5) (124/173)	74,2 (69.0-78.8) (239/322)	63,6 (47.7-77.2) (28/44)	70,3 (61.1-78.2) (83/118)	78,8 (71.1-84.9) (115/146)	92,9 (64.2-99.6) (13/14)
M6	789	26/11/2013	201	98	103	24.1 (1-70)	% (95% CI) (n/N)	75,5 (65.6-83.4) (74/98)	81,6 (72.4-88.3) (84/103)	78,6 (72.2-83.9) (158/201)	72,0 (50.4-87.1) (18/25)	73,2 (61.2-82.7) (52/71)	82,6 (73.0-89.4) (76/92)	92,3 (62.1-99.6) (12/13)
M9	814	18/02/2014	292	144	148	21.7 (0-80)	% (95% CI) (n/N)	46,5 (38.3-55.0) (67/144)	47,3 (39.1-55.6) (70/148)	46,9 (41.1-52.8) (137/292)	40,0 (27.3-54.8) (22/55)	51,1 (40.3-61.9) (45/88)	45,7 (37.2-54.3) (63/138)	63,6 (31.6-87.6) (7/11)
M12	833	15/05/2014	249	114	135	19.3 (0-70)	% (95% CI) (n/N)	59,6 (50.0-68.6) (68/114)	56,3 (47.5-64.7) (76/135)	57,8 (51.4-64.0) (144/249)	39,7 (27.8-52.8) (25/63)	59,1 (46.3-70.8) (39/66)	66,7 (57.0-75.2) (74/111)	66,7 (30.9-91.0) (6/9)
M15	861	06/08/2014	307	152	155	21.3 (0-80)	% (95% CI) (n/N)	92,1 (86.3-95.7) (140/152)	93,5 (88.1-96.7) (145/155)	92,8 (89.2-95.4) (285/307)	94,7 (84.5-98.6) (54/57)	92,0 (84.4-96.2) (92/100)	92,5 (86.4-96.2) (124/134)	93,8 (67.7-99.7) (15/16)
M18	879	30-10-14	195	93	102	19.5 (0-70)	% (95% CI) (n/N)	57,0 (46.3-67.1) (53/93)	74,5 (64.8-82.4) (76/102)	66,2 (59.0-72.7) (129/195)	47,8 (33.1-62.9) (22/46)	76,8 (63.3-86.6) (43/56)	72,1 (61.2-81.0) (62/86)	28,6 (5.1-69.7) (2/7)

Village		KNH												
Survey of blood collection	Total population	Characteristic of blood spot					Antibody prevalence % , (n/N) , (95%CI)							
		Blood Collection date	Number of dry blood spot	Male	Female	Median age (range) (y)	Sex			Age				
							Male	Female	Total	0-4	5-15	15-59	60 up	
M0	348	14/06/2013	325	180	145	29.1 (2-73)	% (95% CI) (n/N)	55,6 (48.0-62.9) (100/180)	51,7 (43.3-60.0) (75/145)	53,8 (48.3-59.3) (175/325)	33,3 (9.0-69.1) (3/9)	40,2 (30.0-51.3) (35/87)	59,3 (52.4-65.8) (128/216)	69,2 (38.9-89.7) (9/13)
M3	417	12/09/2013	277	147	130	28.1 (1-73)	% (95% CI) (n/N)	47,6 (39.4-56.0) (70/147)	51,5 (42.7-60.3) (67/130)	49,5 (43.4-55.5) (137/277)	30,4 (14.1-53.0) (7/23)	38,4 (27.4-50.5) (28/73)	56,7 (49.0-64.2) (97/171)	50,0 (20.1-79.9) (5/10)
M6	435	11/12/2013	263	141	122	28.1 (0-73)	% (95% CI) (n/N)	61,7 (53.1-69.7) (87/141)	52,5 (43.3-61.5) (64/122)	57,4 (51.2-63.4) (151/263)	59,1 (36.7-78.5) (13/22)	55,7 (43.4-67.4) (39/70)	56,5 (48.5-64.2) (91/161)	80,0 (44.2-96.5) (8/10)
M9	449	05/03/2014	301	155	146	25.9 (0-73)	% (95% CI) (n/N)	47,7 (39.7-55.9) (74/155)	49,3 (41.0-57.7) (72/146)	48,5 (42.8-54.3) (146/301)	43,2 (27.5-60.4) (16/37)	45,2 (34.5-56.4) (38/84)	52,1 (44.3-59.8) (88/169)	36,4 (12.4-68.4) (4/11)
M12	469	26/05/2014	298	160	138	25.1 (0-73)	% (95% CI) (n/N)	41,9 (34.2-49.9) (67/160)	39,1 (31.1-47.8) (54/138)	40,6 (35.0-46.4) (121/298)	31,7 (18.6-48.2) (13/41)	47,6 (36.7-58.7) (40/84)	38,4 (31.0-46.4) (63/164)	55,6 (22.7-84.7) (5/9)
M15	485	21/08/2014	300	162	138	26.0 (0-70)	% (95% CI) (n/N)	94,4 (89.4-97.3) (153/162)	96,4 (91.3-98.7) (133/138)	95,3 (92.1-97.3) (286/300)	81,0 (65.4-90.9) (34/42)	97,1 (89.1-99.5) (68/70)	97,8 (94.0-99.3) (174/178)	100,0 (65.6-100.0) (10/10)
M18	495	13-11-14	282	154	128	24.6 (0-70)	% (95% CI) (n/N)	86,4 (79.7-91.2) (133/154)	83,6 (75.8-89.3) (107/128)	85,1 (80.3-89.0) (240/282)	63,3 (48.3-76.2) (31/49)	84,6 (73.1-92.0) (55/65)	91,2 (85.4-94.9) (145/159)	100,0 (62.9-100.0) (9/9)

Village		HKT												
Survey of blood collection	Total population	Characteristic of blood spot					Antibody prevalence % , (n/N) , (95%CI)							
		Blood Collection date	Number of dry blood spot	Male	Female	Median age (range) (y)	Sex			Age				
							Male	Female	Total	0-4	5-15	15-59	60 up	
M0	899	04/07/2013	490	241	249	26.4 (0-94)	% (95% CI) (n/N)	92,9 (88.8-95.7) (224/241)	94,8 (91.0-97.1) (236/249)	93,9 (91.3-95.8) (460/490)	91,7 (71.5-98.6) (22/24)	92,2 (86.8-95.6) (154/167)	95,0 (91.6-97.2) (267/281)	94,4 (70.6-99.7) (17/18)
M3	979	07/10/2013	535	269	266	24.5 (0-89)	% (95% CI) (n/N)	88,5 (80.6-89.4) (230/269)	86,8 (82.0-90.5) (231/266)	86,2 (83.9-89.0) (461/535)	81,6 (67.5-90.8) (40/49)	85,7 (79.6-90.3) (156/182)	86,8 (82.2-90.4) (249/287)	94,1 (69.2-99.7) (16/17)
M6	1029	08/01/2014	500	243	257	22.1 (0-78)	% (95% CI) (n/N)	78,2 (72.4-83.1) (190/243)	85,6 (80.6-89.5) (220/257)	82,0 (78.3-85.2) (410/500)	75,8 (63.4-85.1) (50/66)	81,9 (75.6-86.9) (158/193)	84,8 (79.3-89.0) (195/230)	63,6 (31.6-87.6) (7/11)
M9	1070	01/05/2014	478	240	238	22.4 (0-78)	% (95% CI) (n/N)	87,9 (83.0-91.6) (211/240)	86,6 (81.4-90.5) (206/238)	87,2 (83.8-90.0) (417/478)	77,9 (65.9-86.7) (53/68)	86,9 (80.6-91.4) (146/168)	89,7 (84.9-93.2) (209/233)	100,0 (62.9-100.0) (9/9)
M12	1192	24/06/2014	527	269	258	21.7 (0-89)	% (95% CI) (n/N)	85,1 (80.2-89.0) (229/269)	89,1 (84.6-92.6) (230/258)	87,1 (83.9-89.8) (459/527)	87,6 (78.6-93.4) (78/89)	88,8 (83.1-92.8) (166/187)	85,7 (80.5-89.8) (204/238)	84,6 (53.7-97.3) (11/13)
M15	1217	16-09-14	506	266	240	21.6 (0-78)	% (95% CI) (n/N)	81,2 (75.9-85.6) (216/266)	85,0 (79.7-89.1) (204/240)	83,0 (79.4-86.1) (420/506)	82,1 (71.9-89.3) (69/84)	85,8 (79.6-90.4) (151/176)	81,3 (75.6-85.9) (191/235)	81,8 (47.8-96.8) (9/11)
M18	1236	08-12-14	467	233	234	20.0 (0-78)	% (95% CI) (n/N)	82,8 (77.2-87.3) (193/233)	87,2 (82.1-91.1) (204/234)	85,0 (81.4-88.1) (397/467)	84,0 (73.3-91.1) (63/75)	85,4 (79.6-89.8) (170/199)	84,9 (78.6-89.7) (152/179)	85,7 (56.2-97.5) (12/14)