

SUPPLEMENTARY TABLE S2. PREDICTORS OF INFANT
RECEIVING MOTHER'S OWN MILK FEEDINGS
AT NEONATAL INTENSIVE CARE UNIT DISCHARGE
(UNWEIGHTED MODEL)

| <i>Characteristic, n=205</i> | <i>OR (95% CI)</i> | <i>p</i> |
|--|--------------------|----------|
| Sociodemographic factors | | |
| Maternal age (years) | 1.07 (1.00–1.15) | 0.06 |
| Low socioeconomic status | 0.56 (0.22–1.45) | 0.23 |
| Married | 1.62 (0.64–4.07) | 0.31 |
| Race/ethnicity (white is reference) | | 0.90 |
| Hispanic | 1.03 (0.35–3.02) | 0.96 |
| black | 0.81 (0.25–2.64) | 0.73 |
| Highest education (<high school is reference) | | 0.16 |
| High school | 0.24 (0.07–0.87) | 0.03 |
| Trade or <BA/BS | 0.40 (0.11–1.43) | 0.16 |
| BA/BS or higher | 0.54 (0.13–2.30) | 0.40 |
| Previous breastfeeding experience (yes is reference) | | 0.12 |
| No (primiparous) | 2.47 (1.00–6.12) | 0.05 |
| No (multiparous) | 1.12 (0.33–3.83) | 0.86 |
| Prepregnancy health factors | | |
| Prepregnancy overweight/ obese (BMI ≥25) | 1.62 (0.71–3.68) | 0.25 |
| Hypertension | 0.61 (0.16–2.24) | 0.45 |
| Pregnancy complications | | |
| Multiple gestation | 0.44 (0.13–1.48) | 0.18 |
| Pre-eclampsia/eclampsia | 0.50 (0.11–2.15) | 0.35 |
| Premature rupture of membranes | 1.40 (0.35–5.63) | 0.63 |
| Placental abruption | 0.48 (0.10–2.19) | 0.34 |
| Fetal heart rate abnormalities or poor biophysical profile | 2.37 (0.65–8.66) | 0.19 |
| Intrauterine growth restriction | 0.52 (0.07–3.77) | 0.52 |
| Uterine abnormalities/ cerclage | 0.99 (0.25–3.93) | 0.99 |
| Chorioamnionitis | 0.59 (0.18–1.96) | 0.39 |
| Preterm labor | 0.87 (0.20–3.82) | 0.85 |
| Labor and delivery factors | | |
| Induction of labor | 0.97 (0.26–3.55) | 0.96 |
| Cesarean delivery | 0.85 (0.36–2.02) | 0.72 |
| Gestational age (weeks) | 1.14 (0.96–1.35) | 0.15 |
| MOM | | |
| CTV by day 14 | 8.76 (3.64–21.09) | <0.01* |

* $p < 0.05$.

BA/BS, bachelor of arts/bachelor of science; BMI, body mass index; CI, confidence interval; CTV, coming to volume; MOM, mother's own milk; OR, odds ratio.