

The cellular chloride channels CLIC1 and CLIC4 contribute to virus-mediated cell motility


Gabrielė Stakaitytė^{1,2#}, Nnenna Nwogu^{1,2#}, Jonathan D. Lippiat³, G. Eric Blair¹, Krzysztof Poterlowicz⁴, James R. Boyne⁴, Andrew Macdonald^{1,2}, Jamel Mankouri^{1,2}, and Adrian Whitehouse^{1,2}

¹School of Molecular and Cellular Biology, ²Astbury Centre for Structural Molecular Biology, ³School of Biomedical Sciences, Faculty of Biological Sciences, University of Leeds, Leeds, LS2 9JT, ⁴Centre for Skin Sciences, School of Chemistry and Biosciences, Faculty of Life Sciences, University of Bradford, Bradford, BD7 1DP, United Kingdom


Supplementary Information

Contents:
Supplemental Figures (Figures 1-2)


A(i)




(ii)




B(i)




(ii)





C(i)



(ii)



Supplementary Figure 1. Selection of non-toxic concentrations of each Cl⁻ channel modulator. Cell viability was measured in (A) i293-EGFP and i293-EGFP-ST and (B) MCC13 cells using an MTS-based assay in the presence of increasing concentrations of (i) DIDS, (ii) NPPB and (iii) RIAA inhibitors.



Supplementary Figure 2. Selection of non-toxic concentrations of NPPB channel modulator in MCPyV positive Merkel cell carcinoma cell lines. Cell viability was measured in **(A)** PeTa and **(B)** Waga cells using an MTS-based assay in the presence of increasing concentrations of NPPB inhibitor.