

### S3 Table

Summary of DARTseq analysis. Percentage of observed (Hobs), expected (Hexp) and missing datapoints derived from all and polymorphic DARTseq loci per 14 studied populations are shown

	Total (187)	Baglica (15)	Eskiyaygir (11)	Buyukatli (12)	Gurbuz (18)	Hisarkaya (14)	Kozludere (15)	Kebapci (19)	Kilavuzlu (6)	Kahraman- Maras West (13)	Kokluce (14)	Midyat (13)	Dagbasi (10)	Dogukent (5)	Yesilkoy (22)
mean Hobs	0.03	0.01	0.02	0.02	0.02	0.02	0.04	0.01	0.03	0.02	0.07	0.02	0.03	0.07	0.03
mean Hexp of polymorphic loci	0.18	0.13	0.32	0.27	0.10	0.14	0.28	0.08	0.22	0.26	0.27	0.18	0.32	0.40	0.31
missing data	0.15	0.12	0.13	0.15	0.16	0.16	0.18	0.13	0.18	0.21	0.12	0.17	0.11	0.12	0.14
polymorphic loci	0.50	0.07	0.28	0.17	0.20	0.20	0.26	0.04	0.12	0.26	0.46	0.12	0.27	0.22	0.44
samples in population	187	15	11	12	18	14	15	19	6	13	14	13	10	5	22
number of polymorphic loci	40818	2962	11516	7034	8039	8271	10684	1525	4840	10629	18658	4698	11103	8811	17788

