

Number of Responders Correctly identifying follow up after colonoscopy findings

CORRECT RESPONSE	PRE	POS T		PRE	POS T		PRE	POS T		PRE-TEST				POST-TEST					PRE	POS T	
	PGY 1 n=22	PGY 1 n=20	P	PGY 2 n=15	PGY 2 n=11	P	PGY 3 n=13	PGY 3 n=10	P	PGY 1 n=22	PGY 2 n=15	PGY 3 n=13	P	PGY 1 n=20	PGY 2 n=11	PGY3 n=10	P	Total n=50	Total n=41	P	
10 Years in Small Hyperplastic Rectal Polyps	9 (40.9%)	13 (65.0%)	0.1 19	9 (60.0%)	10 (90.1%)	0.1 78	13(1 00.0%)	9 (90.0%)	0. 43 5	9 (40.9%)	9 (60.0%)	13 (100.0%)	0.0 0 1	13 (65.0%)	10 (90.1%)	9 (90.0%)	0.2 22	31 (62.0%)	32 (78.0%)	0.099	
5 Years in 1 or 2 Small Tubular adenomas	9 (40.9%)	9 (45.0%)	0.7 89	9 (60.0%)	8 (72.7%)	0.6 83	10 (76.9%)	9 (90.0%)	0. 60 4	9 (40.9%)	9 (60.0%)	10 (76.9%)	0. 1 3 6	9 (45.0%)	8 (72.7%)	9 (90.0%)	0.0 42	28 (56.0%)	26 (63.3%)	0.474	
3 Years in 3 to 10 Adenomas	10 (45.5%)	10 (50.0%)	0.7 68	11 (73.3%)	10 (90.1%)	0.3 56	10 (76.9%)	8 (80.0%)		10 (45.5%)	11 (73.3%)	10 (76.9%)	0. 1 1 2	10 (50.0%)	10 (90.1%)	8 (80.0%)	0.0 54	31 (62.0%)	28 (68.2%)	0.099	
1 to 3 Years in > 10 Adenomas	4 (18.2%)	11 (55.0%)	0.0 23	3 (20.0%)	8 (72.7%)	0.0 15	7 (53.8%)	5 (50.0%)		4 (18.2%)	3 (20.0%)	7 (53.8%)	0. 7 3	11 (55.0%)	8 (72.7%)	5 (50.0%)	0.5 55	14 (28.0%)	24 (58.5%)	0.003	
2 to 6 Months in Sessile Adenomas	4 (18.2%)	10 (50.0%)	0.0 49	3 (20.0%)	7 (63.4%)	0.0 43	4 (30.8%)	6 (60.0%)	0. 22 3	4 (18.2%)	3 (20.0%)	4 (30.8%)	0. 6 8 6	10 (50.0%)	7 (63.4%)	6 (60.0%)	0.7 78	11(22 .02%)	23 (56.1%)	0.001	