

Supplementary Online Content

Miura M, Kobayashi T, Kaneko T, et al; the Z-score Project 2nd Stage (ZSP2) Study Group. Association of severity of coronary artery aneurysms in patients with Kawasaki disease and risk of later coronary events. *JAMA Pediatr*. Published online March 5, 2018.
doi:10.1001/jamapediatrics.2018.0030


eFigure 1. Kaplan-Meier survival curves for coronary events classified by the actual internal diameter of coronary artery aneurysms in male and female patients

eFigure 2. Kaplan-Meier survival curves for major severe cardiac events classified by the actual internal diameter of coronary artery aneurysms in male and female patients

This supplementary material has been provided by the authors to give readers additional information about their work.

eFigure 1. Kaplan-Meier survival curves for coronary events classified by the actual internal diameter of coronary artery aneurysms in male and female patients

Coronary event-free survival rate




Number at risk

D < 4	144	65	28	7	1
4 ≤ D < 6	279	146	72	14	1
6 ≤ D < 8	159	89	43	12	1
8 ≤ D	127	49	22	6	1

Time after diagnosis (years)

Coronary event-free survival rate



Number at risk


D < 4, 4 ≤ D < 6	56	26	15	2	0
6 ≤ D < 8	132	67	35	15	0
8 ≤ D	58	34	19	6	0
8 ≤ D	41	29	13	6	3

Time after diagnosis (years)

The survival curves were significantly different among the four groups ($P < .001$, log rank test of equality). Severity classification was based on < 4 mm, ≥ 4 to < 6 mm, ≥ 6 to < 8 mm, and ≥ 8 mm of the actual internal diameter (D) of coronary artery aneurysms.

eFigure 2. Kaplan-Meier survival curves for major severe cardiac events classified by the actual internal diameter of coronary artery aneurysms in male and female patients

Major adverse cardiac event-free survival rate




Number at risk

D < 4	144	65	28	7	1
4 ≤ D < 6	279	149	73	14	1
6 ≤ D < 8	159	100	52	21	2
8 ≤ D	127	66	37	12	3

Time after diagnosis (years)

Major adverse cardiac event-free survival rate



Number at risk

D < 4	56	26	15	2	0
4 ≤ D < 6	132	67	35	15	0
6 ≤ D < 8	58	37	20	6	0
8 ≤ D	41	33	16	7	3

Time after diagnosis (years)

The survival curves were significantly different among the four groups ($P < .001$, log rank test of equality). Severity classification was based on < 4 mm, ≥ 4 to < 6 mm, ≥ 6 to < 8 mm, and ≥ 8 mm of the actual internal diameter (D) of coronary artery aneurysms.