

Supplementary Online Content

Prakash A, Gordon LB, Kleinman ME, et al. Cardiac abnormalities in patients with Hutchinson-Gilford progeria syndrome. *JAMA Cardiol*. Published online February 21, 2018.
doi:10.1001/jamacardio.2017.5235

eTable. Lipid Profile and Glucose Metabolism

This supplementary material has been provided by the authors to give readers additional information about their work.

eTable. Lipid Profile and Glucose Metabolism

Parameter	Median (range)	Fraction (%) abnormal	Normal Range
Total cholesterol (mg/dL)	155 (94, 191)	Borderline 7/27 (26%) High 0/27 (0%)	Desirable <170 High >200
LDL cholesterol (mg/dL)	97 (40, 133)	Borderline 6/27 (22%) High 1/27 (4%)	Desirable <110 High ≥130
HDL cholesterol (mg/dL)	32 (15, 87)	Low 17/27 (63%)	Desirable 36-109
Triglycerides (mg/dL)	73 (40, 406)	Borderline 4/27 (15%) High 2/27 (7%)	Desirable <150 High ≥200
Fasting glucose (mg/dL)	84 (41, 102)	1/27 (4%)	Normal <100
Fasting insulin (μU/mL)	5.1 (0.2, 162.4)	13/27 (48%)	Normal <5
HOMA-IR	1.1 (0.02, 35.7)	9/27 (33%)	Normal <2.5

Data are presented as median (range) or frequency (%). HDL indicates high density lipoprotein; HOMA-IR, homeostasis model assessment-insulin resistance